



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 11:07 AM UTC

PDB ID : 9C39 / pdb_00009c39
EMDB ID : EMD-45162
Title : Bacteriophage Sf14 neck C6 reconstruction
Authors : Subramanian, S.; Kerns, H.R.; Braverman, S.G.; Doore, S.M.
Deposited on : 2024-05-31
Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.















Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14717 (2.90 - 3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	<p>14% 60% 24% 14%</p>
1	B	488	<p>13% 61% 24% 14%</p>
2	C	149	<p>66% 21% 13%</p>
2	D	149	<p>64% 22% 13%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	133	 82% 16% .
4	F	199	 76% 21% ..
5	G	148	 77% 19% .
6	H	450	 77% 20% ..
7	I	166	 5% 73% 21% 6%
7	J	166	 60% 22% . 16%
8	K	110	 7% 68% 30% .
8	L	110	 5% 79% 19% .
8	M	110	 75% 22% ..
8	N	110	 5% 75% 24% .
8	O	110	 5% 68% 28% ..
8	P	110	 16% 73% 24% ..

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 22110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Portal protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	421	Total	C	N	O	S	0	0
			3202	2044	531	610	17		
1	B	421	Total	C	N	O	S	0	0
			3175	2031	527	600	17		

- Molecule 2 is a protein called Phage protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	130	Total	C	N	O	S	0	0
			1018	639	178	198	3		
2	D	129	Total	C	N	O	S	0	0
			1011	639	175	194	3		

- Molecule 3 is a protein called Head-closure protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	130	Total	C	N	O	S	0	0
			1035	651	187	193	4		

- Molecule 4 is a protein called gp119.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	F	195	Total	C	N	O	S	0	0
			1525	964	258	299	4		

- Molecule 5 is a protein called gp40.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	G	142	Total	C	N	O	S	0	0
			1087	682	179	221	5		

- Molecule 6 is a protein called Putative structural protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	H	442	Total	C	N	O	S	0	0
			3212	2032	558	617	5		

- Molecule 7 is a protein called gp127.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	I	156	Total	C	N	O	S	0	0
			1153	733	206	209	5		
7	J	139	Total	C	N	O	S	0	0
			1016	641	179	191	5		


- Molecule 8 is a protein called Phage protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	K	108	Total	C	N	O	S	0	0
			797	506	127	163	1		
8	L	108	Total	C	N	O	S	0	0
			790	503	126	160	1		
8	M	108	Total	C	N	O	S	0	0
			782	499	127	155	1		
8	N	108	Total	C	N	O	S	0	0
			789	502	127	159	1		
8	O	108	Total	C	N	O	S	0	0
			770	492	125	152	1		
8	P	108	Total	C	N	O	S	0	0
			748	473	125	149	1		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Portal protein





- Molecule 1: Portal protein






• Molecule 2: Phage protein





• Molecule 2: Phage protein



• Molecule 3: Head-closure protein





• Molecule 4: gp119








• Molecule 5: gp40




• Molecule 6: Putative structural protein



• Molecule 7: gp127




• Molecule 7: gp127








• Molecule 8: Phage protein





• Molecule 8: Phage protein




• Molecule 8: Phage protein





• Molecule 8: Phage protein



• Molecule 8: Phage protein



• Molecule 8: Phage protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C6	Depositor
Number of particles used	37006	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	33	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	1.203	Depositor
Minimum map value	-0.529	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.063	Depositor
Recommended contour level	0.242	Depositor
Map size (\AA)	489.59998, 489.59998, 489.59998	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.816, 0.816, 0.816	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3272	0.58	1/4439 (0.0%)
1	B	0.31	0/3246	0.60	0/4408
2	C	0.17	0/1034	0.49	0/1394
2	D	0.19	0/1028	0.55	0/1386
3	E	0.21	0/1055	0.56	1/1429 (0.1%)
4	F	0.18	0/1552	0.42	0/2114
5	G	0.19	0/1106	0.42	0/1505
6	H	0.21	0/3268	0.48	0/4460
7	I	0.19	0/1170	0.44	0/1584
7	J	0.30	0/1029	0.63	0/1394
8	K	0.19	0/813	0.49	0/1119
8	L	0.20	0/806	0.51	0/1110
8	M	0.19	0/798	0.48	0/1100
8	N	0.20	0/805	0.51	0/1109
8	O	0.31	0/786	0.63	2/1086 (0.2%)
8	P	0.42	0/763	0.59	0/1056
All	All	0.24	0/22531	0.53	4/30693 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
4	F	0	1
5	G	0	1
7	J	0	3
All	All	0	6

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	O	46	SER	CA-C-N	7.66	130.93	120.67
8	O	46	SER	C-N-CA	7.66	130.93	120.67
1	A	454	PRO	N-CA-CB	-5.50	96.95	103.00
3	E	99	VAL	N-CA-C	5.31	111.90	106.21

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	377	ARG	Sidechain
4	F	52	GLN	Peptide
5	G	123	ARG	Sidechain
7	J	33	ARG	Sidechain
7	J	90	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3202	0	3000	101	0
1	B	3175	0	2959	96	0
2	C	1018	0	997	27	0
2	D	1011	0	996	36	0
3	E	1035	0	997	16	0
4	F	1525	0	1507	33	0
5	G	1087	0	1062	22	0
6	H	3212	0	3060	62	0
7	I	1153	0	1114	26	0
7	J	1016	0	946	24	0
8	K	797	0	784	27	0
8	L	790	0	776	17	0
8	M	782	0	770	22	0
8	N	789	0	776	23	0
8	O	770	0	743	25	0
8	P	748	0	689	20	0
All	All	22110	0	21176	491	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 491 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:P:48:PRO:HB3	8:P:53:ILE:HG12	1.47	0.95
1:B:129:LYS:HA	1:B:146:ILE:HD12	1.52	0.90
1:B:101:MET:SD	1:B:103:HIS:CE1	2.68	0.86
8:P:9:LYS:HZ3	8:P:109:ASN:HD21	1.27	0.81
1:B:101:MET:SD	1:B:103:HIS:HE1	2.03	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/488 (86%)	373 (89%)	40 (10%)	4 (1%)	12	40
1	B	417/488 (86%)	371 (89%)	40 (10%)	6 (1%)	9	31
2	C	128/149 (86%)	122 (95%)	5 (4%)	1 (1%)	16	44
2	D	127/149 (85%)	116 (91%)	11 (9%)	0	100	100
3	E	128/133 (96%)	120 (94%)	7 (6%)	1 (1%)	16	44
4	F	193/199 (97%)	172 (89%)	20 (10%)	1 (0%)	24	54
5	G	140/148 (95%)	127 (91%)	13 (9%)	0	100	100
6	H	438/450 (97%)	380 (87%)	49 (11%)	9 (2%)	5	24
7	I	154/166 (93%)	140 (91%)	13 (8%)	1 (1%)	21	50
7	J	135/166 (81%)	114 (84%)	17 (13%)	4 (3%)	3	18
8	K	106/110 (96%)	98 (92%)	7 (7%)	1 (1%)	14	42
8	L	106/110 (96%)	100 (94%)	6 (6%)	0	100	100
8	M	106/110 (96%)	101 (95%)	5 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
8	N	106/110 (96%)	99 (93%)	7 (7%)	0	100	100
8	O	106/110 (96%)	97 (92%)	8 (8%)	1 (1%)	14	42
8	P	106/110 (96%)	96 (91%)	9 (8%)	1 (1%)	14	42
All	All	2913/3196 (91%)	2626 (90%)	257 (9%)	30 (1%)	15	40

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	PRO
1	A	388	MET
1	B	152	PRO
2	C	106	CYS
4	F	53	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/424 (76%)	312 (97%)	9 (3%)	38	60
1	B	314/424 (74%)	305 (97%)	9 (3%)	37	60
2	C	107/130 (82%)	107 (100%)	0	100	100
2	D	106/130 (82%)	106 (100%)	0	100	100
3	E	109/121 (90%)	109 (100%)	0	100	100
4	F	168/176 (96%)	168 (100%)	0	100	100
5	G	121/127 (95%)	120 (99%)	1 (1%)	73	77
6	H	307/368 (83%)	305 (99%)	2 (1%)	76	78
7	I	117/150 (78%)	117 (100%)	0	100	100
7	J	100/150 (67%)	98 (98%)	2 (2%)	48	65
8	K	92/95 (97%)	91 (99%)	1 (1%)	65	74
8	L	90/95 (95%)	89 (99%)	1 (1%)	65	74
8	M	88/95 (93%)	87 (99%)	1 (1%)	65	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
8	N	90/95 (95%)	90 (100%)	0	100	100
8	O	84/95 (88%)	82 (98%)	2 (2%)	43	62
8	P	76/95 (80%)	75 (99%)	1 (1%)	61	71
All	All	2290/2770 (83%)	2261 (99%)	29 (1%)	59	71

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	145	LEU
8	O	6	ILE
1	B	452	LEU
8	L	59	VAL
1	B	394	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
8	K	35	ASN
8	P	109	ASN
8	M	60	ASN
2	D	105	GLN
7	J	154	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-45162. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 300




Y Index: 300




Z Index: 300


6.2.2 Raw map



X Index: 300



Y Index: 300




Z Index: 300


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 271




Y Index: 271




Z Index: 429


6.3.2 Raw map



X Index: 271



Y Index: 270




Z Index: 169


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.242. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_45162_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 1189 nm³; this corresponds to an approximate mass of 1074 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.36	3.87	3.42
Unmasked-calculated*	4.28	7.02	4.36

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.28 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-45162 and PDB model 9C39. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.242 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.242).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.242) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6940	 0.4160
A	 0.6020	 0.4080
B	 0.5980	 0.3910
C	 0.6780	 0.4410
D	 0.6820	 0.4360
E	 0.7340	 0.4560
F	 0.7560	 0.4510
G	 0.7580	 0.4520
H	 0.7700	 0.4220
I	 0.7420	 0.4190
J	 0.7290	 0.4380
K	 0.7170	 0.3960
L	 0.7070	 0.4110
M	 0.7370	 0.4220
N	 0.7340	 0.4120
O	 0.7240	 0.3700
P	 0.6550	 0.3350

