



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:38 PM UTC

PDB ID : 9C3X / pdb_00009c3x
Title : Crystal structure of biphenyl synthase from *Malus domestica* complexed with triketide-CoA mimetic
Authors : Re, R.N.; Noel, J.P.; Burkart, M.D.
Deposited on : 2024-06-02
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	% 93%
1	B	390	2% 92% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13315 atoms, of which 6200 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BIS3 biphenyl synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	379	Total	C	H	N	O	S	0	13	0
			6076	1927	3052	519	560	18			
1	B	378	Total	C	H	N	O	S	0	11	0
			6051	1919	3042	517	555	18			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP K9MST3
A	0	SER	-	expression tag	UNP K9MST3
A	251	VAL	ALA	engineered mutation	UNP K9MST3
B	-1	GLY	-	expression tag	UNP K9MST3
B	0	SER	-	expression tag	UNP K9MST3
B	251	VAL	ALA	engineered mutation	UNP K9MST3

- Molecule 2 is (3R)-butane-1,3-diol (CCD ID: BU4) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			16	4	10	2		
2	B	1	Total	C	H	O	0	0
			16	4	10	2		

- Molecule 3 is [(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-4-hydroxy-3-(phosphonoxy)oxolan-2-yl]methyl (3R)-3-hydroxy-2,2-dimethyl-4-oxo-4-[(3-oxo-3-{[2-(5-oxo-5-phenylpentanamido)ethyl]amino}propyl)amino]butyl dihydrogen diphosphate (CCD ID: A1AUA) (formula: $C_{32}H_{47}N_8O_{18}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 104	C 32	H 43	N 8	O 18	P 3	0	0
3	B	1	Total 104	C 32	H 43	N 8	O 18	P 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	486	Total	O	0	0
			486	486		
4	B	462	Total	O	0	0
			462	462		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.29Å 113.25Å 63.02Å 90.00° 93.35° 90.00°	Depositor
Resolution (Å)	55.20 – 1.45 55.20 – 1.45	Depositor EDS
% Data completeness (in resolution range)	98.5 (55.20-1.45) 93.0 (55.20-1.45)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 1.45Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.142 , 0.164 0.142 , 0.163	Depositor DCC
R_{free} test set	6818 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	8.4	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13315	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: A1AUA, CSD, BU4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3110	0.60	0/4207
1	B	0.37	0/3083	0.60	0/4170
All	All	0.38	0/6193	0.60	0/8377

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3024	3052	3070	10	0
1	B	3009	3042	3055	14	0
2	A	6	10	10	0	0
2	B	6	10	10	0	0
3	A	61	43	0	1	0
3	B	61	43	0	1	0
4	A	486	0	0	3	2
4	B	462	0	0	5	2
All	All	7115	6200	6145	24	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71[B]:MET:HE3	1:A:92:ILE:HD13	1.64	0.79
1:B:225:PRO:O	1:B:227:ILE:HD12	1.90	0.72
1:B:71[B]:MET:HE3	1:B:92:ILE:HD13	1.74	0.69
1:B:262:LEU:HD21	3:B:401:A1AUA:O04	1.94	0.67
1:A:283:GLU:OE2	4:A:501:HOH:O	2.14	0.64
1:B:296:PHE:CD1	1:B:346[B]:MET:HE1	2.36	0.60
1:B:25[B]:VAL:HG22	1:B:64:TYR:CE1	2.40	0.57
1:A:25[B]:VAL:HG22	1:A:64:TYR:CE1	2.42	0.55
1:A:262:LEU:HD21	3:A:401:A1AUA:O04	2.07	0.54
1:B:285[A]:VAL:HG11	1:B:364:TRP:CZ3	2.45	0.52
1:B:77:ASN:HB2	4:B:821:HOH:O	2.14	0.46
1:A:67:LEU:HD12	1:A:71[B]:MET:HE2	1.97	0.46
1:B:10:GLN:HG2	1:B:10:GLN:O	2.15	0.46
1:B:42:GLU:HG3	4:B:810:HOH:O	2.15	0.46
1:B:289:LYS:HE3	4:B:681:HOH:O	2.15	0.45
1:A:325:ARG:NH1	4:A:511:HOH:O	2.50	0.44
1:B:67:LEU:HD12	1:B:71[B]:MET:HE2	2.00	0.43
1:B:253:ILE:HD12	1:B:253:ILE:N	2.34	0.43
1:A:34:PHE:O	1:A:38:VAL:HG22	2.19	0.42
1:A:71[B]:MET:CE	1:A:92:ILE:HD13	2.43	0.41
1:A:183:VAL:O	1:A:216:ALA:HA	2.21	0.41
1:B:254:ARG:HD2	4:B:539:HOH:O	2.20	0.41
1:A:246[A]:GLU:HG3	4:A:836:HOH:O	2.21	0.40
1:B:268:LYS:NZ	4:B:519:HOH:O	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:807:HOH:O	4:B:860:HOH:O[1_455]	2.05	0.15
4:A:876:HOH:O	4:B:748:HOH:O[1_455]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/390 (100%)	380 (98%)	8 (2%)	0	100	100
1	B	385/390 (99%)	379 (98%)	6 (2%)	0	100	100
All	All	773/780 (99%)	759 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/329 (101%)	331 (100%)	1 (0%)	86	74
1	B	329/329 (100%)	329 (100%)	0	100	100
All	All	661/658 (100%)	660 (100%)	1 (0%)	87	76

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	202	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	27	HIS
1	A	75	ASN
1	B	222	ASN
1	B	252	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	B	159[A]	1	4,7,8	1.83	1 (25%)	1,8,10	0.01	0
1	CSD	A	159[A]	1	4,7,8	1.99	1 (25%)	1,8,10	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	B	159[A]	1	-	0/2/6/8	-
1	CSD	A	159[A]	1	-	0/2/6/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159[A]	CSD	OD1-SG	-3.73	1.44	1.47
1	B	159[A]	CSD	OD1-SG	-3.37	1.44	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	A1AUA	A	401	-	62,64,64	2.84	24 (38%)	88,93,93	1.64	13 (14%)
2	BU4	B	400	-	5,5,5	0.17	0	5,5,5	0.60	0
2	BU4	A	400	-	5,5,5	0.28	0	5,5,5	0.49	0
3	A1AUA	B	401	-	62,64,64	2.85	25 (40%)	88,93,93	1.59	13 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1AUA	A	401	-	-	6/61/77/77	0/4/4/4
2	BU4	B	400	-	-	0/3/3/3	-
2	BU4	A	400	-	-	1/3/3/3	-
3	A1AUA	B	401	-	-	7/61/77/77	0/4/4/4

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	A1AUA	C14-C13	9.13	1.53	1.39
3	B	401	A1AUA	C14-C13	8.42	1.52	1.39
3	B	401	A1AUA	C17-C18	8.05	1.52	1.38
3	A	401	A1AUA	C17-C18	7.63	1.52	1.38
3	B	401	A1AUA	C16-C15	6.34	1.52	1.38
3	A	401	A1AUA	C16-C15	5.99	1.51	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	A1AUA	P01-O07	5.40	1.65	1.59
3	B	401	A1AUA	P01-O07	5.04	1.64	1.59
3	B	401	A1AUA	P02-O07	5.02	1.64	1.59
3	A	401	A1AUA	C08-N01	4.78	1.44	1.33
3	A	401	A1AUA	C18-C13	-4.71	1.32	1.39
3	A	401	A1AUA	C06-N03	4.66	1.44	1.33
3	B	401	A1AUA	O13-C25	4.53	1.52	1.42
3	B	401	A1AUA	C03-N02	4.45	1.43	1.33
3	B	401	A1AUA	C08-N01	4.45	1.43	1.33
3	A	401	A1AUA	O13-C25	4.44	1.52	1.42
3	B	401	A1AUA	P03-O15	4.36	1.67	1.59
3	A	401	A1AUA	P02-O07	4.33	1.64	1.59
3	B	401	A1AUA	C31-N08	4.30	1.45	1.34
3	B	401	A1AUA	C18-C13	-4.24	1.33	1.39
3	A	401	A1AUA	C29-N05	4.22	1.46	1.39
3	B	401	A1AUA	C06-N03	4.21	1.43	1.33
3	A	401	A1AUA	C31-N08	4.09	1.44	1.34
3	A	401	A1AUA	C03-N02	3.95	1.42	1.33
3	B	401	A1AUA	C29-N05	3.90	1.46	1.39
3	A	401	A1AUA	C15-C14	-3.88	1.32	1.38
3	A	401	A1AUA	P03-O15	3.66	1.65	1.59
3	B	401	A1AUA	C15-C14	-3.61	1.32	1.38
3	A	401	A1AUA	C28-N05	3.41	1.38	1.31
3	B	401	A1AUA	C28-N05	3.38	1.38	1.31
3	A	401	A1AUA	O13-C24	3.29	1.52	1.45
3	A	401	A1AUA	C26-C27	-3.27	1.45	1.53
3	B	401	A1AUA	O13-C24	3.14	1.52	1.45
3	B	401	A1AUA	C26-C27	-3.07	1.46	1.53
3	A	401	A1AUA	C16-C17	-2.64	1.32	1.38
3	B	401	A1AUA	C16-C17	-2.56	1.32	1.38
3	A	401	A1AUA	C29-C31	-2.49	1.34	1.41
3	B	401	A1AUA	C28-N04	-2.49	1.33	1.37
3	B	401	A1AUA	C13-C12	2.38	1.53	1.49
3	B	401	A1AUA	C29-C31	-2.38	1.34	1.41
3	A	401	A1AUA	C13-C12	2.31	1.52	1.49
3	B	401	A1AUA	C30-N04	-2.29	1.32	1.37
3	B	401	A1AUA	C32-N07	2.18	1.37	1.33
3	A	401	A1AUA	O04-C08	-2.14	1.19	1.23
3	A	401	A1AUA	C30-N04	-2.13	1.33	1.37
3	A	401	A1AUA	C32-N06	2.12	1.37	1.33
3	A	401	A1AUA	O03-C03	-2.06	1.19	1.23
3	B	401	A1AUA	O05-C12	-2.00	1.18	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	A1AUA	O04-C08	-2.00	1.19	1.23

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	A1AUA	C30-N04-C28	7.94	114.07	105.74
3	B	401	A1AUA	C30-N04-C28	7.64	113.76	105.74
3	A	401	A1AUA	N07-C32-N06	-5.65	120.03	128.58
3	B	401	A1AUA	N07-C32-N06	-5.37	120.46	128.58
3	A	401	A1AUA	N04-C28-N05	-4.61	107.39	113.94
3	B	401	A1AUA	N04-C28-N05	-4.24	107.92	113.94
3	A	401	A1AUA	C29-C30-N07	-3.69	121.64	126.72
3	B	401	A1AUA	C29-C30-N07	-3.46	121.95	126.72
3	B	401	A1AUA	C01-C02-N02	-3.19	101.60	111.54
3	A	401	A1AUA	C01-C02-N02	-3.19	101.62	111.54
3	A	401	A1AUA	N07-C30-N04	3.15	132.53	127.17
3	B	401	A1AUA	N07-C30-N04	2.81	131.95	127.17
3	B	401	A1AUA	C22-C19-C07	2.77	113.49	108.77
3	A	401	A1AUA	C26-C27-C24	2.77	108.08	103.24
3	B	401	A1AUA	C30-C29-N05	-2.69	107.51	110.58
3	A	401	A1AUA	C32-N07-C30	2.49	117.91	111.83
3	B	401	A1AUA	C26-C27-C24	2.40	107.45	103.24
3	A	401	A1AUA	C24-O13-C25	-2.38	104.21	109.47
3	A	401	A1AUA	C30-C29-N05	-2.34	107.91	110.58
3	B	401	A1AUA	C04-C05-N03	-2.33	107.04	112.00
3	A	401	A1AUA	C09-C08-N01	2.33	120.58	116.34
3	B	401	A1AUA	C32-N07-C30	2.25	117.32	111.83
3	B	401	A1AUA	C31-C29-C30	2.25	120.24	117.18
3	B	401	A1AUA	C24-O13-C25	-2.21	104.58	109.47
3	A	401	A1AUA	C01-N01-C08	-2.19	118.74	122.82
3	A	401	A1AUA	C31-C29-C30	2.15	120.11	117.18

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	A1AUA	C11-C12-C13-C18
3	A	401	A1AUA	O05-C12-C13-C18
3	A	401	A1AUA	C11-C12-C13-C14
3	A	401	A1AUA	O05-C12-C13-C14
3	B	401	A1AUA	C10-C11-C12-O05
3	B	401	A1AUA	C10-C11-C12-C13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	400	BU4	O1-C1-C2-C3
3	B	401	A1AUA	C11-C12-C13-C14
3	A	401	A1AUA	C19-C20-O06-P01
3	B	401	A1AUA	C19-C20-O06-P01
3	B	401	A1AUA	C03-C04-C05-N03
3	B	401	A1AUA	O05-C12-C13-C14
3	A	401	A1AUA	P02-O07-P01-O09
3	B	401	A1AUA	P02-O07-P01-O09


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	A1AUA	1	0
3	B	401	A1AUA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1AUA A 401





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/390 (96%)	-0.58	2 (0%) 87 90	5, 11, 27, 76	12 (3%)
1	B	377/390 (96%)	-0.49	7 (1%) 66 68	5, 11, 32, 59	10 (2%)
All	All	755/780 (96%)	-0.54	9 (1%) 76 80	5, 11, 30, 76	22 (2%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	388	CYS	6.7
1	B	10	GLN	4.3
1	A	387	ARG	3.6
1	B	227	ILE	3.5
1	B	225	PRO	3.2
1	B	387	ARG	3.0
1	B	222	ASN	2.6
1	B	226	GLU	2.3
1	B	46	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	159[A]	8/9	0.98	0.05	9,10,13,22	0
1	CSD	B	159[A]	8/9	0.99	0.04	8,10,13,22	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BU4	A	400	6/6	0.92	0.09	13,19,22,24	0
3	A1AUA	A	401	61/61	0.94	0.08	13,20,33,39	0
3	A1AUA	B	401	61/61	0.95	0.07	12,20,29,33	0
2	BU4	B	400	6/6	0.96	0.07	12,20,24,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around A1AUA B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.