



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:19 AM UTC

PDB ID : 1C4K / pdb_00001c4k
Title : ORNITHINE DECARBOXYLASE MUTANT (GLY121TYR)
Authors : Vitali, J.; Hackert, M.L.
Deposited on : 1999-08-26
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	730	2% 78% 18% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (ORNITHINE DECARBOXYLASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	728	5826	3723	978	1106	19	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	121	TYR	GLY	engineered mutation	UNP P43099

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	14	8	1	4	1	0	0

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	250	Total	O	0	0
			250	250		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN (ORNITHINE DECARBOXYLASE)



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	111.80Å 111.80Å 135.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	7.00 – 2.70 7.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	94.4 (7.00-2.70) 89.4 (7.00-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 2.68Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.212 , 0.281 0.204 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	46.1	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6122	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	4/5975 (0.1%)	1.18	57/8106 (0.7%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	394	SER	C-O	7.08	1.33	1.23
1	A	395	THR	N-CA	-6.63	1.37	1.46
1	A	396	SER	N-CA	-5.34	1.40	1.46
1	A	626	MET	SD-CE	-5.15	1.66	1.79

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	270	LYS	N-CA-C	-12.29	97.76	113.12
1	A	19	ASP	N-CA-C	-11.43	99.66	112.57
1	A	374	LYS	N-CA-C	10.30	122.51	111.28
1	A	152	THR	N-CA-C	9.36	121.48	111.28
1	A	477	LYS	N-CA-C	8.70	120.76	111.28
1	A	239	VAL	N-CA-C	-8.18	94.06	106.42
1	A	20	ARG	N-CA-C	-7.62	101.85	112.45
1	A	307	GLY	N-CA-C	7.59	122.08	112.83
1	A	338	ASP	N-CA-C	7.38	120.40	111.40
1	A	584	GLU	N-CA-C	6.93	118.63	111.14
1	A	88	ALA	N-CA-C	6.90	117.38	108.34
1	A	478	TRP	N-CA-C	6.84	119.60	111.33
1	A	109	PRO	N-CA-C	6.83	124.84	113.78
1	A	648	ILE	N-CA-C	6.80	120.56	111.44
1	A	218	PHE	N-CA-C	6.63	120.47	109.46
1	A	640	ASN	N-CA-C	6.53	118.99	109.07
1	A	540	LEU	N-CA-C	6.28	119.64	109.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	305	ARG	N-CA-C	6.25	121.08	112.90
1	A	205	VAL	N-CA-C	5.99	116.65	110.36
1	A	33	VAL	N-CA-C	5.92	117.11	108.58
1	A	543	MET	N-CA-C	5.86	118.07	108.52
1	A	276	ALA	N-CA-C	5.83	118.38	111.33
1	A	347	ILE	N-CA-C	5.78	116.17	107.80
1	A	482	GLU	N-CA-C	5.73	120.11	113.18
1	A	151	GLU	N-CA-C	5.72	117.52	111.28
1	A	471	ARG	N-CA-C	5.64	118.60	109.40
1	A	339	ASP	N-CA-C	5.62	122.78	110.80
1	A	232	ALA	N-CA-C	5.58	117.37	111.28
1	A	87	ASP	N-CA-C	-5.54	96.71	107.62
1	A	17	ASP	N-CA-C	5.50	118.06	107.44
1	A	162	ASP	N-CA-C	-5.50	99.08	110.80
1	A	13	ARG	N-CA-C	5.49	117.98	111.33
1	A	298	ASN	N-CA-C	-5.49	96.33	107.41
1	A	222	ASN	N-CA-C	5.47	117.95	110.24
1	A	210	VAL	N-CA-C	5.45	116.57	108.23
1	A	135	TYR	N-CA-C	-5.45	104.99	111.69
1	A	63	LYS	N-CA-C	-5.41	106.35	113.12
1	A	80	ILE	N-CA-C	-5.39	99.20	106.85
1	A	289	LEU	N-CA-C	5.36	117.81	111.33
1	A	645	LEU	N-CA-C	5.36	118.61	111.75
1	A	171	HIS	N-CA-C	5.32	122.14	110.80
1	A	297	TYR	N-CA-C	5.32	116.79	110.19
1	A	219	ASP	N-CA-C	-5.32	100.64	109.46
1	A	163	VAL	N-CA-C	5.30	120.38	109.34
1	A	340	LEU	N-CA-C	5.30	118.19	109.76
1	A	413	GLU	N-CA-C	5.28	118.10	110.28
1	A	27	GLY	N-CA-C	-5.25	100.73	113.18
1	A	391	LEU	N-CA-C	-5.18	105.32	111.69
1	A	18	THR	N-CA-C	-5.16	99.81	110.80
1	A	201	ALA	N-CA-C	-5.15	105.74	112.23
1	A	636	ILE	CB-CA-C	-5.12	105.23	112.14
1	A	237	ARG	CA-C-N	5.11	125.01	119.85
1	A	237	ARG	C-N-CA	5.11	125.01	119.85
1	A	394	SER	N-CA-CB	-5.11	103.06	110.36
1	A	62	THR	N-CA-C	5.09	117.40	109.52
1	A	311	ASP	N-CA-C	-5.06	105.46	111.69
1	A	28	SER	N-CA-C	5.05	118.00	110.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5826	0	5609	59	0
2	A	14	0	6	0	0
3	A	32	0	12	0	0
4	A	250	0	0	0	0
All	All	6122	0	5627	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:479:HIS:HD2	1:A:481:TYR:H	1.40	0.70
1:A:376:GLN:HE21	1:A:378:ARG:HE	1.40	0.68
1:A:361:SER:O	1:A:362:GLN:HB2	1.98	0.64
1:A:376:GLN:NE2	1:A:378:ARG:HE	1.94	0.64
1:A:441:MET:HG2	1:A:563:GLN:HG3	1.80	0.63
1:A:632:ARG:O	1:A:636:ILE:HG12	2.02	0.59
1:A:408:ASN:O	1:A:412:GLN:HG2	2.03	0.59
1:A:646:ASN:HD22	1:A:647:LYS:HG3	1.67	0.59
1:A:592:ILE:HD12	1:A:593:ARG:H	1.68	0.58
1:A:26:VAL:HG11	1:A:45:VAL:HG13	1.85	0.58
1:A:651:GLU:HB3	1:A:716:VAL:HG11	1.87	0.57
1:A:246:ASN:HB2	1:A:247:PRO:HD2	1.88	0.55
1:A:139:HIS:HD2	1:A:141:ALA:H	1.54	0.54
1:A:180:LYS:O	1:A:184:ARG:HG2	2.08	0.54
1:A:261:GLU:HG3	1:A:309:LEU:HD11	1.90	0.53
1:A:451:ASN:N	1:A:451:ASN:HD22	2.08	0.52
1:A:237:ARG:HH12	1:A:271:VAL:HG22	1.75	0.52
1:A:296:ILE:HD12	1:A:320:VAL:HG11	1.93	0.51
1:A:18:THR:C	1:A:20:ARG:H	2.18	0.50
1:A:318:ALA:HA	1:A:352:SER:H	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:544:THR:O	1:A:547:GLU:HG2	2.13	0.48
1:A:609:PHE:CD1	1:A:609:PHE:C	2.92	0.48
1:A:657:ALA:HB1	1:A:681:PHE:CZ	2.49	0.48
1:A:473:GLU:O	1:A:479:HIS:HE1	1.96	0.48
1:A:565:LEU:HD13	1:A:575:VAL:HG22	1.95	0.47
1:A:640:ASN:HB3	1:A:654:LEU:HD12	1.95	0.47
1:A:468:ASP:HA	1:A:471:ARG:HD2	1.97	0.47
1:A:271:VAL:HG12	1:A:273:PRO:HD2	1.97	0.47
1:A:564:ARG:HA	1:A:567:GLU:HG2	1.97	0.47
1:A:354:HIS:HB3	1:A:360:PHE:O	2.16	0.46
1:A:580:TYR:C	1:A:582:ALA:H	2.24	0.46
1:A:570:ALA:H	1:A:592:ILE:HD11	1.81	0.45
1:A:270:LYS:NZ	1:A:624:GLN:HE22	2.14	0.45
1:A:311:ASP:HA	1:A:378:ARG:HH11	1.81	0.45
1:A:246:ASN:HD21	1:A:248:TYR:HB2	1.81	0.45
1:A:21:VAL:HG12	1:A:22:VAL:N	2.31	0.44
1:A:168:LEU:HD12	1:A:168:LEU:N	2.32	0.44
1:A:509:ASP:CG	1:A:510:TYR:H	2.25	0.44
1:A:358:ALA:O	1:A:408:ASN:HB2	2.18	0.44
1:A:237:ARG:HH22	1:A:271:VAL:CG2	2.30	0.44
1:A:246:ASN:ND2	1:A:248:TYR:H	2.16	0.43
1:A:226:VAL:HG11	1:A:286:VAL:HG11	2.01	0.43
1:A:544:THR:C	1:A:546:ALA:H	2.27	0.43
1:A:246:ASN:HD22	1:A:246:ASN:C	2.25	0.43
1:A:569:ASP:HA	1:A:592:ILE:HD11	2.01	0.43
1:A:710:VAL:HG13	1:A:710:VAL:O	2.19	0.42
1:A:685:GLN:NE2	1:A:711:VAL:HA	2.34	0.42
1:A:272:ASP:OD1	1:A:272:ASP:N	2.52	0.42
1:A:353:VAL:O	1:A:357:GLN:HB2	2.19	0.42
1:A:169:LEU:HD21	1:A:395:THR:HA	2.00	0.42
1:A:194:VAL:HG12	1:A:196:GLY:H	1.83	0.41
1:A:4:LEU:O	1:A:21:VAL:HG11	2.19	0.41
1:A:473:GLU:O	1:A:479:HIS:CE1	2.74	0.41
1:A:543:MET:HE1	1:A:552:MET:SD	2.61	0.41
1:A:425:LEU:HD23	1:A:425:LEU:HA	1.91	0.41
1:A:447:PRO:HA	1:A:448:PRO:HD3	1.92	0.41
1:A:685:GLN:HE21	1:A:711:VAL:HA	1.86	0.40
1:A:688:ILE:HG12	1:A:695:ALA:HB1	2.04	0.40
1:A:176:VAL:HG22	1:A:180:LYS:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	724/730 (99%)	681 (94%)	35 (5%)	8 (1%)	11	29

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	339	ASP
1	A	27	GLY
1	A	131	GLN
1	A	171	HIS
1	A	729	ASN
1	A	268	ALA
1	A	320	VAL
1	A	545	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	616/618 (100%)	580 (94%)	36 (6%)	18	42

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ARG
1	A	17	ASP
1	A	18	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	20	ARG
1	A	39	MET
1	A	86	PHE
1	A	106	ILE
1	A	116	GLU
1	A	152	THR
1	A	202	ASN
1	A	211	SER
1	A	245	ARG
1	A	246	ASN
1	A	262	LYS
1	A	263	LYS
1	A	267	LEU
1	A	374	LYS
1	A	376	GLN
1	A	395	THR
1	A	449	VAL
1	A	451	ASN
1	A	465	ASN
1	A	477	LYS
1	A	482	GLU
1	A	588	ASN
1	A	592	ILE
1	A	609	PHE
1	A	617	LEU
1	A	626	MET
1	A	640	ASN
1	A	646	ASN
1	A	648	ILE
1	A	658	LEU
1	A	701	VAL
1	A	716	VAL
1	A	726	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	ASN
1	A	139	HIS
1	A	187	ASN
1	A	202	ASN
1	A	222	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	246	ASN
1	A	351	GLN
1	A	354	HIS
1	A	357	GLN
1	A	362	GLN
1	A	376	GLN
1	A	451	ASN
1	A	479	HIS
1	A	488	GLN
1	A	504	ASN
1	A	522	ASN
1	A	554	ASN
1	A	583	ASN
1	A	588	ASN
1	A	606	ASN
1	A	624	GLN
1	A	640	ASN
1	A	646	ASN
1	A	685	GLN
1	A	724	ASN
1	A	730	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	A	955	1	14,14,16	2.18	6 (42%)	19,20,23	2.43	8 (42%)
3	GTP	A	999	-	33,34,34	1.20	4 (12%)	50,54,54	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	955	1	-	1/6/6/8	0/1/1/1
3	GTP	A	999	-	-	2/22/38/38	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	955	PLP	C4A-C4	4.44	1.59	1.51
2	A	955	PLP	P-O4P	-3.41	1.49	1.60
2	A	955	PLP	C2-N1	3.20	1.40	1.34
3	A	999	GTP	PB-O3A	3.19	1.62	1.59
3	A	999	GTP	PB-O3B	2.69	1.62	1.59
2	A	955	PLP	P-O2P	-2.67	1.44	1.54
3	A	999	GTP	PG-O3G	2.64	1.64	1.54
3	A	999	GTP	PA-O3A	2.40	1.62	1.59
2	A	955	PLP	P-O1P	-2.23	1.43	1.50
2	A	955	PLP	P-O3P	-2.20	1.46	1.54

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	955	PLP	O2P-P-O4P	7.63	126.57	106.67
2	A	955	PLP	C6-C5-C4	3.25	120.73	117.97
2	A	955	PLP	O3P-P-O4P	-2.79	99.40	106.67
2	A	955	PLP	C2A-C2-N1	2.52	121.42	117.68
2	A	955	PLP	C5-C6-N1	-2.39	119.94	123.83
2	A	955	PLP	O4P-C5A-C5	-2.23	105.18	109.36
2	A	955	PLP	C6-N1-C2	2.19	120.11	117.53
2	A	955	PLP	C3-C2-N1	-2.16	119.50	121.61

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	999	GTP	PB-O3B-PG-O2G
2	A	955	PLP	C5A-O4P-P-O1P
3	A	999	GTP	PB-O3B-PG-O1G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	728/730 (99%)	-0.42	12 (1%) 70 68	10, 34, 87, 100	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	ALA	6.2
1	A	19	ASP	3.4
1	A	27	GLY	3.1
1	A	269	ALA	3.0
1	A	277	LYS	2.8
1	A	18	THR	2.7
1	A	646	ASN	2.6
1	A	160	ASN	2.6
1	A	730	ASN	2.6
1	A	71	ASP	2.4
1	A	116	GLU	2.1
1	A	44	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GTP	A	999	32/32	0.96	0.07	36,36,36,36	0
2	PLP	A	955	14/16	0.97	0.06	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.