



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 12:00 PM UTC

PDB ID : 5C4A / pdb_00005c4a
Title : Crystal structure of a transcribing RNA Polymerase II complex reveals a complete transcription bubble
Authors : Barnes, C.O.; Calero, M.; Malik, I.; Saphr, H.; Zhang, Q.; Pullara, F.; Kaplan, C.D.; Calero, G.
Deposited on : 2015-06-17
Resolution : 4.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1004 (4.52-3.88)
Clashscore	190562	1019 (4.50-3.90)
Ramachandran outliers	187476	1020 (4.56-3.84)
Sidechain outliers	187428	1006 (4.56-3.84)
RSRZ outliers	180081	1001 (4.52-3.88)
RNA backbone	3983	1026 (5.04-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	5% 62% 17% • 18%
2	B	1224	11% 70% 20% • 5%
3	C	318	6% 66% 15% • 17%
4	D	221	21% 63% 13% 24%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	215	
6	F	155	
7	G	179	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	R	9	
14	S	56	
15	U	56	

2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 32574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1425	Total	C	N	O	S	0	0	0
			11206	7057	1960	2127	62			

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1162	Total	C	N	O	S	0	0	0
			9215	5816	1617	1726	56			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	265	Total	C	N	O	S	0	0	0
			2086	1312	347	414	13			

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	168	Total	C	N	O	S	0	0	0
			1331	822	237	270	2			

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	87	Total	C	N	O	S	0	0	0
			705	451	119	132	3			

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	170	Total	C	N	O	S	0	0	0
			1331	857	220	246	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	172	LEU	-	expression tag	UNP P34087
G	173	GLU	-	expression tag	UNP P34087
G	174	HIS	-	expression tag	UNP P34087
G	175	HIS	-	expression tag	UNP P34087
G	176	HIS	-	expression tag	UNP P34087
G	177	HIS	-	expression tag	UNP P34087
G	178	HIS	-	expression tag	UNP P34087
G	179	HIS	-	expression tag	UNP P34087

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	135	Total	C	N	O	S	0	0	0
			1080	679	182	214	5			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	114	Total	C	N	O	S	0	0	0
			927	571	168	178	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	66	Total	C	N	O	S	0	0	0
			540	345	94	95	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	115	Total	C	N	O	S	0	0	0
			924	593	157	172	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	44	Total	C	N	O	S	0	0	0
			352	217	70	61	4			

- Molecule 13 is a RNA chain called RNA (5'-R(P*UP*CP*GP*AP*GP*AP*GP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	R	9	Total	C	N	O	P	0	0	0
			197	88	40	60	9			

- Molecule 14 is a DNA chain called Scaffold 2 Non-template Strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	S	16	Total	C	N	O	P	0	0	0
			331	158	58	99	16			

- Molecule 15 is a DNA chain called Scaffold 2 Template Strand.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	U	29	Total	C	N	O	P	0	0	0
			587	280	107	171	29			

- Molecule 16 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	2	Total	Zn	0	0
			2	2		
16	B	1	Total	Zn	0	0
			1	1		
16	C	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		
16	J	1	Total	Zn	0	0
			1	1		
16	L	1	Total	Zn	0	0
			1	1		


- Molecule 17 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	2	Total	Mg	0	0
			2	2		




• Molecule 2: DNA-directed RNA polymerase II subunit RPB2







- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2




- Molecule 7: DNA-directed RNA polymerase II subunit RPB7



- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 9: DNA-directed RNA polymerase II subunit RPB9





- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5






- Molecule 11: DNA-directed RNA polymerase II subunit RPB11




- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4





- Molecule 13: RNA (5'-R(P*UP*CP*GP*AP*GP*AP*GP*GP*A)-3')



- Molecule 14: Scaffold 2 Non-template Strand



- Molecule 15: Scaffold 2 Template Strand



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	219.83Å 396.71Å 273.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.91 – 4.20 39.91 – 4.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.91-4.20) 99.6 (39.91-4.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 4.13Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.224 , 0.242 0.245 , 0.263	Depositor DCC
R_{free} test set	6719 reflections (7.74%)	wwPDB-VP
Wilson B-factor (Å ²)	100.3	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 134.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.019 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.026 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	32574	wwPDB-VP
Average B, all atoms (Å ²)	139.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	2/11407 (0.0%)	1.25	17/15428 (0.1%)
2	B	0.96	4/9390 (0.0%)	1.22	27/12662 (0.2%)
3	C	0.91	0/2124	1.18	1/2879 (0.0%)
4	D	0.89	0/1339	1.25	0/1793
5	E	0.92	1/1788 (0.1%)	1.17	2/2406 (0.1%)
6	F	0.92	0/717	1.17	0/967
7	G	0.93	0/1358	1.19	5/1830 (0.3%)
8	H	0.87	0/1097	1.09	1/1484 (0.1%)
9	I	0.88	0/945	1.13	0/1273
10	J	0.87	0/549	1.17	1/738 (0.1%)
11	K	0.86	0/942	1.19	1/1272 (0.1%)
12	L	0.93	0/354	1.14	0/468
13	R	0.51	0/221	0.62	0/343
14	S	0.40	0/370	0.65	0/570
15	U	0.43	0/657	0.57	0/1009
All	All	0.92	7/33258 (0.0%)	1.19	55/45122 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	264	SER	C-N	15.25	1.51	1.33
2	B	924	GLU	CA-C	11.95	1.58	1.52
1	A	417	TYR	CA-C	7.57	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	265	SER	C-N	6.13	1.42	1.33
5	E	124	VAL	CA-CB	5.53	1.57	1.54

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	934	LYS	N-CA-C	7.72	119.39	110.97
2	B	1139	ILE	N-CA-C	-7.57	103.47	110.82
1	A	864	ILE	N-CA-C	-6.83	105.18	111.67
2	B	129	PHE	CA-CB-CG	6.49	120.29	113.80
2	B	918	ILE	CA-C-N	6.48	133.37	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	265	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11206	0	11265	140	0
2	B	9215	0	9210	114	0
3	C	2086	0	2045	24	0
4	D	1331	0	1345	8	0
5	E	1752	0	1776	9	0
6	F	705	0	731	8	0
7	G	1331	0	1350	20	0
8	H	1080	0	1049	10	0
9	I	927	0	883	7	0
10	J	540	0	555	9	0
11	K	924	0	934	3	0
12	L	352	0	377	2	0
13	R	197	0	96	4	0
14	S	331	0	183	2	0
15	U	587	0	326	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	C	1	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	A	2	0	0	0	0
All	All	32574	0	32125	328	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 328 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:869:GLY:HA3	1:A:1366:ARG:HG2	1.46	0.98
3:C:39:ALA:HA	3:C:164:ALA:HB3	1.64	0.80
1:A:768:GLN:HG2	1:A:816:HIS:HA	1.64	0.79
1:A:311:GLN:HB2	1:A:312:PRO:HD3	1.67	0.77
1:A:1107:VAL:HA	1:A:1108:ALA:HB3	1.68	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1417/1733 (82%)	1222 (86%)	147 (10%)	48 (3%)	3	21
2	B	1146/1224 (94%)	977 (85%)	124 (11%)	45 (4%)	2	19
3	C	263/318 (83%)	235 (89%)	20 (8%)	8 (3%)	3	23
4	D	164/221 (74%)	150 (92%)	8 (5%)	6 (4%)	2	20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	E	212/215 (99%)	197 (93%)	9 (4%)	6 (3%)	4	25
6	F	85/155 (55%)	74 (87%)	10 (12%)	1 (1%)	10	42
7	G	166/179 (93%)	147 (89%)	14 (8%)	5 (3%)	3	23
8	H	129/146 (88%)	111 (86%)	10 (8%)	8 (6%)	1	13
9	I	112/122 (92%)	100 (89%)	10 (9%)	2 (2%)	6	33
10	J	64/70 (91%)	57 (89%)	6 (9%)	1 (2%)	7	37
11	K	113/120 (94%)	109 (96%)	4 (4%)	0	100	100
12	L	42/70 (60%)	31 (74%)	9 (21%)	2 (5%)	2	17
All	All	3913/4573 (86%)	3410 (87%)	371 (10%)	132 (3%)	3	21

5 of 132 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	LYS
1	A	66	LYS
1	A	76	GLU
1	A	194	ALA
1	A	286	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1244/1520 (82%)	1116 (90%)	128 (10%)	7	23
2	B	1002/1061 (94%)	888 (89%)	114 (11%)	5	21
3	C	233/274 (85%)	217 (93%)	16 (7%)	14	37
4	D	146/200 (73%)	134 (92%)	12 (8%)	10	32
5	E	196/197 (100%)	185 (94%)	11 (6%)	19	42
6	F	77/137 (56%)	73 (95%)	4 (5%)	21	44
7	G	151/160 (94%)	138 (91%)	13 (9%)	10	30
8	H	118/128 (92%)	111 (94%)	7 (6%)	18	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
9	I	108/116 (93%)	102 (94%)	6 (6%)	19	42
10	J	61/65 (94%)	56 (92%)	5 (8%)	10	32
11	K	99/102 (97%)	90 (91%)	9 (9%)	9	29
12	L	39/57 (68%)	33 (85%)	6 (15%)	2	14
All	All	3474/4017 (86%)	3143 (90%)	331 (10%)	8	26

5 of 331 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	1138	MET
7	G	13	LEU
2	B	1182	CYS
4	D	71	LYS
8	H	40	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	481	GLN
2	B	951	GLN
11	K	76	GLN
8	H	21	ASN
2	B	499	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
13	R	7/9 (77%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1425/1733 (82%)	0.66	87 (6%) 27 27	50, 116, 208, 300	0
2	B	1162/1224 (94%)	0.90	133 (11%) 10 14	50, 131, 241, 300	0
3	C	265/318 (83%)	0.77	19 (7%) 21 23	50, 121, 191, 240	0
4	D	168/221 (76%)	1.53	47 (27%) 1 4	78, 178, 284, 300	0
5	E	214/215 (99%)	0.70	12 (5%) 30 28	65, 148, 226, 282	0
6	F	87/155 (56%)	0.49	3 (3%) 48 38	51, 94, 173, 238	0
7	G	170/179 (94%)	1.11	35 (20%) 2 6	57, 128, 270, 300	0
8	H	135/146 (92%)	1.00	19 (14%) 6 10	99, 164, 236, 292	0
9	I	114/122 (93%)	0.98	13 (11%) 10 14	78, 155, 214, 280	0
10	J	66/70 (94%)	0.61	3 (4%) 38 33	67, 121, 194, 218	0
11	K	115/120 (95%)	0.54	3 (2%) 57 43	50, 116, 179, 237	0
12	L	44/70 (62%)	1.15	7 (15%) 5 8	82, 157, 207, 241	0
13	R	9/9 (100%)	0.69	0 100 100	158, 185, 278, 293	0
14	S	16/56 (28%)	2.28	10 (62%) 0 1	111, 278, 300, 300	0
15	U	29/56 (51%)	1.82	10 (34%) 1 3	111, 262, 300, 300	0
All	All	4019/4694 (85%)	0.83	401 (9%) 12 16	50, 129, 238, 300	0

The worst 5 of 401 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	245	GLU	8.2
7	G	94	CYS	6.9
2	B	735	ALA	6.0
4	D	217	LEU	5.6
4	D	149	THR	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
17	MG	A	1804	1/1	0.62	0.28	71,71,71,71	0
17	MG	A	1803	1/1	0.96	0.07	54,54,54,54	0
16	ZN	A	1802	1/1	0.98	0.10	68,68,68,68	0
16	ZN	I	202	1/1	0.98	0.10	155,155,155,155	0
16	ZN	A	1801	1/1	0.99	0.05	101,101,101,101	0
16	ZN	J	101	1/1	0.99	0.10	90,90,90,90	0
16	ZN	L	101	1/1	0.99	0.08	91,91,91,91	0
16	ZN	B	1301	1/1	0.99	0.07	60,60,60,60	0
16	ZN	I	201	1/1	0.99	0.07	115,115,115,115	0
16	ZN	C	401	1/1	1.00	0.06	82,82,82,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.