



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:19 PM UTC

PDB ID : 6C4T / pdb_00006c4t
Title : Staphylopine dehydrogenase (SaODH) - NADP⁺ bound
Authors : McFarlane, J.S.; Davis, C.L.; Lamb, A.L.
Deposited on : 2018-01-12
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6860 atoms, of which 3329 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Staphylopin dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	428	6797	2230	3321	566	656	24	0	0	0

There are 19 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	HIS	-	expression tag	UNP A0A1Q4GXD5
A	-16	HIS	-	expression tag	UNP A0A1Q4GXD5
A	-15	HIS	-	expression tag	UNP A0A1Q4GXD5
A	-14	HIS	-	expression tag	UNP A0A1Q4GXD5
A	-13	HIS	-	expression tag	UNP A0A1Q4GXD5
A	-12	HIS	-	expression tag	UNP A0A1Q4GXD5
A	-11	SER	-	expression tag	UNP A0A1Q4GXD5
A	-10	SER	-	expression tag	UNP A0A1Q4GXD5
A	-9	GLY	-	expression tag	UNP A0A1Q4GXD5
A	-8	ARG	-	expression tag	UNP A0A1Q4GXD5
A	-7	GLU	-	expression tag	UNP A0A1Q4GXD5
A	-6	ASN	-	expression tag	UNP A0A1Q4GXD5
A	-5	LEU	-	expression tag	UNP A0A1Q4GXD5
A	-4	TYR	-	expression tag	UNP A0A1Q4GXD5
A	-3	PHE	-	expression tag	UNP A0A1Q4GXD5
A	-2	GLN	-	expression tag	UNP A0A1Q4GXD5
A	-1	GLY	-	expression tag	UNP A0A1Q4GXD5
A	0	HIS	-	expression tag	UNP A0A1Q4GXD5
A	?	-	LEU	deletion	UNP A0A1Q4GXD5

- Molecule 2 is [(2R,3R,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-3-HYDROXY-4-(PHOSPHONOXY)TETRAHYDROFURAN-2-YL]METHYL [(2R,3S,4S)-3,4-DIHYDROXYTETRAHYDROFURAN-2-YL]METHYL DIHYDROGEN DIPHOSPHATE (CCD ID: NA7) (formula: C₁₅H₂₄N₅O₁₆P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			39	15	5	16	3		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total	O	0	0
			10	10		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Staphylopin dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	186.10Å 49.26Å 59.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.58 – 2.49 38.58 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.58-2.49) 99.2 (38.58-2.49)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.98 (at 2.48Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.219 , 0.266 0.219 , 0.267	Depositor DCC
R_{free} test set	1960 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å ²)	50.7	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 32.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6860	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA7, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/3551	0.74	0/4799

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3476	3321	3469	61	0
2	A	39	0	20	3	0
3	A	6	8	8	0	0
4	A	10	0	0	0	0
All	All	3531	3329	3497	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:VAL:HG21	2:A:501:NA7:H51'	1.52	0.91
1:A:285:TYR:OH	1:A:322:ASP:OD1	1.98	0.80
1:A:57:ILE:HG12	1:A:62:HIS:HB2	1.65	0.78
1:A:57:ILE:CG1	1:A:62:HIS:HB2	2.20	0.71
1:A:356:ARG:NH1	2:A:501:NA7:HN6'	2.09	0.68
1:A:273:SER:OG	1:A:301:GLU:OE2	2.11	0.67
1:A:169:LYS:NZ	1:A:197:GLU:OE1	2.21	0.66
1:A:12:VAL:CG2	2:A:501:NA7:H51'	2.25	0.66
1:A:15:GLN:HE22	1:A:64:HIS:HB3	1.65	0.62
1:A:217:PRO:O	1:A:251:LEU:HD11	2.02	0.58
1:A:226:LYS:O	1:A:230:GLU:O	2.22	0.57
1:A:15:GLN:NE2	1:A:64:HIS:HB3	2.18	0.56
1:A:3:LYS:HE3	1:A:83:GLU:O	2.06	0.56
1:A:257:LEU:O	1:A:261:GLU:HG3	2.06	0.55
1:A:150:THR:HA	1:A:162:LEU:O	2.09	0.53
1:A:280:MET:O	1:A:284:ASN:HB2	2.09	0.53
1:A:153:VAL:O	1:A:153:VAL:HG12	2.10	0.50
1:A:327:PRO:HG3	1:A:333:TYR:CZ	2.47	0.50
1:A:35:SER:HB2	1:A:76:ASP:OD1	2.11	0.50
1:A:397:LYS:HG2	1:A:405:LEU:HD12	1.92	0.50
1:A:379:THR:HB	1:A:382:ILE:HB	1.93	0.50
1:A:171:TYR:OH	1:A:283:GLU:OE1	2.27	0.49
1:A:223:PHE:CZ	1:A:236:VAL:HG21	2.47	0.49
1:A:172:MET:HB2	1:A:196:LEU:HD11	1.95	0.49
1:A:285:TYR:OH	1:A:322:ASP:CG	2.55	0.49
1:A:263:MET:SD	1:A:273:SER:HB2	2.52	0.49
1:A:214:VAL:C	1:A:217:PRO:HD2	2.38	0.49
1:A:107:THR:O	1:A:111:VAL:HG23	2.13	0.49
1:A:63:GLN:H	1:A:63:GLN:CD	2.21	0.48
1:A:397:LYS:HG2	1:A:405:LEU:CD1	2.43	0.48
1:A:31:VAL:HG21	1:A:77:VAL:HA	1.95	0.48
1:A:121:PHE:CE1	1:A:203:LEU:HD11	2.48	0.47
1:A:199:VAL:CG1	1:A:204:HIS:CG	2.98	0.47
1:A:255:MET:CE	1:A:312:LEU:HD22	2.44	0.47
1:A:258:MET:HE3	1:A:382:ILE:HD11	1.97	0.47
1:A:34:ALA:HB3	1:A:75:LYS:HG2	1.98	0.46
1:A:34:ALA:O	1:A:41:LYS:CG	2.65	0.45
1:A:57:ILE:CD1	1:A:62:HIS:HB2	2.46	0.45
1:A:126:ILE:HG12	1:A:421:VAL:HG12	1.99	0.45
1:A:255:MET:HE2	1:A:312:LEU:HD13	1.97	0.44
1:A:133:LYS:HE3	1:A:423:LYS:NZ	2.33	0.43
1:A:319:ILE:O	1:A:319:ILE:CG2	2.66	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:282:LYS:NZ	1:A:294:GLU:OE2	2.48	0.43
1:A:262:MET:SD	1:A:368:ILE:HG21	2.59	0.43
1:A:35:SER:OG	1:A:76:ASP:OD2	2.34	0.42
1:A:124:GLN:HG3	1:A:202:PRO:CG	2.49	0.42
1:A:305:ASP:O	1:A:309:GLU:HG3	2.18	0.42
1:A:371:ILE:O	1:A:375:LEU:HD13	2.20	0.42
1:A:12:VAL:HG22	1:A:146:TYR:CE2	2.54	0.42
1:A:199:VAL:HG11	1:A:204:HIS:CG	2.54	0.42
1:A:126:ILE:HD11	1:A:367:MET:HE2	2.02	0.42
1:A:41:LYS:O	1:A:45:GLN:HG3	2.20	0.41
1:A:142:SER:HB2	1:A:202:PRO:CB	2.50	0.41
1:A:199:VAL:CG1	1:A:200:GLU:N	2.83	0.41
1:A:299:HIS:O	1:A:300:PHE:C	2.64	0.41
1:A:328:ASP:HB2	1:A:332:HIS:O	2.20	0.41
1:A:417:ASP:O	1:A:421:VAL:HG13	2.20	0.41
1:A:57:ILE:HD11	1:A:62:HIS:HB2	2.03	0.41
1:A:172:MET:HE2	1:A:185:ILE:CG2	2.51	0.40
1:A:280:MET:HB3	1:A:315:ARG:CZ	2.52	0.40
1:A:153:VAL:O	1:A:153:VAL:CG1	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/450 (95%)	406 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	391/411 (95%)	378 (97%)	13 (3%)	33 61

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	GLU
1	A	57	ILE
1	A	63	GLN
1	A	87	VAL
1	A	135	SER
1	A	150	THR
1	A	297	ILE
1	A	315	ARG
1	A	384	GLN
1	A	407	SER
1	A	418	LYS
1	A	420	LEU
1	A	421	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	GLN
1	A	113	HIS
1	A	384	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NA7	A	501	-	42,42,42	4.15	21 (50%)	60,65,65	2.48	17 (28%)
3	GOL	A	502	-	5,5,5	0.54	0	5,5,5	1.42	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NA7	A	501	-	-	8/27/56/56	0/4/4/4
3	GOL	A	502	-	-	2/4/4/4	-

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NA7	O4D-C4D	10.57	1.61	1.44
2	A	501	NA7	C3B-C4B	-9.85	1.28	1.53
2	A	501	NA7	C2B-C1B	-9.66	1.29	1.53
2	A	501	NA7	C3D-C4D	-8.49	1.31	1.53
2	A	501	NA7	PA-O3	8.30	1.68	1.59
2	A	501	NA7	P2B-O2B	7.06	1.71	1.59
2	A	501	NA7	O4B-C1B	5.74	1.55	1.42
2	A	501	NA7	C3B-C2B	5.48	1.65	1.53
2	A	501	NA7	C6A-N6A	5.32	1.47	1.34
2	A	501	NA7	O4D-C1D	-5.09	1.33	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NA7	O4B-C4B	4.92	1.55	1.45
2	A	501	NA7	O3D-C3D	3.35	1.51	1.43
2	A	501	NA7	O2D-C2D	-2.83	1.37	1.43
2	A	501	NA7	PN-O3	2.79	1.62	1.59
2	A	501	NA7	PN-O5D	2.39	1.68	1.59
2	A	501	NA7	C1D-C2D	2.39	1.56	1.51
2	A	501	NA7	C5A-C4A	-2.33	1.34	1.39
2	A	501	NA7	C2A-N1A	2.28	1.38	1.33
2	A	501	NA7	C8A-N7A	2.26	1.36	1.31
2	A	501	NA7	C1B-N9A	2.24	1.52	1.46
2	A	501	NA7	C4A-N3A	2.10	1.38	1.34

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NA7	N6A-C6A-N1A	-6.86	103.10	118.38
2	A	501	NA7	C4A-N9A-C1B	-6.04	112.50	126.63
2	A	501	NA7	C5A-C6A-N6A	5.80	137.65	123.29
2	A	501	NA7	N9A-C8A-N7A	-5.61	105.98	113.94
2	A	501	NA7	N3A-C2A-N1A	-5.53	120.20	128.58
2	A	501	NA7	C5A-C4A-N3A	-5.32	119.39	126.72
2	A	501	NA7	C1B-N9A-C8A	4.79	137.72	127.09
2	A	501	NA7	C5A-N7A-C8A	4.04	109.81	103.45
2	A	501	NA7	N3A-C4A-N9A	3.99	133.96	127.17
2	A	501	NA7	C4A-N9A-C8A	3.85	109.78	105.74
2	A	501	NA7	O4B-C1B-C2B	-3.58	100.43	106.59
2	A	501	NA7	C2A-N3A-C4A	3.56	120.53	111.83
2	A	501	NA7	C4B-O4B-C1B	-2.86	103.15	109.47
2	A	501	NA7	C6A-C5A-C4A	2.61	120.75	117.18
3	A	502	GOL	C3-C2-C1	-2.51	102.61	111.80
2	A	501	NA7	C5D-C4D-C3D	-2.47	106.32	115.21
2	A	501	NA7	C4A-C5A-N7A	-2.46	107.77	110.58
2	A	501	NA7	C1D-O4D-C4D	-2.10	103.42	108.06

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NA7	PN-O3-PA-O5B
2	A	501	NA7	C5D-O5D-PN-O3
2	A	501	NA7	C5D-O5D-PN-O1N
3	A	502	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	501	NA7	O4D-C4D-C5D-O5D
2	A	501	NA7	C3D-C4D-C5D-O5D
2	A	501	NA7	C3B-C2B-O2B-P2B
3	A	502	GOL	O1-C1-C2-O2
2	A	501	NA7	C1B-C2B-O2B-P2B
2	A	501	NA7	C4B-C5B-O5B-PA

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NA7	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/450 (95%)	0.38	20 (4%) 36 32	44, 53, 67, 96	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	4.9
1	A	153	VAL	2.7
1	A	39	LYS	2.7
1	A	428	ASN	2.7
1	A	168	LYS	2.6
1	A	373	LYS	2.6
1	A	427	ILE	2.6
1	A	413	LEU	2.5
1	A	285	TYR	2.5
1	A	151	ARG	2.5
1	A	63	GLN	2.4
1	A	64	HIS	2.3
1	A	98	ASP	2.3
1	A	412	ASN	2.2
1	A	402	ASP	2.2
1	A	297	ILE	2.2
1	A	193	LYS	2.2
1	A	319	ILE	2.1
1	A	155	LYS	2.1
1	A	349	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NA7	A	501	39/39	0.83	0.15	75,76,79,81	0
3	GOL	A	502	6/6	0.83	0.18	52,52,73,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.