



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:46 PM UTC

PDB ID : 9C4Y / pdb_00009c4y
Title : Menin mutant - T349M
Authors : Clegg, B.D.; Cierpicki, T.; Grembecka, J.
Deposited on : 2024-06-05
Resolution : 1.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	489	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	606	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8158 atoms, of which 3787 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Menin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	468	Total	C	H	N	O	S	127	19	0
			7470	2388	3731	641	691	19			

There are 39 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP O00255
A	-3	GLY	-	expression tag	UNP O00255
A	-2	SER	-	expression tag	UNP O00255
A	-1	SER	-	expression tag	UNP O00255
A	0	SER	-	expression tag	UNP O00255
A	?	-	ILE	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	THR	deletion	UNP O00255
A	?	-	ASN	deletion	UNP O00255
A	?	-	VAL	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	GLU	deletion	UNP O00255
A	?	-	LEU	deletion	UNP O00255
A	?	-	THR	deletion	UNP O00255
A	?	-	PHE	deletion	UNP O00255
A	?	-	GLN	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	SER	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	ALA	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	ASP	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	GLY	deletion	UNP O00255
A	344	MET	THR	engineered mutation	UNP O00255
A	?	-	GLU	deletion	UNP O00255

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLU	deletion	UNP O00255
A	?	-	ARG	deletion	UNP O00255
A	?	-	PRO	deletion	UNP O00255
A	?	-	GLY	deletion	UNP O00255
A	?	-	GLU	deletion	UNP O00255
A	?	-	GLN	deletion	UNP O00255
A	?	-	SER	deletion	UNP O00255
A	?	-	GLN	deletion	UNP O00255
A	?	-	GLY	deletion	UNP O00255
A	?	-	THR	deletion	UNP O00255
A	?	-	GLN	deletion	UNP O00255
A	541	ALA	THR	variant	UNP O00255

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	4	1
			20	4	12	4		
2	A	1	Total	C	H	O	2	0
			10	2	6	2		

- Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	2	0
			31	8	18	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	2	0
			17	4	10	3		
4	A	1	Total	C	H	O	2	0
			17	4	10	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	588	Total	O	0	0
			588	588		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Menin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.02Å 80.27Å 124.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.92 – 1.31 28.92 – 1.31	Depositor EDS
% Data completeness (in resolution range)	99.6 (28.92-1.31) 99.6 (28.92-1.31)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.31Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.162 , 0.183 0.162 , 0.183	Depositor DCC
R_{free} test set	5836 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	11.3	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8158	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PEG, SO4, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/3901 (0.0%)	1.17	15/5291 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	412	HIS	CE1-NE2	-5.33	1.27	1.32

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	333	ASN	CA-CB-CG	9.81	122.41	112.60
1	A	335[A]	ARG	N-CA-CB	-9.43	96.39	110.07
1	A	335[B]	ARG	N-CA-CB	-9.43	96.39	110.07
1	A	355	ARG	N-CA-C	7.75	121.84	112.38
1	A	282	ASN	CA-CB-CG	6.95	119.55	112.60
1	A	205	ASP	CA-CB-CG	6.88	119.48	112.60
1	A	438	THR	CA-CB-OG1	-6.25	100.22	109.60
1	A	456	ARG	CD-NE-CZ	5.83	132.56	124.40
1	A	549	PRO	N-CA-CB	5.55	109.10	103.00
1	A	297	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	562[A]	LYS	CA-C-N	5.28	127.61	120.38
1	A	562[A]	LYS	C-N-CA	5.28	127.61	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	562[B]	LYS	CA-C-N	5.28	127.61	120.38
1	A	562[B]	LYS	C-N-CA	5.28	127.61	120.38
1	A	285	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	21	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3739	3731	3606	32	0
2	A	12	18	18	0	0
3	A	13	18	18	0	0
4	A	14	20	20	8	0
5	A	5	0	0	0	0
6	A	588	0	0	11	0
All	All	4371	3787	3662	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:335[A]:ARG:NH2	1:A:336[A]:GLU:OE1	2.00	0.95
1:A:335[A]:ARG:CZ	1:A:336[A]:GLU:OE1	2.16	0.93
1:A:335[A]:ARG:HD2	6:A:982:HOH:O	1.75	0.84
1:A:355:ARG:O	1:A:358:GLU:OE1	2.02	0.78
1:A:374:ASN:HB3	6:A:1088:HOH:O	1.95	0.64
1:A:46:HIS:CE1	4:A:606:PEG:H41	2.36	0.60
1:A:366:GLU:HG3	6:A:1038:HOH:O	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:606:PEG:C4	6:A:744:HOH:O	2.53	0.57
1:A:46:HIS:NE2	4:A:606:PEG:H41	2.19	0.57
1:A:46:HIS:HE1	4:A:606:PEG:H11	1.72	0.54
1:A:322[B]:MET:CE	4:A:604:PEG:H41	2.38	0.54
1:A:2:GLY:HA2	6:A:882:HOH:O	2.11	0.51
1:A:366:GLU:HG2	6:A:1179:HOH:O	2.11	0.50
1:A:438:THR:HB	6:A:742:HOH:O	2.12	0.49
1:A:437:ALA:HB1	1:A:576:LYS:HG2	1.95	0.49
1:A:96[A]:GLN:O	1:A:96[A]:GLN:HG2	2.12	0.48
1:A:96[A]:GLN:NE2	1:A:128:SER:OG	2.47	0.48
1:A:318:ILE:HG23	1:A:344[B]:MET:HG3	1.96	0.47
1:A:322[B]:MET:CE	4:A:604:PEG:C4	2.94	0.46
1:A:335[A]:ARG:HD2	6:A:756:HOH:O	2.14	0.46
1:A:353:TYR:O	1:A:429:THR:HG23	2.16	0.45
1:A:233:LYS:HG3	1:A:272:HIS:CE1	2.52	0.44
1:A:108:ARG:HH11	1:A:108:ARG:HD3	1.68	0.43
1:A:570:ILE:HG12	6:A:1093:HOH:O	2.19	0.43
1:A:96[A]:GLN:O	1:A:96[A]:GLN:CG	2.65	0.43
4:A:606:PEG:H42	6:A:744:HOH:O	2.17	0.43
1:A:454[B]:LYS:O	1:A:549:PRO:HD2	2.19	0.43
1:A:322[B]:MET:HE2	4:A:604:PEG:H42	2.01	0.43
1:A:108:ARG:HD2	1:A:170:LEU:CD2	2.50	0.41
1:A:452:ARG:NH1	1:A:570:ILE:HD11	2.35	0.41
1:A:51:ASN:ND2	1:A:53:VAL:H	2.19	0.40
1:A:30:GLU:O	6:A:701:HOH:O	2.22	0.40
1:A:360:ILE:HD12	1:A:360:ILE:HA	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/489 (99%)	477 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/410 (97%)	398 (100%)	1 (0%)	86	67

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	429	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	ASN
1	A	212	ASN
1	A	282	ASN
1	A	331	ASN
1	A	333	ASN
1	A	339	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PG4	A	602	-	12,12,12	0.72	0	11,11,11	1.10	2 (18%)
4	PEG	A	606	-	6,6,6	0.65	0	5,5,5	0.83	0
2	EDO	A	601[A]	-	3,3,3	0.15	0	2,2,2	0.16	0
4	PEG	A	604	-	6,6,6	0.14	0	5,5,5	0.18	0
2	EDO	A	601[B]	-	3,3,3	0.13	0	2,2,2	0.08	0
5	SO4	A	605	-	4,4,4	0.58	0	6,6,6	0.65	0
2	EDO	A	603	-	3,3,3	0.24	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	A	602	-	-	3/10/10/10	-
4	PEG	A	606	-	-	2/4/4/4	-
2	EDO	A	601[A]	-	-	1/1/1/1	-
4	PEG	A	604	-	-	2/4/4/4	-
2	EDO	A	601[B]	-	-	1/1/1/1	-
2	EDO	A	603	-	-	1/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	PG4	O4-C6-C5	2.60	122.20	110.35
3	A	602	PG4	C5-O3-C4	2.31	123.39	113.26

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	PG4	O3-C5-C6-O4
4	A	606	PEG	O1-C1-C2-O2
4	A	606	PEG	O2-C3-C4-O4
4	A	604	PEG	O2-C3-C4-O4
3	A	602	PG4	O1-C1-C2-O2
2	A	601[B]	EDO	O1-C1-C2-O2
3	A	602	PG4	C6-C5-O3-C4
2	A	601[A]	EDO	O1-C1-C2-O2
2	A	603	EDO	O1-C1-C2-O2
4	A	604	PEG	C1-C2-O2-C3

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	606	PEG	5	0
4	A	604	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	468/489 (95%)	-0.02	27 (5%)	29 33	4, 11, 27, 43	10 (2%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	460	VAL	7.2
1	A	461	PRO	6.2
1	A	111	GLY	4.0
1	A	2	GLY	4.0
1	A	566	VAL	3.7
1	A	110	GLY	3.7
1	A	134	PHE	3.7
1	A	333	ASN	3.6
1	A	581	ALA	3.6
1	A	587	MET	3.2
1	A	126	TRP	3.2
1	A	570	ILE	3.1
1	A	332	ARG	3.0
1	A	568	THR	2.7
1	A	136	ASP	2.7
1	A	30	GLU	2.6
1	A	458	VAL	2.6
1	A	109	GLU	2.6
1	A	567	ALA	2.5
1	A	429	THR	2.3
1	A	549	PRO	2.3
1	A	550	VAL	2.2
1	A	202	GLY	2.2
1	A	582	GLN	2.2
1	A	299	LEU	2.1
1	A	204	GLU	2.1
1	A	563	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	601[A]	4/4	0.71	0.16	27,27,29,29	10
2	EDO	A	601[B]	4/4	0.71	0.16	29,29,31,31	10
4	PEG	A	606	7/7	0.75	0.18	24,34,36,38	2
4	PEG	A	604	7/7	0.76	0.14	42,46,54,54	2
5	SO4	A	605	5/5	0.82	0.17	23,24,29,41	0
3	PG4	A	602	13/13	0.85	0.14	18,27,40,41	2
2	EDO	A	603	4/4	0.89	0.12	37,38,39,39	2

6.5 Other polymers [i](#)

There are no such residues in this entry.