



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 11:59 PM UTC

PDB ID : 4C5R / pdb_00004c5r
Title : Structural Investigations into the Stereochemistry and Activity of a Phenylalanine-2,3-Aminomutase from *Taxus chinensis*
Authors : Wybenga, G.G.; Szymanski, W.; Wu, B.; Feringa, B.L.; Janssen, D.B.; Dijkstra, B.W.
Deposited on : 2013-09-16
Resolution : 2.14 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	705	 2% 80% 9% • 10%
1	B	705	 1% 80% 10% • 9%
1	C	705	 1% 79% 10% • 9%
1	D	705	 1% 81% 8% • 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLALANINE AMMONIA-LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	635	Total	C	N	O	S	0	0	0
			4926	3129	847	927	23			
1	B	640	Total	C	N	O	S	0	0	0
			4967	3152	855	937	23			
1	C	640	Total	C	N	O	S	0	0	0
			4968	3153	855	937	23			
1	D	636	Total	C	N	O	S	0	0	0
			4935	3132	848	932	23			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	175	MDO	ALA	chromophore	UNP Q68G84
A	175	MDO	SER	chromophore	UNP Q68G84
A	175	MDO	GLY	chromophore	UNP Q68G84
B	175	MDO	ALA	chromophore	UNP Q68G84
B	175	MDO	SER	chromophore	UNP Q68G84
B	175	MDO	GLY	chromophore	UNP Q68G84
C	175	MDO	ALA	chromophore	UNP Q68G84
C	175	MDO	SER	chromophore	UNP Q68G84
C	175	MDO	GLY	chromophore	UNP Q68G84
D	175	MDO	ALA	chromophore	UNP Q68G84
D	175	MDO	SER	chromophore	UNP Q68G84
D	175	MDO	GLY	chromophore	UNP Q68G84
A	-19	MET	-	expression tag	UNP Q68G84
A	-18	GLY	-	expression tag	UNP Q68G84
A	-17	SER	-	expression tag	UNP Q68G84
A	-16	SER	-	expression tag	UNP Q68G84
A	-15	HIS	-	expression tag	UNP Q68G84
A	-14	HIS	-	expression tag	UNP Q68G84
A	-13	HIS	-	expression tag	UNP Q68G84
A	-12	HIS	-	expression tag	UNP Q68G84
A	-11	HIS	-	expression tag	UNP Q68G84

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	HIS	-	expression tag	UNP Q68G84
A	-9	SER	-	expression tag	UNP Q68G84
A	-8	SER	-	expression tag	UNP Q68G84
A	-7	GLY	-	expression tag	UNP Q68G84
A	-6	LEU	-	expression tag	UNP Q68G84
A	-5	VAL	-	expression tag	UNP Q68G84
A	-4	PRO	-	expression tag	UNP Q68G84
A	-3	ARG	-	expression tag	UNP Q68G84
A	-2	GLY	-	expression tag	UNP Q68G84
A	-1	SER	-	expression tag	UNP Q68G84
A	0	HIS	-	expression tag	UNP Q68G84
B	-19	MET	-	expression tag	UNP Q68G84
B	-18	GLY	-	expression tag	UNP Q68G84
B	-17	SER	-	expression tag	UNP Q68G84
B	-16	SER	-	expression tag	UNP Q68G84
B	-15	HIS	-	expression tag	UNP Q68G84
B	-14	HIS	-	expression tag	UNP Q68G84
B	-13	HIS	-	expression tag	UNP Q68G84
B	-12	HIS	-	expression tag	UNP Q68G84
B	-11	HIS	-	expression tag	UNP Q68G84
B	-10	HIS	-	expression tag	UNP Q68G84
B	-9	SER	-	expression tag	UNP Q68G84
B	-8	SER	-	expression tag	UNP Q68G84
B	-7	GLY	-	expression tag	UNP Q68G84
B	-6	LEU	-	expression tag	UNP Q68G84
B	-5	VAL	-	expression tag	UNP Q68G84
B	-4	PRO	-	expression tag	UNP Q68G84
B	-3	ARG	-	expression tag	UNP Q68G84
B	-2	GLY	-	expression tag	UNP Q68G84
B	-1	SER	-	expression tag	UNP Q68G84
B	0	HIS	-	expression tag	UNP Q68G84
C	-19	MET	-	expression tag	UNP Q68G84
C	-18	GLY	-	expression tag	UNP Q68G84
C	-17	SER	-	expression tag	UNP Q68G84
C	-16	SER	-	expression tag	UNP Q68G84
C	-15	HIS	-	expression tag	UNP Q68G84
C	-14	HIS	-	expression tag	UNP Q68G84
C	-13	HIS	-	expression tag	UNP Q68G84
C	-12	HIS	-	expression tag	UNP Q68G84
C	-11	HIS	-	expression tag	UNP Q68G84
C	-10	HIS	-	expression tag	UNP Q68G84
C	-9	SER	-	expression tag	UNP Q68G84

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-8	SER	-	expression tag	UNP Q68G84
C	-7	GLY	-	expression tag	UNP Q68G84
C	-6	LEU	-	expression tag	UNP Q68G84
C	-5	VAL	-	expression tag	UNP Q68G84
C	-4	PRO	-	expression tag	UNP Q68G84
C	-3	ARG	-	expression tag	UNP Q68G84
C	-2	GLY	-	expression tag	UNP Q68G84
C	-1	SER	-	expression tag	UNP Q68G84
C	0	HIS	-	expression tag	UNP Q68G84
D	-19	MET	-	expression tag	UNP Q68G84
D	-18	GLY	-	expression tag	UNP Q68G84
D	-17	SER	-	expression tag	UNP Q68G84
D	-16	SER	-	expression tag	UNP Q68G84
D	-15	HIS	-	expression tag	UNP Q68G84
D	-14	HIS	-	expression tag	UNP Q68G84
D	-13	HIS	-	expression tag	UNP Q68G84
D	-12	HIS	-	expression tag	UNP Q68G84
D	-11	HIS	-	expression tag	UNP Q68G84
D	-10	HIS	-	expression tag	UNP Q68G84
D	-9	SER	-	expression tag	UNP Q68G84
D	-8	SER	-	expression tag	UNP Q68G84
D	-7	GLY	-	expression tag	UNP Q68G84
D	-6	LEU	-	expression tag	UNP Q68G84
D	-5	VAL	-	expression tag	UNP Q68G84
D	-4	PRO	-	expression tag	UNP Q68G84
D	-3	ARG	-	expression tag	UNP Q68G84
D	-2	GLY	-	expression tag	UNP Q68G84
D	-1	SER	-	expression tag	UNP Q68G84
D	0	HIS	-	expression tag	UNP Q68G84

- Molecule 2 is (3S)-3-amino-2,2-difluoro-3-phenylpropanoic acid (CCD ID: BQ7) (formula: $C_9H_9F_2NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			14	9	2	1	2		
2	B	1	Total	C	F	N	O	0	0
			14	9	2	1	2		
2	C	1	Total	C	F	N	O	0	0
			14	9	2	1	2		
2	D	1	Total	C	F	N	O	0	0
			14	9	2	1	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	181	Total	O	0	0
			181	181		
4	B	154	Total	O	0	0
			154	154		
4	C	161	Total	O	0	0
			161	161		
4	D	145	Total	O	0	0
			145	145		

- Molecule 1: PHENYLALANINE AMMONIA-LYASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	99.39Å 145.48Å 99.56Å 90.00° 99.54° 90.00°	Depositor
Resolution (Å)	49.09 – 2.14 49.09 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.09-2.14) 99.4 (49.09-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.172 , 0.210 0.173 , 0.209	Depositor DCC
R_{free} test set	7602 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 31.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	20505	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MDO, BQ7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	6/4996 (0.1%)	0.86	4/6778 (0.1%)
1	B	0.75	7/5038 (0.1%)	0.87	0/6835
1	C	0.75	7/5039 (0.1%)	0.86	2/6836 (0.0%)
1	D	0.73	3/5005 (0.1%)	0.88	1/6790 (0.0%)
All	All	0.74	23/20078 (0.1%)	0.87	7/27239 (0.0%)

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	368	GLY	C-O	-7.13	1.18	1.24
1	C	368	GLY	C-O	-7.12	1.18	1.24
1	B	360	HIS	CG-ND1	-6.92	1.30	1.38
1	A	456	GLN	C-O	-6.84	1.14	1.23
1	C	360	HIS	CG-ND1	-6.67	1.30	1.38

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	535	VAL	N-CA-CB	5.88	115.55	110.08
1	D	29	VAL	CB-CA-C	5.43	118.05	110.99
1	A	322	TYR	N-CA-C	5.36	117.12	111.28
1	C	29	VAL	CB-CA-C	5.29	117.87	110.99
1	A	369	ALA	N-CA-C	5.23	117.53	110.06

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4926	0	4988	42	0
1	B	4967	0	5021	43	0
1	C	4968	0	5023	49	0
1	D	4935	0	4987	33	0
2	A	14	0	7	0	0
2	B	14	0	7	0	0
2	C	14	0	7	0	0
2	D	14	0	7	1	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	181	0	0	7	0
4	B	154	0	0	0	0
4	C	161	0	0	4	0
4	D	145	0	0	3	0
All	All	20505	0	20063	143	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 143 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:506:VAL:HG21	1:B:599:VAL:HG11	1.45	0.96
1:A:352:ALA:HB3	1:B:279:VAL:CG1	2.03	0.87
1:C:381:ASP:OD2	1:C:448:THR:HB	1.74	0.86
1:A:279:VAL:HG13	1:B:352:ALA:HB3	1.57	0.84
1:C:279:VAL:HG13	1:D:352:ALA:HB3	1.60	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	622/705 (88%)	613 (99%)	9 (1%)	0	100	100
1	B	627/705 (89%)	615 (98%)	12 (2%)	0	100	100
1	C	627/705 (89%)	616 (98%)	11 (2%)	0	100	100
1	D	623/705 (88%)	610 (98%)	13 (2%)	0	100	100
All	All	2499/2820 (89%)	2454 (98%)	45 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	537/592 (91%)	518 (96%)	19 (4%)	32	31
1	B	542/592 (92%)	523 (96%)	19 (4%)	32	31
1	C	542/592 (92%)	521 (96%)	21 (4%)	28	26
1	D	538/592 (91%)	516 (96%)	22 (4%)	27	24
All	All	2159/2368 (91%)	2078 (96%)	81 (4%)	29	28

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	640	VAL
1	D	392	LYS
1	D	26	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	164	SER
1	D	553	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	47	HIS
1	D	162	ASN
1	D	674	GLN
1	D	671	HIS
1	B	360	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MDO	D	175	2,1	11,13,14	2.91	4 (36%)	15,18,20	1.81	4 (26%)
1	MDO	C	175	2,1	11,13,14	2.81	5 (45%)	15,18,20	1.90	3 (20%)
1	MDO	B	175	2,1	11,13,14	2.90	5 (45%)	15,18,20	2.07	4 (26%)
1	MDO	A	175	2,1	11,13,14	2.88	4 (36%)	15,18,20	2.01	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MDO	D	175	2,1	-	2/4/23/24	0/1/1/1
1	MDO	C	175	2,1	-	2/4/23/24	0/1/1/1
1	MDO	B	175	2,1	-	2/4/23/24	0/1/1/1
1	MDO	A	175	2,1	-	3/4/23/24	0/1/1/1

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	175	MDO	O2-C2	6.72	1.36	1.23
1	A	175	MDO	O2-C2	6.66	1.36	1.23
1	B	175	MDO	O2-C2	6.21	1.35	1.23
1	C	175	MDO	O2-C2	5.65	1.34	1.23
1	B	175	MDO	O3-C3	4.39	1.44	1.20

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	175	MDO	CA2-C2-N3	5.24	107.90	103.50
1	B	175	MDO	O2-C2-CA2	-4.67	128.04	131.02
1	B	175	MDO	CA2-C2-N3	4.48	107.27	103.50
1	A	175	MDO	O2-C2-CA2	-4.41	128.21	131.02
1	A	175	MDO	CA2-C2-N3	4.01	106.87	103.50

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	175	MDO	C3-CA3-N3-C2
1	A	175	MDO	N2-C1-CA1-CB
1	B	175	MDO	N2-C1-CA1-CB
1	C	175	MDO	N2-C1-CA1-CB
1	D	175	MDO	N2-C1-CA1-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BQ7	B	1678	1	13,14,14	1.40	2 (15%)	17,20,20	0.94	1 (5%)
2	BQ7	A	1678	1	13,14,14	1.37	1 (7%)	17,20,20	0.92	1 (5%)
2	BQ7	C	1678	1	13,14,14	1.42	2 (15%)	17,20,20	0.88	1 (5%)
3	GOL	B	1679	-	5,5,5	0.31	0	5,5,5	0.81	0
3	GOL	A	1679	-	5,5,5	0.13	0	5,5,5	0.51	0
2	BQ7	D	1678	1	13,14,14	1.33	1 (7%)	17,20,20	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BQ7	B	1678	1	-	3/14/16/16	0/1/1/1
2	BQ7	A	1678	1	-	4/14/16/16	0/1/1/1
2	BQ7	C	1678	1	-	2/14/16/16	0/1/1/1
3	GOL	B	1679	-	-	2/4/4/4	-
3	GOL	A	1679	-	-	3/4/4/4	-
2	BQ7	D	1678	1	-	3/14/16/16	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1678	BQ7	CG-CB	3.49	1.56	1.52
2	B	1678	BQ7	CG-CB	3.44	1.56	1.52
2	A	1678	BQ7	CG-CB	2.95	1.55	1.52
2	C	1678	BQ7	CG-CB	2.82	1.55	1.52
2	C	1678	BQ7	FS-CA	-2.31	1.32	1.36

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1678	BQ7	FS-CA-CB	3.17	113.44	109.23
2	A	1678	BQ7	FS-CA-CB	2.60	112.69	109.23
2	D	1678	BQ7	O-C-CA	2.44	118.32	112.54
2	C	1678	BQ7	O-C-CA	2.22	117.80	112.54
2	B	1678	BQ7	O-C-CA	2.02	117.32	112.54

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1678	BQ7	O-C-CA-FS
2	A	1678	BQ7	C-CA-CB-CG
2	A	1678	BQ7	FR-CA-CB-CG
2	B	1678	BQ7	C-CA-CB-CG
2	B	1678	BQ7	FR-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1678	BQ7	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	634/705 (89%)	-0.14	16 (2%) 58 62	11, 24, 49, 60	0
1	B	639/705 (90%)	-0.22	7 (1%) 78 81	12, 24, 40, 54	0
1	C	639/705 (90%)	-0.27	7 (1%) 78 81	12, 22, 42, 54	0
1	D	635/705 (90%)	-0.23	9 (1%) 73 77	11, 22, 45, 57	0
All	All	2547/2820 (90%)	-0.22	39 (1%) 72 76	11, 23, 44, 60	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	576	THR	3.8
1	B	55	ALA	3.7
1	D	566	LEU	3.5
1	A	586	LYS	3.2
1	A	677	LEU	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MDO	B	175	13/14	0.95	0.07	18,21,23,25	0
1	MDO	C	175	13/14	0.95	0.06	15,16,18,19	0
1	MDO	A	175	13/14	0.96	0.06	18,19,20,22	0
1	MDO	D	175	13/14	0.96	0.06	16,18,19,21	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BQ7	A	1678	14/14	0.85	0.12	26,28,36,37	0
3	GOL	A	1679	6/6	0.88	0.12	28,30,32,36	0
2	BQ7	B	1678	14/14	0.90	0.09	27,31,37,40	0
3	GOL	B	1679	6/6	0.90	0.11	28,32,32,35	0
2	BQ7	C	1678	14/14	0.93	0.08	24,25,30,31	0
2	BQ7	D	1678	14/14	0.93	0.08	27,29,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.