



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 07:29 AM UTC

PDB ID : 6C56 / pdb_00006c56
Title : Crystal structure of mutant human geranylgeranyl pyrophosphate synthase (Y246D) in its Apo form
Authors : Park, J.; Ta, V.; Tsantrizos, Y.S.; Berghuis, A.M.
Deposited on : 2018-01-15
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	
1	B	322	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranylgeranyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	0	0
			2228	1432	366	423	7			
1	B	267	Total	C	N	O	S	0	0	0
			2088	1350	337	393	8			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP O95749
A	-20	HIS	-	expression tag	UNP O95749
A	-19	HIS	-	expression tag	UNP O95749
A	-18	HIS	-	expression tag	UNP O95749
A	-17	HIS	-	expression tag	UNP O95749
A	-16	HIS	-	expression tag	UNP O95749
A	-15	HIS	-	expression tag	UNP O95749
A	-14	SER	-	expression tag	UNP O95749
A	-13	SER	-	expression tag	UNP O95749
A	-12	GLY	-	expression tag	UNP O95749
A	-11	VAL	-	expression tag	UNP O95749
A	-10	ASP	-	expression tag	UNP O95749
A	-9	LEU	-	expression tag	UNP O95749
A	-8	GLY	-	expression tag	UNP O95749
A	-7	THR	-	expression tag	UNP O95749
A	-6	GLU	-	expression tag	UNP O95749
A	-5	ASN	-	expression tag	UNP O95749
A	-4	LEU	-	expression tag	UNP O95749
A	-3	TYR	-	expression tag	UNP O95749
A	-2	PHE	-	expression tag	UNP O95749
A	-1	GLN	-	expression tag	UNP O95749
A	0	SER	-	expression tag	UNP O95749
A	246	ASP	TYR	engineered mutation	UNP O95749
B	-21	MET	-	initiating methionine	UNP O95749
B	-20	HIS	-	expression tag	UNP O95749

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	HIS	-	expression tag	UNP O95749
B	-18	HIS	-	expression tag	UNP O95749
B	-17	HIS	-	expression tag	UNP O95749
B	-16	HIS	-	expression tag	UNP O95749
B	-15	HIS	-	expression tag	UNP O95749
B	-14	SER	-	expression tag	UNP O95749
B	-13	SER	-	expression tag	UNP O95749
B	-12	GLY	-	expression tag	UNP O95749
B	-11	VAL	-	expression tag	UNP O95749
B	-10	ASP	-	expression tag	UNP O95749
B	-9	LEU	-	expression tag	UNP O95749
B	-8	GLY	-	expression tag	UNP O95749
B	-7	THR	-	expression tag	UNP O95749
B	-6	GLU	-	expression tag	UNP O95749
B	-5	ASN	-	expression tag	UNP O95749
B	-4	LEU	-	expression tag	UNP O95749
B	-3	TYR	-	expression tag	UNP O95749
B	-2	PHE	-	expression tag	UNP O95749
B	-1	GLN	-	expression tag	UNP O95749
B	0	SER	-	expression tag	UNP O95749
B	246	ASP	TYR	engineered mutation	UNP O95749

- Molecule 2 is {[(2-phenylthieno[2,3-d]pyrimidin-4-yl)amino]methylene}bis(phosphonic acid) (CCD ID: VT0) (formula: C₁₃H₁₃N₃O₆P₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	S	0	0
			25	13	3	6	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	18	Total	O	0	0
			18	18		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Geranylgeranyl pyrophosphate synthase



- Molecule 1: Geranylgeranyl pyrophosphate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	185.10Å 185.10Å 115.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.39 – 2.80 54.39 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (54.39-2.80) 99.9 (54.39-2.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.185 , 0.238 0.186 , 0.236	Depositor DCC
R_{free} test set	1270 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	91.7	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 73.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4380	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VT0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.47	17/2275 (0.7%)	1.40	18/3094 (0.6%)
1	B	1.37	10/2134 (0.5%)	1.38	18/2901 (0.6%)
All	All	1.42	27/4409 (0.6%)	1.39	36/5995 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	14	GLU	C-O	-8.22	1.16	1.24
1	A	240	ASN	N-CA	7.95	1.55	1.46
1	A	242	ASP	N-CA	7.17	1.54	1.46
1	A	237	ARG	CA-C	7.12	1.60	1.53
1	B	29	THR	N-CA	6.68	1.55	1.45
1	A	174	LYS	CA-C	6.66	1.61	1.52
1	A	211	GLY	C-O	-6.58	1.14	1.23
1	A	23	PRO	N-CA	6.54	1.55	1.47
1	B	86	PRO	C-O	-6.42	1.15	1.24
1	A	252	GLU	N-CA	6.39	1.54	1.46
1	A	237	ARG	C-O	6.25	1.32	1.23
1	B	92	ALA	C-O	-6.11	1.17	1.24
1	B	82	ILE	C-O	-6.08	1.17	1.23
1	A	241	ILE	CA-C	5.92	1.61	1.52
1	A	249	HIS	N-CA	5.69	1.53	1.46
1	B	195	SER	CA-C	5.56	1.60	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	234	LEU	CA-C	5.53	1.59	1.52
1	A	23	PRO	CA-C	5.45	1.59	1.52
1	B	29	THR	CA-C	5.35	1.59	1.52
1	A	152	THR	CA-CB	5.29	1.61	1.53
1	A	235	ARG	N-CA	5.20	1.52	1.45
1	B	10	ARG	CA-CB	5.13	1.61	1.53
1	B	206	GLU	N-CA	5.13	1.52	1.46
1	A	151	LYS	C-O	5.05	1.30	1.24
1	B	171	GLU	CA-C	5.02	1.59	1.52
1	B	8	VAL	C-O	-5.01	1.18	1.24
1	A	252	GLU	CA-C	5.00	1.59	1.52

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	LEU	N-CA-C	8.37	120.40	111.28
1	B	10	ARG	N-CA-C	-7.37	103.33	111.36
1	B	10	ARG	CB-CA-C	7.33	123.31	110.85
1	B	196	LYS	N-CA-C	7.13	119.31	107.20
1	A	31	LEU	N-CA-C	7.00	118.80	111.03
1	B	31	LEU	N-CA-C	7.00	118.60	110.97
1	A	73	ARG	NE-CZ-NH2	6.38	124.94	119.20
1	A	-1	GLN	OE1-CD-NE2	-6.21	116.39	122.60
1	B	67	GLU	N-CA-C	6.11	117.94	111.28
1	A	224	ARG	N-CA-CB	5.91	120.55	110.50
1	B	157	GLY	N-CA-C	-5.84	105.72	112.73
1	A	73	ARG	NE-CZ-NH1	-5.76	115.74	121.50
1	A	154	GLY	CA-C-N	5.74	127.97	120.28
1	A	154	GLY	C-N-CA	5.74	127.97	120.28
1	A	19	LEU	N-CA-C	5.73	117.61	111.36
1	A	20	LEU	CB-CA-C	-5.73	101.28	110.79
1	A	202	LYS	N-CA-C	-5.67	103.63	110.44
1	B	207	ASP	N-CA-C	5.63	117.09	111.07
1	B	135	ASN	N-CA-C	5.47	118.96	112.72
1	B	112	VAL	N-CA-C	-5.47	105.51	110.82
1	A	131	TYR	N-CA-C	5.43	117.62	111.11
1	B	74	ARG	NE-CZ-NH2	5.32	123.99	119.20
1	B	29	THR	N-CA-C	5.28	120.59	113.72
1	B	43	GLU	CA-CB-CG	5.28	124.66	114.10
1	A	237	ARG	CA-C-N	5.28	130.37	121.14
1	A	237	ARG	C-N-CA	5.28	130.37	121.14
1	B	22	LEU	CA-C-N	5.26	125.26	119.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	22	LEU	C-N-CA	5.26	125.26	119.90
1	B	90	ASN	N-CA-CB	-5.23	102.40	110.20
1	A	243	ILE	N-CA-C	-5.21	98.50	109.34
1	A	265	LYS	N-CA-C	-5.19	104.88	111.11
1	A	290	HIS	N-CA-C	-5.10	105.61	111.07
1	B	153	GLY	N-CA-C	-5.04	106.68	112.73
1	B	89	ILE	CB-CA-C	-5.04	105.44	112.04
1	A	130	ILE	N-CA-C	5.03	115.75	110.62
1	B	117	ARG	NE-CZ-NH2	5.02	123.72	119.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	-1	GLN	Sidechain
1	A	223	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2086	30	0
1	B	2088	0	1987	20	0
2	B	25	0	0	0	0
3	A	21	0	0	0	0
3	B	18	0	0	0	0
All	All	4380	0	4073	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:HIS:O	1:A:251:LEU:N	2.12	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:0:SER:O	1:B:4:THR:HG22	1.80	0.81
1:A:205:CYS:HB2	1:A:241:ILE:O	1.81	0.79
1:A:249:HIS:O	1:A:250:TYR:C	2.30	0.74
1:A:133:ARG:NH2	1:A:211:GLY:O	2.22	0.72
1:A:115:PHE:CE2	1:A:119:LEU:HD12	2.30	0.65
1:A:97:PHE:CE1	1:B:119:LEU:HD13	2.38	0.58
1:A:89:ILE:HD11	1:B:89:ILE:HD11	1.87	0.56
1:A:249:HIS:O	1:A:252:GLU:N	2.38	0.55
1:B:137:THR:HG22	1:B:138:CYS:H	1.71	0.54
1:A:129:ASP:OD1	1:A:133:ARG:NH1	2.37	0.52
1:A:192:ASN:OD1	1:A:192:ASN:C	2.52	0.52
1:A:14:GLU:HB2	1:A:15:PRO:HD3	1.92	0.52
1:A:272:TYR:CE1	1:A:285:VAL:HG13	2.46	0.50
1:A:5:GLN:O	1:A:9:GLN:HG2	2.12	0.50
1:A:129:ASP:OD1	1:A:129:ASP:C	2.55	0.49
1:A:141:GLU:OE1	1:A:219:HIS:HD2	1.95	0.49
1:B:18:TYR:O	1:B:21:GLN:HG3	2.13	0.48
1:B:147:MET:HE3	1:B:151:LYS:CD	2.43	0.48
1:B:147:MET:HE3	1:B:151:LYS:HD3	1.95	0.48
1:A:205:CYS:CB	1:A:241:ILE:O	2.57	0.48
1:A:249:HIS:C	1:A:251:LEU:N	2.70	0.48
1:B:144:TYR:OH	1:B:189:ASP:OD2	2.27	0.48
1:A:115:PHE:CZ	1:A:119:LEU:HD12	2.49	0.48
1:B:45:LYS:HD2	1:B:166:PHE:CE1	2.49	0.48
1:A:210:GLU:HA	1:A:210:GLU:OE1	2.15	0.47
1:A:138:CYS:HB2	1:A:238:THR:HG21	1.97	0.47
1:B:17:LYS:O	1:B:18:TYR:C	2.53	0.47
1:B:292:SER:O	1:B:293:LYS:C	2.58	0.46
1:A:32:SER:O	1:A:33:GLN:C	2.59	0.46
1:A:239:GLU:HA	1:A:242:ASP:HB2	1.99	0.45
1:B:137:THR:HG22	1:B:138:CYS:N	2.31	0.44
1:A:259:TYR:HA	1:A:262:ASN:HD22	1.81	0.44
1:A:115:PHE:CZ	1:A:119:LEU:CD1	3.00	0.44
1:B:174:LYS:HA	1:B:177:LEU:HD12	2.01	0.42
1:A:246:ASP:O	1:A:247:CYS:C	2.62	0.42
1:B:45:LYS:HD2	1:B:166:PHE:CD1	2.54	0.42
1:A:138:CYS:CB	1:A:238:THR:HG21	2.50	0.42
1:A:151:LYS:HB3	1:A:152:THR:H	1.42	0.41
1:B:219:HIS:CD2	1:B:259:TYR:CE2	3.08	0.41
1:B:209:THR:HB	1:B:212:LYS:O	2.20	0.41
1:A:145:LYS:HG2	1:A:182:LEU:HD22	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:THR:CG2	1:B:138:CYS:H	2.33	0.41
1:B:140:THR:OG1	1:B:143:GLU:HG3	2.21	0.41
1:A:215:PHE:HB3	1:A:216:PRO:HD3	2.02	0.40
1:A:247:CYS:C	1:A:249:HIS:N	2.77	0.40
1:B:169:TYR:CE2	1:B:171:GLU:HB2	2.56	0.40
1:B:191:ALA:O	1:B:192:ASN:C	2.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/322 (87%)	251 (90%)	26 (9%)	3 (1%)	11	36
1	B	259/322 (80%)	240 (93%)	18 (7%)	1 (0%)	30	60
All	All	539/644 (84%)	491 (91%)	44 (8%)	4 (1%)	18	47

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	151	LYS
1	A	250	TYR
1	A	239	GLU
1	B	294	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/292 (77%)	214 (95%)	11 (5%)	22	55
1	B	217/292 (74%)	209 (96%)	8 (4%)	30	65
All	All	442/584 (76%)	423 (96%)	19 (4%)	26	60

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLU
1	A	27	VAL
1	A	31	LEU
1	A	32	SER
1	A	44	ASP
1	A	87	SER
1	A	114	LEU
1	A	121	GLU
1	A	152	THR
1	A	188	ASP
1	A	221	ILE
1	B	19	LEU
1	B	31	LEU
1	B	43	GLU
1	B	70	SER
1	B	156	PHE
1	B	188	ASP
1	B	217	THR
1	B	249	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	80	HIS
1	A	90	ASN
1	A	93	ASN
1	A	126	GLN
1	A	150	GLN
1	A	219	HIS
1	A	262	ASN
1	B	5	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	93	ASN
1	B	124	GLN
1	B	135	ASN
1	B	150	GLN
1	B	194	HIS
1	B	219	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	VT0	B	401	-	27,27,27	4.63	13 (48%)	34,41,41	4.47	13 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VT0	B	401	-	-	0/20/20/20	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	VT0	C6-SAE	-15.44	1.56	1.73
2	B	401	VT0	PAM-OAR	8.68	1.64	1.49
2	B	401	VT0	C5-C4	-7.51	1.32	1.42
2	B	401	VT0	PAL-OAO	7.22	1.61	1.49
2	B	401	VT0	PAL-OAP	6.45	1.65	1.54
2	B	401	VT0	PAM-OAS	5.91	1.64	1.54
2	B	401	VT0	C5-C6	-4.25	1.36	1.41
2	B	401	VT0	PAM-OAQ	-3.94	1.48	1.54
2	B	401	VT0	CAT-C2	-3.11	1.39	1.48
2	B	401	VT0	PAL-OAN	-3.09	1.49	1.54
2	B	401	VT0	CAK-NAJ	2.71	1.49	1.46
2	B	401	VT0	C5-CAC	-2.71	1.35	1.44
2	B	401	VT0	PAM-CAK	2.67	1.85	1.83

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	VT0	C6-SAE-CAD	21.78	97.08	90.62
2	B	401	VT0	C5-C6-N1	-5.86	122.60	126.55
2	B	401	VT0	SAE-C6-N1	5.73	129.18	122.51
2	B	401	VT0	C6-C5-CAC	4.15	115.45	111.76
2	B	401	VT0	C5-C6-SAE	-3.91	108.22	111.38
2	B	401	VT0	PAM-CAK-PAL	-3.79	109.69	116.24
2	B	401	VT0	CAT-C2-N1	3.68	123.48	117.40
2	B	401	VT0	N1-C2-N3	-3.42	119.51	125.13
2	B	401	VT0	CAC-CAD-SAE	-3.38	106.95	113.90
2	B	401	VT0	C4-N3-C2	2.69	121.22	116.36
2	B	401	VT0	OAQ-PAM-CAK	2.53	111.05	104.77
2	B	401	VT0	OAR-PAM-CAK	-2.48	105.54	110.20
2	B	401	VT0	OAN-PAL-OAP	2.28	113.78	107.58

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/322 (88%)	-0.14	8 (2%) 55 45	68, 106, 149, 169	0
1	B	267/322 (82%)	-0.25	5 (1%) 66 57	62, 102, 151, 183	0
All	All	553/644 (85%)	-0.19	13 (2%) 59 49	62, 104, 151, 183	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	224	ARG	3.7
1	B	197	GLU	3.7
1	A	-3	TYR	3.5
1	B	28	ARG	3.2
1	A	233	ILE	3.1
1	A	237	ARG	3.1
1	A	222	TRP	2.6
1	A	21	GLN	2.6
1	A	203	SER	2.5
1	B	24	GLY	2.5
1	B	206	GLU	2.4
1	A	195	SER	2.3
1	B	245	LYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	VT0	B	401	25/25	0.91	0.10	75,86,96,99	25

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around VT0 B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.