



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:39 PM UTC

PDB ID : 7C5M / pdb_00007c5m
Title : Crystal Structure of C150A+H177A mutant of Glyceraldehyde-3-phosphate-dehydrogenase1 from Escherichia coli complexed with G3P at 1.8 Angstrom resolution
Authors : Zhang, L.; Liu, M.R.; Bao, L.Y.; Yao, Y.C.; Bostrom, I.K.; Wang, Y.D.; Chen, A.Q.; Li, J.X.; Gu, S.H.; Ji, C.N.
Deposited on : 2020-05-20
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	352	92% 5%
1	P	352	91% 5%
1	Q	352	91% 5%
1	R	352	88% 7% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	3PG	O	402	-	X	-	-
5	G3H	Q	402	-	X	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	O	334	Total	C	N	O	S	0	2	0
			2531	1599	425	501	6			
1	P	334	Total	C	N	O	S	0	2	1
			2528	1598	424	500	6			
1	Q	336	Total	C	N	O	S	0	2	1
			2538	1605	426	501	6			
1	R	333	Total	C	N	O	S	0	2	0
			2528	1599	426	497	6			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	-18	HIS	-	expression tag	UNP A0A140NCK4
O	-17	HIS	-	expression tag	UNP A0A140NCK4
O	-16	HIS	-	expression tag	UNP A0A140NCK4
O	-15	HIS	-	expression tag	UNP A0A140NCK4
O	-14	HIS	-	expression tag	UNP A0A140NCK4
O	-13	HIS	-	expression tag	UNP A0A140NCK4
O	-12	SER	-	expression tag	UNP A0A140NCK4
O	-11	SER	-	expression tag	UNP A0A140NCK4
O	-10	GLY	-	expression tag	UNP A0A140NCK4
O	-9	LEU	-	expression tag	UNP A0A140NCK4
O	-8	VAL	-	expression tag	UNP A0A140NCK4
O	-7	PRO	-	expression tag	UNP A0A140NCK4
O	-6	ARG	-	expression tag	UNP A0A140NCK4
O	-5	GLY	-	expression tag	UNP A0A140NCK4
O	-4	SER	-	expression tag	UNP A0A140NCK4
O	-3	HIS	-	expression tag	UNP A0A140NCK4
O	-2	MET	-	expression tag	UNP A0A140NCK4
O	-1	ALA	-	expression tag	UNP A0A140NCK4
O	0	SER	-	expression tag	UNP A0A140NCK4
O	150	ALA	CYS	engineered mutation	UNP A0A140NCK4
O	177	ALA	HIS	engineered mutation	UNP A0A140NCK4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
P	-18	HIS	-	expression tag	UNP A0A140NCK4
P	-17	HIS	-	expression tag	UNP A0A140NCK4
P	-16	HIS	-	expression tag	UNP A0A140NCK4
P	-15	HIS	-	expression tag	UNP A0A140NCK4
P	-14	HIS	-	expression tag	UNP A0A140NCK4
P	-13	HIS	-	expression tag	UNP A0A140NCK4
P	-12	SER	-	expression tag	UNP A0A140NCK4
P	-11	SER	-	expression tag	UNP A0A140NCK4
P	-10	GLY	-	expression tag	UNP A0A140NCK4
P	-9	LEU	-	expression tag	UNP A0A140NCK4
P	-8	VAL	-	expression tag	UNP A0A140NCK4
P	-7	PRO	-	expression tag	UNP A0A140NCK4
P	-6	ARG	-	expression tag	UNP A0A140NCK4
P	-5	GLY	-	expression tag	UNP A0A140NCK4
P	-4	SER	-	expression tag	UNP A0A140NCK4
P	-3	HIS	-	expression tag	UNP A0A140NCK4
P	-2	MET	-	expression tag	UNP A0A140NCK4
P	-1	ALA	-	expression tag	UNP A0A140NCK4
P	0	SER	-	expression tag	UNP A0A140NCK4
P	150	ALA	CYS	engineered mutation	UNP A0A140NCK4
P	177	ALA	HIS	engineered mutation	UNP A0A140NCK4
Q	-18	HIS	-	expression tag	UNP A0A140NCK4
Q	-17	HIS	-	expression tag	UNP A0A140NCK4
Q	-16	HIS	-	expression tag	UNP A0A140NCK4
Q	-15	HIS	-	expression tag	UNP A0A140NCK4
Q	-14	HIS	-	expression tag	UNP A0A140NCK4
Q	-13	HIS	-	expression tag	UNP A0A140NCK4
Q	-12	SER	-	expression tag	UNP A0A140NCK4
Q	-11	SER	-	expression tag	UNP A0A140NCK4
Q	-10	GLY	-	expression tag	UNP A0A140NCK4
Q	-9	LEU	-	expression tag	UNP A0A140NCK4
Q	-8	VAL	-	expression tag	UNP A0A140NCK4
Q	-7	PRO	-	expression tag	UNP A0A140NCK4
Q	-6	ARG	-	expression tag	UNP A0A140NCK4
Q	-5	GLY	-	expression tag	UNP A0A140NCK4
Q	-4	SER	-	expression tag	UNP A0A140NCK4
Q	-3	HIS	-	expression tag	UNP A0A140NCK4
Q	-2	MET	-	expression tag	UNP A0A140NCK4
Q	-1	ALA	-	expression tag	UNP A0A140NCK4
Q	0	SER	-	expression tag	UNP A0A140NCK4
Q	150	ALA	CYS	engineered mutation	UNP A0A140NCK4
Q	177	ALA	HIS	engineered mutation	UNP A0A140NCK4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
R	-18	HIS	-	expression tag	UNP A0A140NCK4
R	-17	HIS	-	expression tag	UNP A0A140NCK4
R	-16	HIS	-	expression tag	UNP A0A140NCK4
R	-15	HIS	-	expression tag	UNP A0A140NCK4
R	-14	HIS	-	expression tag	UNP A0A140NCK4
R	-13	HIS	-	expression tag	UNP A0A140NCK4
R	-12	SER	-	expression tag	UNP A0A140NCK4
R	-11	SER	-	expression tag	UNP A0A140NCK4
R	-10	GLY	-	expression tag	UNP A0A140NCK4
R	-9	LEU	-	expression tag	UNP A0A140NCK4
R	-8	VAL	-	expression tag	UNP A0A140NCK4
R	-7	PRO	-	expression tag	UNP A0A140NCK4
R	-6	ARG	-	expression tag	UNP A0A140NCK4
R	-5	GLY	-	expression tag	UNP A0A140NCK4
R	-4	SER	-	expression tag	UNP A0A140NCK4
R	-3	HIS	-	expression tag	UNP A0A140NCK4
R	-2	MET	-	expression tag	UNP A0A140NCK4
R	-1	ALA	-	expression tag	UNP A0A140NCK4
R	0	SER	-	expression tag	UNP A0A140NCK4
R	150	ALA	CYS	engineered mutation	UNP A0A140NCK4
R	177	ALA	HIS	engineered mutation	UNP A0A140NCK4

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	O	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	P	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	Q	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	R	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is 3-PHOSPHOGLYCERIC ACID (CCD ID: 3PG) (formula: $C_3H_7O_7P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	O	1	Total	C	O	P	0	0
			11	3	7	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	O	1	Total	O	P	0	0
			5	4	1		
4	O	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is GLYCERALDEHYDE-3-PHOSPHATE (CCD ID: G3H) (formula: $C_3H_7O_6P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	P	1	Total	C	O	P	0	0
			10	3	6	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	Q	1	Total	C	O	P	0	0
			10	3	6	1		
5	R	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	P	1	Total	C	O	0	1
			8	4	4		
6	Q	1	Total	C	O	0	0
			4	2	2		
6	Q	1	Total	C	O	0	1
			8	4	4		
6	R	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	O	419	Total	O	0	0
			419	419		
7	P	388	Total	O	0	0
			388	388		
7	Q	375	Total	O	0	0
			375	375		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	R	292	Total 292	O 292	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase

Chain O:  92% 5%




- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase

Chain P:  91% 5%




- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase

Chain Q:  91% 5%



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase

Chain R:  88% 7% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.92Å 89.92Å 342.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 1.80 45.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.00-1.80) 99.9 (45.00-1.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.145 , 0.189 (Not available) , 0.195	Depositor DCC
R_{free} test set	6457 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11850	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G3H, EDO, NAD, 3PG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	O	1.25	2/2571 (0.1%)	1.11	3/3489 (0.1%)
1	P	1.21	0/2568	1.10	6/3484 (0.2%)
1	Q	1.23	3/2578 (0.1%)	1.11	1/3498 (0.0%)
1	R	1.25	2/2569 (0.1%)	1.14	3/3486 (0.1%)
All	All	1.24	7/10286 (0.1%)	1.12	13/13957 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	171	GLY	N-CA	5.70	1.51	1.45
1	Q	75	TYR	N-CA	5.54	1.52	1.45
1	R	29	VAL	N-CA	-5.24	1.40	1.46
1	Q	13	GLY	N-CA	5.17	1.52	1.45
1	Q	325	ARG	CA-CB	-5.03	1.45	1.53

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	18	ARG	NE-CZ-NH2	6.89	125.41	119.20
1	O	268	GLU	CB-CG-CD	6.03	122.85	112.60
1	P	268	GLU	CB-CG-CD	5.97	122.75	112.60
1	P	83	ILE	N-CA-CB	5.82	115.49	110.08
1	O	317	TYR	N-CA-C	5.78	117.58	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	2531	0	2556	4	0
1	P	2528	0	2551	8	0
1	Q	2538	0	2566	10	0
1	R	2528	0	2556	15	0
2	O	44	0	26	2	0
2	P	44	0	26	2	0
2	Q	44	0	26	1	0
2	R	44	0	26	1	0
3	O	11	0	4	1	0
4	O	10	0	0	0	0
5	P	10	0	5	1	0
5	Q	10	0	5	2	0
5	R	10	0	5	2	0
6	P	8	0	12	3	0
6	Q	12	0	18	1	0
6	R	4	0	6	0	0
7	O	419	0	0	1	1
7	P	388	0	0	4	0
7	Q	375	0	0	3	0
7	R	292	0	0	1	2
All	All	11850	0	10388	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:Q:402:G3H:H31	7:Q:557:HOH:O	1.53	1.04
1:P:182:THR:OG1	5:P:402:G3H:H32	1.95	0.66
1:Q:229:HIS:HD2	7:Q:814:HOH:O	1.87	0.58
1:R:323:LEU:C	1:R:323:LEU:HD23	2.28	0.58
1:P:323:LEU:C	1:P:323:LEU:HD23	2.29	0.57

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:O:803:HOH:O	7:R:629:HOH:O[1_655]	1.96	0.24
7:R:532:HOH:O	7:R:532:HOH:O[7_465]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	334/352 (95%)	322 (96%)	12 (4%)	0	100	100
1	P	334/352 (95%)	322 (96%)	12 (4%)	0	100	100
1	Q	336/352 (96%)	325 (97%)	11 (3%)	0	100	100
1	R	333/352 (95%)	320 (96%)	13 (4%)	0	100	100
All	All	1337/1408 (95%)	1289 (96%)	48 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	O	274/287 (96%)	274 (100%)	0	100	100
1	P	273/287 (95%)	272 (100%)	1 (0%)	84	83
1	Q	274/287 (96%)	272 (99%)	2 (1%)	76	73
1	R	273/287 (95%)	270 (99%)	3 (1%)	65	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1094/1148 (95%)	1088 (100%)	6 (0%)	81 80

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	46	LYS
1	R	229[A]	HIS
1	R	229[B]	HIS
1	Q	4	VAL
1	P	209	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	R	153	ASN
1	Q	257	ASN
1	Q	163	HIS
1	Q	153	ASN
1	Q	229	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	3PG	O	402	-	9,10,10	1.84	3 (33%)	11,14,14	1.51	2 (18%)
4	PO4	O	404	-	4,4,4	0.75	0	6,6,6	1.01	0
6	EDO	P	403[B]	-	3,3,3	0.48	0	2,2,2	0.26	0
4	PO4	O	403	-	4,4,4	0.92	0	6,6,6	2.29	3 (50%)
5	G3H	P	402	-	8,9,9	2.35	2 (25%)	7,12,12	1.43	1 (14%)
2	NAD	P	401	-	46,48,48	1.61	11 (23%)	64,73,73	1.67	12 (18%)
2	NAD	R	401	-	46,48,48	1.76	7 (15%)	64,73,73	1.82	17 (26%)
6	EDO	Q	403	-	3,3,3	0.44	0	2,2,2	0.51	0
5	G3H	Q	402	-	8,9,9	2.34	4 (50%)	7,12,12	2.82	5 (71%)
6	EDO	P	403[A]	-	3,3,3	0.52	0	2,2,2	0.52	0
2	NAD	Q	401	-	46,48,48	1.50	8 (17%)	64,73,73	1.79	14 (21%)
6	EDO	Q	404[A]	-	3,3,3	0.44	0	2,2,2	0.36	0
6	EDO	R	403	-	3,3,3	0.47	0	2,2,2	0.83	0
6	EDO	Q	404[B]	-	3,3,3	0.41	0	2,2,2	0.73	0
5	G3H	R	402	-	8,9,9	1.91	2 (25%)	7,12,12	1.77	2 (28%)
2	NAD	O	401	-	46,48,48	2.14	8 (17%)	64,73,73	1.77	17 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3PG	O	402	-	-	10/10/10/10	-
6	EDO	P	403[B]	-	-	1/1/1/1	-
5	G3H	P	402	-	-	7/7/8/8	-
2	NAD	P	401	-	-	4/30/62/62	0/5/5/5
2	NAD	R	401	-	-	5/30/62/62	0/5/5/5
6	EDO	Q	403	-	-	1/1/1/1	-
5	G3H	Q	402	-	-	4/7/8/8	-
6	EDO	P	403[A]	-	-	0/1/1/1	-
2	NAD	Q	401	-	-	5/30/62/62	0/5/5/5
6	EDO	Q	404[A]	-	-	1/1/1/1	-
6	EDO	R	403	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	Q	404[B]	-	-	1/1/1/1	-
5	G3H	R	402	-	-	6/7/8/8	-
2	NAD	O	401	-	-	4/30/62/62	0/5/5/5

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	O	401	NAD	PA-O3	11.13	1.71	1.59
2	R	401	NAD	PA-O3	5.50	1.65	1.59
5	P	402	G3H	C3-C2	5.28	1.59	1.51
2	P	401	NAD	C5A-C4A	5.00	1.48	1.39
2	P	401	NAD	PA-O3	4.87	1.64	1.59

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	401	NAD	N3A-C2A-N1A	-6.34	118.98	128.58
5	Q	402	G3H	O2-C2-C1	5.22	120.26	109.03
2	O	401	NAD	N3A-C4A-N9A	4.83	135.38	127.17
2	P	401	NAD	N3A-C2A-N1A	-4.61	121.60	128.58
2	P	401	NAD	N3A-C4A-N9A	4.36	134.58	127.17

There are no chirality outliers.

5 of 50 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	O	401	NAD	O4D-C1D-N1N-C2N
2	O	401	NAD	O4D-C1D-N1N-C6N
2	O	401	NAD	C2D-C1D-N1N-C2N
2	O	401	NAD	C2D-C1D-N1N-C6N
2	P	401	NAD	O4D-C1D-N1N-C2N

There are no ring outliers.

10 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	O	402	3PG	1	0
6	P	403[B]	EDO	3	0
5	P	402	G3H	1	0
2	P	401	NAD	2	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	R	401	NAD	1	0
5	Q	402	G3H	2	0
2	Q	401	NAD	1	0
6	Q	404[A]	EDO	1	0
5	R	402	G3H	2	0
2	O	401	NAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














Ligand G3H P 402








Ligand NAD P 401














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	O	334/352 (94%)	-0.57	0 100 100	12, 20, 32, 47	2 (0%)
1	P	334/352 (94%)	-0.55	1 (0%) 90 90	12, 21, 35, 60	2 (0%)
1	Q	336/352 (95%)	-0.40	5 (1%) 72 72	10, 22, 39, 56	2 (0%)
1	R	333/352 (94%)	0.19	11 (3%) 49 49	11, 29, 50, 62	2 (0%)
All	All	1337/1408 (94%)	-0.33	17 (1%) 75 75	10, 22, 43, 62	8 (0%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	5	GLY	2.8
1	R	90	ALA	2.7
1	R	333	LEU	2.6
1	Q	1	MET	2.6
1	Q	-1	ALA	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	Q	403	4/4	0.72	0.15	54,56,56,59	0
6	EDO	R	403	4/4	0.77	0.17	45,46,52,53	0
4	PO4	O	404	5/5	0.81	0.17	58,64,65,98	0
6	EDO	P	403[A]	4/4	0.84	0.14	25,32,32,36	4
6	EDO	P	403[B]	4/4	0.84	0.14	23,24,25,26	4
6	EDO	Q	404[B]	4/4	0.87	0.15	24,28,29,32	4
6	EDO	Q	404[A]	4/4	0.87	0.15	26,28,28,28	4
5	G3H	R	402	10/10	0.90	0.12	34,42,53,59	0
3	3PG	O	402	11/11	0.93	0.11	28,33,57,66	0
5	G3H	P	402	10/10	0.94	0.10	28,33,47,49	0
4	PO4	O	403	5/5	0.94	0.12	32,35,37,41	0
2	NAD	R	401	44/44	0.95	0.08	25,31,34,36	0
2	NAD	O	401	44/44	0.97	0.06	18,20,23,27	0
2	NAD	P	401	44/44	0.97	0.06	18,22,25,27	0
5	G3H	Q	402	10/10	0.98	0.05	21,27,31,33	0
2	NAD	Q	401	44/44	0.98	0.04	16,21,23,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around EDO Q 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO R 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 O 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO P 403 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO P 403 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO Q 404 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EDO Q 404 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around G3H R 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3PG O 402:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around G3H P 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 O 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD R 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD O 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAD P 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around G3H Q 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.