



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 03:28 AM UTC

PDB ID : 8C5V / pdb_00008c5v
EMDB ID : EMD-15641
Title : Chemotaxis core signalling unit from E protein lysed E. coli cells
Authors : Cassidy, C.K.; Qin, Z.; Zhang, P.
Deposited on : 2023-01-10
Resolution : 12.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 12.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	93 (11.50 - 12.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	
1	B	391	
2	C	131	
2	D	131	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	143	41% 91% 9%
3	F	143	43% 84% 16%
3	G	143	34% 88% 12%
3	H	143	38% 90% 10%
4	I	516	28% 89% 11%
4	J	516	27% 87% 13%
4	K	516	26% 90% 10%
4	L	516	34% 86% 14%
4	M	516	30% 89% 11%
4	N	516	35% 88% 12%
4	O	516	30% 88% 12%
4	P	516	28% 89% 11%
4	Q	516	27% 88% 12%
4	R	516	33% 86% 14%
4	S	516	29% 89% 11%
4	T	516	35% 89% 11%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 59156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chemotaxis protein CheA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	391	Total	C	N	O	S	0	0
			2972	1851	532	576	13		
1	B	391	Total	C	N	O	S	0	0
			2972	1851	532	576	13		

- Molecule 2 is a protein called Chemotaxis protein CheA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	131	Total	C	N	O	S	0	0
			1042	653	169	213	7		
2	D	131	Total	C	N	O	S	0	0
			1042	653	169	213	7		

- Molecule 3 is a protein called Chemotaxis protein CheW.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	143	Total	C	N	O	S	0	0
			1103	704	181	216	2		
3	F	143	Total	C	N	O	S	0	0
			1103	704	181	216	2		
3	G	143	Total	C	N	O	S	0	0
			1103	704	181	216	2		
3	H	143	Total	C	N	O	S	0	0
			1103	704	181	216	2		

- Molecule 4 is a protein called Methyl-accepting chemotaxis protein I.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	I	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	J	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
4	K	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	L	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	M	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	N	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	O	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	P	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	Q	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	R	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	S	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		
4	T	516	Total	C	N	O	S	0	0
			3893	2404	693	780	16		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chemotaxis protein CheA





- Molecule 1: Chemotaxis protein CheA









• Molecule 2: Chemotaxis protein CheA





• Molecule 2: Chemotaxis protein CheA




• Molecule 3: Chemotaxis protein CheW






• Molecule 3: Chemotaxis protein CheW










- Molecule 3: Chemotaxis protein CheW





- Molecule 3: Chemotaxis protein CheW





- Molecule 4: Methyl-accepting chemotaxis protein I





- Molecule 4: Methyl-accepting chemotaxis protein I




• Molecule 4: Methyl-accepting chemotaxis protein I





• Molecule 4: Methyl-accepting chemotaxis protein I








• Molecule 4: Methyl-accepting chemotaxis protein I





• Molecule 4: Methyl-accepting chemotaxis protein I








• Molecule 4: Methyl-accepting chemotaxis protein I





• Molecule 4: Methyl-accepting chemotaxis protein I









• Molecule 4: Methyl-accepting chemotaxis protein I





• Molecule 4: Methyl-accepting chemotaxis protein I



• Molecule 4: Methyl-accepting chemotaxis protein I



• Molecule 4: Methyl-accepting chemotaxis protein I



4 Experimental information

Property	Value	Source
EM reconstruction method	SUBTOMOGRAM AVERAGING	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of subtomograms used	5100	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; emClarity	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	100	Depositor
Minimum defocus (nm)	3000	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.466	Depositor
Minimum map value	-0.299	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.041	Depositor
Recommended contour level	0.09	Depositor
Map size (Å)	527.04, 527.04, 527.04	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.745, 2.745, 2.745	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/3002	0.18	0/4058
1	B	0.07	0/3002	0.18	0/4058
2	C	0.08	0/1058	0.16	0/1426
2	D	0.07	0/1058	0.17	0/1426
3	E	0.10	0/1115	0.20	0/1515
3	F	0.09	0/1115	0.19	0/1515
3	G	0.10	0/1115	0.20	0/1515
3	H	0.08	0/1115	0.17	0/1515
4	I	0.07	0/3931	0.17	0/5316
4	J	0.06	0/3931	0.14	0/5316
4	K	0.10	0/3931	0.21	1/5316 (0.0%)
4	L	0.06	0/3931	0.14	0/5316
4	M	0.08	0/3931	0.19	2/5316 (0.0%)
4	N	0.06	0/3931	0.15	0/5316
4	O	0.07	0/3931	0.17	0/5316
4	P	0.06	0/3931	0.15	0/5316
4	Q	0.06	0/3931	0.15	0/5316
4	R	0.06	0/3931	0.15	0/5316
4	S	0.06	0/3931	0.15	0/5316
4	T	0.06	0/3931	0.15	0/5316
All	All	0.07	0/59752	0.17	3/80820 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
3	E	0	1
3	H	0	1
All	All	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	M	213	GLY	O-C-N	-7.46	114.95	122.19
4	K	441	THR	OG1-CB-CG2	5.84	120.99	109.30
4	M	213	GLY	CA-C-O	5.26	126.23	120.66

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	312	ARG	Sidechain
3	E	29	TYR	Sidechain
3	H	29	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2972	0	3064	24	0
1	B	2972	0	3064	34	0
2	C	1042	0	996	6	0
2	D	1042	0	996	5	0
3	E	1103	0	1133	7	0
3	F	1103	0	1133	16	0
3	G	1103	0	1133	10	0
3	H	1103	0	1133	8	0
4	I	3893	0	3891	31	0
4	J	3893	0	3891	38	0
4	K	3893	0	3891	33	0
4	L	3893	0	3891	44	0
4	M	3893	0	3891	35	0
4	N	3893	0	3891	39	0
4	O	3893	0	3891	37	0
4	P	3893	0	3891	33	0
4	Q	3893	0	3891	35	0
4	R	3893	0	3891	41	0
4	S	3893	0	3891	38	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	T	3893	0	3891	35	0
All	All	59156	0	59344	500	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 500 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:P:73:ARG:NH1	4:P:77:ASP:OD1	2.21	0.74
4:I:64:ARG:NH1	4:I:68:ASN:OD1	2.21	0.74
4:R:73:ARG:NH1	4:R:77:ASP:OD1	2.21	0.73
4:Q:73:ARG:NH1	4:Q:77:ASP:OD1	2.21	0.73
4:M:73:ARG:NH1	4:M:77:ASP:OD1	2.22	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/391 (100%)	356 (92%)	31 (8%)	2 (0%)	24	63
1	B	389/391 (100%)	359 (92%)	28 (7%)	2 (0%)	24	63
2	C	129/131 (98%)	127 (98%)	2 (2%)	0	100	100
2	D	129/131 (98%)	125 (97%)	4 (3%)	0	100	100
3	E	141/143 (99%)	127 (90%)	13 (9%)	1 (1%)	18	56
3	F	141/143 (99%)	122 (86%)	18 (13%)	1 (1%)	18	56
3	G	141/143 (99%)	127 (90%)	12 (8%)	2 (1%)	9	40
3	H	141/143 (99%)	125 (89%)	14 (10%)	2 (1%)	9	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	I	514/516 (100%)	499 (97%)	13 (2%)	2 (0%)	30	67
4	J	514/516 (100%)	507 (99%)	6 (1%)	1 (0%)	43	78
4	K	514/516 (100%)	505 (98%)	9 (2%)	0	100	100
4	L	514/516 (100%)	508 (99%)	6 (1%)	0	100	100
4	M	514/516 (100%)	509 (99%)	5 (1%)	0	100	100
4	N	514/516 (100%)	504 (98%)	8 (2%)	2 (0%)	30	67
4	O	514/516 (100%)	510 (99%)	4 (1%)	0	100	100
4	P	514/516 (100%)	508 (99%)	4 (1%)	2 (0%)	30	67
4	Q	514/516 (100%)	509 (99%)	2 (0%)	3 (1%)	21	59
4	R	514/516 (100%)	503 (98%)	11 (2%)	0	100	100
4	S	514/516 (100%)	504 (98%)	10 (2%)	0	100	100
4	T	514/516 (100%)	508 (99%)	6 (1%)	0	100	100
All	All	7768/7808 (100%)	7542 (97%)	206 (3%)	20 (0%)	37	72

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	614	TYR
4	I	146	GLY
3	F	94	GLY
4	J	218	LEU
4	Q	218	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/329 (100%)	328 (100%)	1 (0%)	86	86
1	B	329/329 (100%)	329 (100%)	0	100	100
2	C	111/111 (100%)	111 (100%)	0	100	100
2	D	111/111 (100%)	111 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	E	124/124 (100%)	124 (100%)	0	100	100
3	F	124/124 (100%)	124 (100%)	0	100	100
3	G	124/124 (100%)	124 (100%)	0	100	100
3	H	124/124 (100%)	124 (100%)	0	100	100
4	I	408/408 (100%)	407 (100%)	1 (0%)	87	87
4	J	408/408 (100%)	408 (100%)	0	100	100
4	K	408/408 (100%)	408 (100%)	0	100	100
4	L	408/408 (100%)	407 (100%)	1 (0%)	87	87
4	M	408/408 (100%)	408 (100%)	0	100	100
4	N	408/408 (100%)	408 (100%)	0	100	100
4	O	408/408 (100%)	408 (100%)	0	100	100
4	P	408/408 (100%)	408 (100%)	0	100	100
4	Q	408/408 (100%)	408 (100%)	0	100	100
4	R	408/408 (100%)	408 (100%)	0	100	100
4	S	408/408 (100%)	407 (100%)	1 (0%)	87	87
4	T	408/408 (100%)	408 (100%)	0	100	100
All	All	6272/6272 (100%)	6268 (100%)	4 (0%)	87	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	SER
4	I	1	MET
4	L	1	MET
4	S	43	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 78 such sidechains are listed below:

Mol	Chain	Res	Type
4	Q	504	GLN
4	T	48	GLN
4	R	39	ASN
4	S	31	ASN
4	T	231	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-15641. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 96




Y Index: 96




Z Index: 96


6.2.2 Raw map



X Index: 96



Y Index: 96




Z Index: 96


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 105




Y Index: 96




Z Index: 67


6.3.2 Raw map



X Index: 105



Y Index: 96




Z Index: 67


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 3413 nm³; this corresponds to an approximate mass of 3083 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.083 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.083 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	12.00	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	23.58	26.18	25.25

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 23.58 differs from the reported value 12.0 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-15641 and PDB model 8C5V. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.09 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.09).

9.4 Atom inclusion [i](#)



At the recommended contour level, 66% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.09) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6270	 0.0600
A	 0.5390	 0.0450
B	 0.5430	 0.0520
C	 0.1780	 -0.0050
D	 0.1920	 0.0030
E	 0.5670	 0.0490
F	 0.5500	 0.0460
G	 0.6440	 0.0520
H	 0.5790	 0.0340
I	 0.6930	 0.0700
J	 0.6790	 0.0640
K	 0.7080	 0.0670
L	 0.6170	 0.0590
M	 0.6620	 0.0740
N	 0.6050	 0.0600
O	 0.6850	 0.0690
P	 0.6770	 0.0660
Q	 0.7070	 0.0630
R	 0.6290	 0.0650
S	 0.6690	 0.0760
T	 0.6040	 0.0580

