



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:29 AM UTC

PDB ID : 2C67 / pdb_00002c67
Title : MAO inhibition by rasagiline analogues
Authors : Binda, C.; Hubalek, F.; Li, M.; Herzig, Y.; Sterling, J.; Edmondson, D.E.;
Mattevi, A.
Deposited on : 2005-11-07
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	8% 90% 5% . .
1	B	520	8% 87% 8% . 5%



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			11	10	1		
3	B	1	Total	C	N	0	0
			11	10	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	299	Total	O	0	0
			299	299		
4	B	328	Total	O	0	0
			328	328		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AMINE OXIDASE (FLAVIN-CONTAINING) B



• Molecule 1: AMINE OXIDASE (FLAVIN-CONTAINING) B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.27Å 222.45Å 86.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.70 15.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.7 (15.00-1.70) 97.5 (15.00-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.194 , 0.212 (Not available) , 0.214	Depositor DCC
R_{free} test set	3348 reflections (2.48%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.013 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8666	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, RM1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/4068	0.71	1/5522 (0.0%)
1	B	0.47	0/4037	0.71	0/5479
All	All	0.47	0/8105	0.71	1/11001 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	LYS	CA-CB-CG	5.60	125.29	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3971	0	3967	32	0
1	B	3940	0	3937	41	1
2	A	53	0	29	1	0
2	B	53	0	29	1	0
3	A	11	0	13	1	0
3	B	11	0	13	1	0
4	A	299	0	0	3	1
4	B	328	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8666	0	7988	60	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:ALA:HB1	1:B:280:MET:HE1	1.43	0.97
1:A:280:MET:HE1	1:B:353:ALA:HB1	1.48	0.95
1:A:28:LEU:HD11	1:A:454:MET:HE1	1.49	0.93
1:B:28:LEU:HD11	1:B:454:MET:HE1	1.50	0.91
1:A:251:ASN:HD22	1:A:251:ASN:H	1.32	0.76
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.34	0.75
1:B:251:ASN:HD22	1:B:251:ASN:H	1.33	0.74
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.37	0.72
1:A:353:ALA:CB	1:B:280:MET:HE1	2.21	0.70
1:B:49:GLN:NE2	4:B:2032:HOH:O	2.26	0.67
1:B:271:LYS:HE2	4:B:2264:HOH:O	2.02	0.58
1:B:309:LYS:HG3	4:B:2203:HOH:O	2.05	0.56
1:A:280:MET:HE1	1:B:353:ALA:CB	2.29	0.55
1:A:321:GLU:H	1:A:321:GLU:CD	2.14	0.55
1:A:350:ARG:HD3	1:B:410:TYR:OH	2.08	0.54
1:B:451:LEU:HD23	1:B:454:MET:CE	2.38	0.54
1:A:451:LEU:HD23	1:A:454:MET:CE	2.40	0.52
1:A:268:LEU:HD23	1:B:270:MET:HE1	1.92	0.52
1:A:248:GLU:OE2	1:B:252:HIS:NE2	2.43	0.51
1:A:251:ASN:H	1:A:251:ASN:ND2	2.07	0.51
1:B:321:GLU:H	1:B:321:GLU:CD	2.17	0.51
1:A:58:GLY:HA2	2:A:1502:FAD:C4X	2.42	0.50
1:A:451:LEU:HA	1:A:454:MET:HE2	1.94	0.50
1:A:270:MET:HE1	1:B:268:LEU:HD23	1.94	0.49
1:A:49:GLN:HB3	4:A:2026:HOH:O	2.11	0.49
1:B:163:GLN:HB3	1:B:318:ASP:OD2	2.12	0.49
1:B:451:LEU:HD23	1:B:454:MET:HE1	1.94	0.48
1:B:58:GLY:HA2	2:B:1497:FAD:C4X	2.44	0.48
1:A:233:ARG:HG3	1:A:251:ASN:HD21	1.79	0.48
1:B:309:LYS:NZ	1:B:373:GLY:O	2.47	0.47
1:B:233:ARG:HG3	1:B:251:ASN:HD21	1.80	0.47
1:B:251:ASN:H	1:B:251:ASN:ND2	2.08	0.47
1:A:280:MET:HG3	1:B:389:CYS:CB	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:451:LEU:HD23	1:A:454:MET:HE1	1.95	0.47
1:B:451:LEU:HA	1:B:454:MET:HE2	1.97	0.47
1:B:412:ARG:HG2	4:B:2275:HOH:O	2.14	0.46
1:A:410:TYR:OH	1:B:350:ARG:HD3	2.15	0.46
1:B:387:ASN:O	1:B:390:GLU:HG2	2.16	0.46
1:A:451:LEU:HA	1:A:454:MET:CE	2.45	0.46
1:A:252:HIS:NE2	1:B:248:GLU:OE2	2.45	0.45
1:A:389:CYS:CB	1:B:280:MET:HG3	2.47	0.45
1:B:126:GLY:O	1:B:190:LYS:HG3	2.17	0.45
1:A:163:GLN:HB3	1:A:318:ASP:OD2	2.17	0.44
1:B:117:ASN:HD22	1:B:120:ARG:NH2	2.10	0.44
1:B:451:LEU:HA	1:B:454:MET:CE	2.47	0.44
1:B:412:ARG:HD2	4:B:2274:HOH:O	2.17	0.43
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.10	0.43
1:B:398:TYR:CE2	3:B:1498:RM1:H9	2.54	0.43
1:A:398:TYR:CE2	3:A:1503:RM1:H9	2.55	0.42
1:A:93:LYS:HB2	4:A:2068:HOH:O	2.20	0.41
1:B:8:VAL:HG23	1:B:257:ALA:HB2	2.03	0.41
1:B:290:PRO:HG2	1:B:401:TYR:CZ	2.54	0.41
1:B:190:LYS:C	1:B:190:LYS:HD3	2.46	0.41
1:B:323:PRO:HD2	1:B:367:LEU:HD22	2.03	0.41
1:B:280:MET:HE3	1:B:280:MET:HB3	1.78	0.40
1:A:280:MET:HG3	1:B:389:CYS:HB3	2.03	0.40
1:A:412:ARG:HD2	4:A:2247:HOH:O	2.20	0.40
1:A:190:LYS:CD	1:A:190:LYS:C	2.95	0.40
1:A:290:PRO:HG2	1:A:401:TYR:CZ	2.56	0.40
1:B:237:TYR:HB3	1:B:248:GLU:HB3	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2027:HOH:O	4:A:2027:HOH:O[3_655]	1.40	0.80
1:B:25:ASP:O	1:B:49:GLN:NE2[4_565]	1.96	0.24

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/520 (96%)	486 (98%)	11 (2%)	0	100	100
1	B	492/520 (95%)	480 (98%)	12 (2%)	0	100	100
All	All	989/1040 (95%)	966 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/444 (96%)	416 (97%)	11 (3%)	40	23
1	B	424/444 (96%)	413 (97%)	11 (3%)	40	23
All	All	851/888 (96%)	829 (97%)	22 (3%)	40	23

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	190	LYS
1	A	209	LYS
1	A	251	ASN
1	A	280	MET
1	A	321	GLU
1	A	379	GLU
1	A	397	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	413	VAL
1	A	414	LEU
1	A	459	GLU
1	B	49	GLN
1	B	190	LYS
1	B	209	LYS
1	B	251	ASN
1	B	280	MET
1	B	321	GLU
1	B	379	GLU
1	B	397	CYS
1	B	413	VAL
1	B	414	LEU
1	B	459	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	251	ASN
1	A	464	GLN
1	B	117	ASN
1	B	251	ASN
1	B	452	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	B	1497	1	58,58,58	1.07	4 (6%)	85,89,89	1.55	16 (18%)
2	FAD	A	1502	1	58,58,58	1.06	4 (6%)	85,89,89	1.54	16 (18%)
3	RM1	B	1498	-	12,12,12	2.07	5 (41%)	14,16,16	1.64	3 (21%)
3	RM1	A	1503	-	12,12,12	2.16	5 (41%)	14,16,16	1.77	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	1497	1	-	0/34/50/50	0/6/6/6
2	FAD	A	1502	1	-	0/34/50/50	0/6/6/6
3	RM1	B	1498	-	-	0/0/11/11	0/2/2/2
3	RM1	A	1503	-	-	0/0/11/11	0/2/2/2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1503	RM1	C7-C8	-4.30	1.34	1.39
3	B	1498	RM1	C3-C8	-4.08	1.32	1.39
3	A	1503	RM1	C3-C8	-4.01	1.32	1.39
3	B	1498	RM1	C7-C8	-3.85	1.35	1.39
2	B	1497	FAD	C4X-N5	3.70	1.38	1.30
2	A	1502	FAD	C4X-N5	3.70	1.38	1.30
2	B	1497	FAD	C10-N1	2.95	1.39	1.33
2	A	1502	FAD	C10-N1	2.84	1.39	1.33
3	A	1503	RM1	C4-C3	-2.83	1.35	1.39
3	B	1498	RM1	C4-C3	-2.71	1.35	1.39
2	B	1497	FAD	C2A-N1A	2.57	1.38	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1502	FAD	C2A-N1A	2.48	1.38	1.33
2	A	1502	FAD	C8A-N7A	2.47	1.36	1.31
3	A	1503	RM1	C6-C7	-2.27	1.35	1.38
3	B	1498	RM1	C5-C4	-2.27	1.35	1.38
3	A	1503	RM1	C5-C4	-2.26	1.35	1.38
2	B	1497	FAD	C2A-N3A	2.22	1.37	1.33
3	B	1498	RM1	C6-C7	-2.09	1.35	1.38

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1497	FAD	N3A-C2A-N1A	-5.85	119.73	128.58
2	A	1502	FAD	N3A-C2A-N1A	-5.83	119.75	128.58
3	A	1503	RM1	C1-C9-N10	-4.36	106.50	115.35
2	A	1502	FAD	N9A-C8A-N7A	-4.05	108.19	113.94
2	B	1497	FAD	N9A-C8A-N7A	-3.97	108.31	113.94
3	B	1498	RM1	C1-C9-N10	-3.82	107.59	115.35
2	B	1497	FAD	C5A-N7A-C8A	3.38	108.76	103.45
2	B	1497	FAD	C5A-C4A-N3A	-3.36	122.08	126.72
2	B	1497	FAD	C9A-C5X-N5	-3.33	118.93	122.45
2	A	1502	FAD	C5A-C4A-N3A	-3.31	122.16	126.72
2	A	1502	FAD	C5A-N7A-C8A	3.19	108.46	103.45
2	A	1502	FAD	C9A-C5X-N5	-3.09	119.17	122.45
2	B	1497	FAD	C2A-N3A-C4A	3.01	119.17	111.83
2	A	1502	FAD	C2A-N3A-C4A	2.98	119.10	111.83
2	A	1502	FAD	C4-N3-C2	-2.90	120.50	125.64
2	B	1497	FAD	C4-N3-C2	-2.88	120.52	125.64
2	B	1497	FAD	C5'-C4'-C3'	-2.80	106.93	112.22
2	B	1497	FAD	C10-C4X-N5	-2.54	119.62	124.81
2	B	1497	FAD	C4A-C5A-N7A	-2.50	107.72	110.58
2	B	1497	FAD	C4X-C4-N3	2.50	119.61	113.25
2	A	1502	FAD	C4X-C4-N3	2.47	119.54	113.25
2	A	1502	FAD	C4X-C10-N10	2.43	119.96	116.48
2	A	1502	FAD	C4A-C5A-N7A	-2.42	107.82	110.58
2	A	1502	FAD	C5'-C4'-C3'	-2.40	107.68	112.22
2	A	1502	FAD	C10-C4X-N5	-2.40	119.91	124.81
2	A	1502	FAD	C9A-N10-C10	-2.39	117.11	120.75
2	B	1497	FAD	C4X-C10-N10	2.31	119.79	116.48
2	B	1497	FAD	O4-C4-C4X	-2.27	120.55	126.53
2	A	1502	FAD	C4A-N9A-C8A	2.24	108.09	105.74
3	B	1498	RM1	C6-C7-C8	-2.22	118.33	120.99
3	B	1498	RM1	C7-C8-C9	-2.16	126.20	129.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1503	RM1	C10-N10-C9	2.15	120.13	113.70
2	B	1497	FAD	C9A-N10-C10	-2.14	117.49	120.75
2	B	1497	FAD	C4A-N9A-C8A	2.14	107.98	105.74
2	A	1502	FAD	O4-C4-C4X	-2.08	121.03	126.53
2	A	1502	FAD	C2A-N1A-C6A	2.07	122.14	118.73
2	B	1497	FAD	N3A-C4A-N9A	2.07	130.68	127.17
3	A	1503	RM1	C6-C7-C8	-2.05	118.53	120.99

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1497	FAD	1	0
2	A	1502	FAD	1	0
3	B	1498	RM1	1	0
3	A	1503	RM1	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	499/520 (95%)	0.07	43 (8%)	16 17	6, 12, 29, 52	0
1	B	494/520 (95%)	0.06	40 (8%)	18 19	6, 12, 27, 46	0
All	All	993/1040 (95%)	0.06	83 (8%)	17 18	6, 12, 29, 52	0

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	495	LEU	6.3
1	B	496	ILE	6.2
1	A	501	ILE	5.8
1	A	495	LEU	5.0
1	B	494	ARG	4.6
1	A	103	PHE	4.5
1	B	103	PHE	4.4
1	B	492	LEU	4.2
1	A	493	LEU	4.0
1	B	252	HIS	4.0
1	A	481	PHE	3.9
1	A	500	THR	3.8
1	B	49	GLN	3.8
1	B	107	TRP	3.8
1	B	318	ASP	3.7
1	A	252	HIS	3.6
1	A	319	GLY	3.6
1	B	493	LEU	3.6
1	A	498	LEU	3.6
1	A	107	TRP	3.5
1	A	496	ILE	3.5
1	A	254	MET	3.4
1	A	480	THR	3.3
1	B	491	GLY	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	494	ARG	3.2
1	A	489	VAL	3.2
1	A	491	GLY	3.2
1	B	243	GLU	3.2
1	B	489	VAL	3.0
1	A	492	LEU	3.0
1	A	490	PRO	3.0
1	A	321	GLU	3.0
1	B	3	ASN	3.0
1	A	486	LEU	2.9
1	A	484	ARG	2.9
1	B	479	THR	2.9
1	A	482	LEU	2.9
1	A	3	ASN	2.9
1	A	354	ARG	2.8
1	B	481	PHE	2.8
1	B	92	VAL	2.8
1	A	49	GLN	2.8
1	B	482	LEU	2.8
1	A	477	ILE	2.8
1	A	478	THR	2.7
1	B	242	ARG	2.7
1	A	370	LYS	2.6
1	B	490	PRO	2.6
1	B	119	TRP	2.5
1	A	102	PRO	2.5
1	A	499	THR	2.5
1	A	487	PRO	2.5
1	B	486	LEU	2.4
1	B	475	GLN	2.4
1	A	483	GLU	2.4
1	B	485	HIS	2.4
1	A	101	GLY	2.3
1	B	414	LEU	2.3
1	B	93	LYS	2.3
1	B	224	LEU	2.3
1	B	244	ASN	2.3
1	B	157	TRP	2.3
1	B	319	GLY	2.3
1	A	95	LYS	2.3
1	B	476	PRO	2.3
1	B	379	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	254	MET	2.3
1	B	163	GLN	2.2
1	B	354	ARG	2.2
1	A	227	ASP	2.2
1	B	321	GLU	2.2
1	A	100	ARG	2.2
1	B	112	TYR	2.2
1	A	479	THR	2.2
1	A	157	TRP	2.2
1	A	475	GLN	2.2
1	A	93	LYS	2.1
1	B	26	SER	2.1
1	B	484	ARG	2.1
1	A	106	VAL	2.1
1	B	110	ILE	2.1
1	A	497	GLY	2.1
1	A	242	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.


6.4 Ligands ⓘ


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	RM1	B	1498	11/11	0.90	0.08	12,13,13,14	0
3	RM1	A	1503	11/11	0.93	0.08	12,13,14,14	0
2	FAD	A	1502	53/53	0.98	0.04	6,7,8,9	0
2	FAD	B	1497	53/53	0.98	0.04	6,7,8,9	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.