



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:12 PM UTC

PDB ID : 2C6E / pdb_00002c6e
Title : Aurora A kinase activated mutant (T287D) in complex with a 5- aminopyrimidinyl quinazoline inhibitor
Authors : Paupit, R.A.; Pannifer, A.D.; Breed, J.; McMiken, H.H.J.; Rowsell, S.; Anderson, M.
Deposited on : 2005-11-09
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	
1	B	283	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4197 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE/THREONINE-PROTEIN KINASE 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	0	0
			2040	1317	352	366	5			
1	B	246	Total	C	N	O	S	0	0	0
			1977	1275	339	358	5			

- Molecule 2 is N-{5-[(7-[(2S)-2-HYDROXY-3-PIPERIDIN-1-YLPROPYL]OXY}-6-METHOXYQUINAZOLIN-4-YL)AMINO]PYRIMIDIN-2-YL}BENZAMIDE (CCD ID: HPM) (formula: C₂₈H₃₁N₇O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			39	28	7	4		
2	B	1	Total	C	N	O	0	0
			39	28	7	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	56	Total 56	O 56	0	0
3	B	46	Total 46	O 46	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SERINE/THREONINE-PROTEIN KINASE 6



• Molecule 1: SERINE/THREONINE-PROTEIN KINASE 6



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.60Å 88.42Å 67.83Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	53.00 – 2.10 53.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (53.00-2.10) 72.5 (53.00-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.10Å)	Xtriage
Refinement program	CNX 2000	Depositor
R, R_{free}	0.200 , 0.270 0.220 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.245	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.038 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4197	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HPM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2091	1.07	13/2829 (0.5%)
1	B	0.46	1/2025 (0.0%)	0.97	11/2743 (0.4%)
All	All	0.46	1/4116 (0.0%)	1.02	24/5572 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	251	VAL	C-N	-7.05	1.23	1.33

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	VAL	O-C-N	-16.03	100.18	122.62
1	B	175	HIS	N-CA-C	-11.95	98.72	113.18
1	B	251	VAL	O-C-N	-11.14	108.29	122.88
1	A	175	HIS	N-CA-C	-10.59	100.37	113.18
1	A	251	VAL	CA-C-N	10.48	136.19	123.19
1	A	251	VAL	C-N-CA	10.48	136.19	123.19
1	B	148	LEU	N-CA-C	-7.45	100.00	110.50
1	A	325	LYS	CA-C-N	6.91	124.70	119.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	325	LYS	C-N-CA	6.91	124.70	119.66
1	B	326	PRO	CA-C-N	6.38	126.30	119.28
1	B	326	PRO	C-N-CA	6.38	126.30	119.28
1	A	212	ALA	CA-C-N	6.25	125.98	119.05
1	A	212	ALA	C-N-CA	6.25	125.98	119.05
1	A	171	ALA	N-CA-C	-6.18	105.60	113.01
1	B	177	LEU	N-CA-C	-6.03	104.83	111.82
1	A	253	HIS	N-CA-C	6.01	118.99	111.24
1	A	148	LEU	N-CA-C	-5.97	100.77	110.20
1	A	133	GLU	N-CA-C	-5.47	100.26	109.07
1	B	209	LEU	N-CA-C	5.31	118.24	109.85
1	A	327	PRO	N-CA-C	5.22	120.39	113.86
1	B	213	PRO	N-CA-C	5.22	121.38	113.81
1	B	127	TRP	N-CA-C	5.12	118.22	110.64
1	B	152	LYS	N-CA-C	5.07	116.49	111.07
1	B	259	GLU	N-CA-C	-5.03	106.66	112.89

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	251	VAL	Mainchain
1	B	251	VAL	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2040	0	2017	43	0
1	B	1977	0	1942	60	0
2	A	39	0	31	2	0
2	B	39	0	31	1	0
3	A	56	0	0	2	0
3	B	46	0	0	2	0
All	All	4197	0	4021	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 13.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:LEU:HD22	3:A:2035:HOH:O	1.57	1.02
1:B:169:GLU:HA	1:B:173:VAL:HA	1.51	0.92
1:B:272:ALA:O	1:B:273:ASP:HB2	1.79	0.82
1:B:189:HIS:HE1	1:B:191:ASN:HD22	1.33	0.77
1:B:150:ARG:HH11	1:B:155:LYS:HZ3	1.33	0.74
1:B:184:GLN:HG2	2:B:1388:HPM:H32	1.71	0.71
1:B:165:LYS:O	1:B:169:GLU:HG2	1.90	0.70
1:B:150:ARG:HH11	1:B:155:LYS:NZ	1.90	0.69
1:A:252:ILE:HD12	1:A:252:ILE:N	2.08	0.69
1:B:258:PRO:HD2	1:B:259:GLU:OE2	1.94	0.68
1:B:257:LYS:HB3	1:B:259:GLU:OE2	1.95	0.67
1:A:168:LEU:O	1:A:174:GLU:HG2	1.96	0.66
1:B:143:PHE:CD2	1:B:168:LEU:HD23	2.30	0.66
1:A:219:ARG:O	1:A:223:LYS:HD3	1.96	0.66
1:B:200:HIS:HB2	1:B:204:ARG:O	1.95	0.65
1:B:257:LYS:HD3	1:B:290:GLY:HA3	1.79	0.65
1:A:315:GLY:HA3	1:A:363:LEU:HD21	1.78	0.65
1:B:366:ASN:HD22	1:B:369:GLN:HG3	1.63	0.64
1:B:253:HIS:O	1:B:254:ARG:HB2	1.95	0.64
1:A:155:LYS:HG2	1:A:155:LYS:O	1.98	0.63
1:B:366:ASN:HB3	1:B:369:GLN:HE21	1.63	0.63
1:A:180:GLU:HG2	1:A:276:TRP:HB2	1.79	0.63
1:B:366:ASN:ND2	1:B:369:GLN:HG3	2.16	0.61
1:A:183:ILE:O	1:A:187:LEU:HD13	2.01	0.61
1:B:357:ASP:O	1:B:361:ARG:HG3	2.02	0.60
1:A:325:LYS:HB2	1:A:326:PRO:HD2	1.83	0.60
1:B:326:PRO:HG2	1:B:329:GLU:HB2	1.84	0.60
1:B:143:PHE:HA	1:B:167:GLN:HG2	1.84	0.59
1:A:247:HIS:CE1	1:A:307:GLU:HB2	2.38	0.59
1:A:254:ARG:HG2	1:A:309:VAL:HG21	1.85	0.57
1:B:143:PHE:CD2	1:B:167:GLN:HG3	2.40	0.57
1:B:352:THR:HG22	1:B:354:GLY:H	1.71	0.55
1:A:167:GLN:O	1:A:170:LYS:HG3	2.06	0.55
1:B:382:ILE:HG22	1:B:387:SER:HB2	1.88	0.55
1:A:326:PRO:HG2	1:A:329:GLU:HB2	1.88	0.54
1:B:335:GLU:HA	1:B:335:GLU:OE2	2.08	0.54
1:A:253:HIS:O	1:A:254:ARG:HB2	2.07	0.54
1:B:177:LEU:C	1:B:177:LEU:HD13	2.33	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ILE:O	1:A:256:ILE:HG23	2.06	0.54
1:B:172:GLY:O	1:B:173:VAL:C	2.51	0.53
1:B:366:ASN:HD22	1:B:369:GLN:HE21	1.56	0.53
1:B:169:GLU:HA	1:B:173:VAL:CA	2.34	0.53
1:B:155:LYS:O	1:B:155:LYS:HD3	2.09	0.53
1:A:138:LEU:HD11	1:A:148:LEU:HB2	1.92	0.52
1:B:179:ARG:HH11	1:B:179:ARG:HG2	1.74	0.52
1:B:256:ILE:HG23	1:B:256:ILE:O	2.09	0.52
1:B:237:THR:HG21	1:B:387:SER:C	2.35	0.52
1:A:153:GLN:NE2	1:A:153:GLN:HA	2.25	0.52
1:A:153:GLN:HA	1:A:153:GLN:HE21	1.74	0.52
1:A:352:THR:HG22	1:A:354:GLY:H	1.75	0.51
1:B:174:GLU:HG3	1:B:176:GLN:HG3	1.93	0.51
1:B:326:PRO:HB2	3:B:2031:HOH:O	2.11	0.51
1:A:150:ARG:HD2	1:A:155:LYS:HG3	1.92	0.50
1:A:254:ARG:CZ	1:A:305:HIS:NE2	2.74	0.50
1:B:259:GLU:OE2	1:B:259:GLU:N	2.35	0.50
1:B:165:LYS:NZ	1:B:200:HIS:HD2	2.10	0.50
1:A:366:ASN:HD22	1:A:366:ASN:C	2.19	0.50
1:B:366:ASN:HB3	1:B:369:GLN:NE2	2.27	0.49
1:B:385:ASN:O	1:B:386:SER:HB3	2.11	0.49
1:B:352:THR:HG23	3:B:2038:HOH:O	2.13	0.49
1:A:144:GLY:HA3	1:A:162:VAL:O	2.13	0.49
1:B:155:LYS:HZ1	1:B:157:ILE:HD11	1.76	0.48
1:B:255:ASP:OD1	1:B:290:GLY:N	2.45	0.48
1:A:154:SER:O	1:A:155:LYS:HB3	2.13	0.48
1:B:150:ARG:HD3	1:B:155:LYS:NZ	2.29	0.47
1:A:254:ARG:CG	1:A:309:VAL:HG21	2.44	0.47
1:A:189:HIS:CD2	1:A:191:ASN:H	2.32	0.47
1:B:304:MET:SD	1:B:305:HIS:N	2.81	0.47
1:B:189:HIS:CE1	1:B:191:ASN:HD22	2.23	0.46
1:A:251:VAL:C	1:A:252:ILE:HD12	2.40	0.46
1:A:359:ILE:O	1:A:363:LEU:HG	2.15	0.46
1:A:257:LYS:NZ	1:A:290:GLY:HA2	2.31	0.46
1:B:200:HIS:HB3	1:B:205:VAL:HA	1.98	0.45
1:A:315:GLY:HA3	1:A:363:LEU:CD2	2.46	0.45
1:B:257:LYS:NZ	1:B:260:ASN:HD21	2.14	0.45
1:B:332:THR:HG23	1:B:335:GLU:HB2	1.98	0.45
1:B:177:LEU:O	1:B:181:VAL:HG23	2.17	0.45
1:A:316:VAL:HG23	3:A:2035:HOH:O	2.15	0.45
1:B:253:HIS:O	1:B:254:ARG:CB	2.65	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:GLU:HA	1:A:174:GLU:OE1	2.16	0.45
1:A:189:HIS:HD2	1:A:191:ASN:H	1.64	0.45
1:A:252:ILE:N	1:A:252:ILE:CD1	2.77	0.45
1:A:184:GLN:HG2	2:A:1389:HPM:H32	2.00	0.44
1:B:216:THR:HG22	1:B:262:LEU:CD2	2.48	0.43
1:B:184:GLN:HB3	1:B:195:LEU:HD12	2.00	0.43
1:A:155:LYS:HB3	1:A:155:LYS:NZ	2.33	0.43
1:B:385:ASN:O	1:B:386:SER:CB	2.66	0.43
1:A:150:ARG:HD2	1:A:155:LYS:CG	2.49	0.43
1:B:257:LYS:HB2	1:B:260:ASN:ND2	2.34	0.43
1:B:216:THR:HG22	1:B:262:LEU:HD23	2.01	0.43
1:B:376:VAL:O	1:B:382:ILE:HD12	2.19	0.42
1:A:170:LYS:C	1:A:172:GLY:N	2.78	0.42
1:B:352:THR:HG22	1:B:353:GLU:N	2.34	0.42
1:B:155:LYS:O	1:B:155:LYS:CD	2.67	0.42
1:B:244:SER:HA	1:B:373:LEU:HD12	2.02	0.42
1:A:214:LEU:HD12	1:A:214:LEU:N	2.35	0.41
1:B:237:THR:OG1	1:B:387:SER:HA	2.20	0.41
1:A:366:ASN:C	1:A:366:ASN:ND2	2.78	0.41
1:B:266:ALA:N	1:B:268:GLU:OE2	2.51	0.41
1:A:153:GLN:HE21	1:A:153:GLN:CA	2.33	0.41
1:A:224:LEU:O	1:A:225:SER:CB	2.69	0.41
1:A:363:LEU:HD23	1:A:363:LEU:N	2.36	0.41
2:A:1389:HPM:H5C2	2:A:1389:HPM:H8	1.95	0.40
1:B:366:ASN:HA	1:B:367:PRO:HD2	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	247/283 (87%)	235 (95%)	10 (4%)	2 (1%)	16 12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	242/283 (86%)	222 (92%)	11 (4%)	9 (4%)	2	1
All	All	489/566 (86%)	457 (94%)	21 (4%)	11 (2%)	5	2

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	256	ILE
1	B	173	VAL
1	A	174	GLU
1	B	255	ASP
1	B	256	ILE
1	B	303	ARG
1	B	253	HIS
1	B	386	SER
1	B	174	GLU
1	B	254	ARG
1	B	172	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/250 (86%)	207 (96%)	8 (4%)	30	33
1	B	207/250 (83%)	195 (94%)	12 (6%)	18	16
All	All	422/500 (84%)	402 (95%)	20 (5%)	23	24

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	LEU
1	A	155	LYS
1	A	158	LEU
1	A	195	LEU
1	A	252	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	255	ASP
1	A	268	GLU
1	A	338	LYS
1	B	138	LEU
1	B	169	GLU
1	B	174	GLU
1	B	177	LEU
1	B	184	GLN
1	B	187	LEU
1	B	195	LEU
1	B	231	ARG
1	B	268	GLU
1	B	304	MET
1	B	307	GLU
1	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	GLN
1	A	175	HIS
1	A	176	GLN
1	A	184	GLN
1	A	189	HIS
1	A	191	ASN
1	A	222	GLN
1	A	365	HIS
1	A	366	ASN
1	A	369	GLN
1	B	126	GLN
1	B	176	GLN
1	B	191	ASN
1	B	200	HIS
1	B	260	ASN
1	B	331	ASN
1	B	366	ASN
1	B	369	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HPM	A	1389	-	43,43,43	2.09	10 (23%)	57,58,58	1.73	11 (19%)
2	HPM	B	1388	-	43,43,43	2.17	13 (30%)	57,58,58	1.83	14 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HPM	A	1389	-	-	3/23/31/31	0/5/5/5
2	HPM	B	1388	-	-	4/23/31/31	0/5/5/5

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1388	HPM	C20-N21	7.62	1.47	1.36
2	A	1389	HPM	C20-N21	7.12	1.47	1.36
2	B	1388	HPM	C14-C15	4.32	1.49	1.42
2	A	1389	HPM	C14-C15	3.82	1.48	1.42
2	A	1389	HPM	C25-N28	3.80	1.43	1.38
2	B	1388	HPM	C25-N28	3.59	1.43	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1389	HPM	C16-C11	3.49	1.42	1.36
2	A	1389	HPM	C31-C30	3.26	1.44	1.39
2	B	1388	HPM	C31-C30	3.14	1.44	1.39
2	A	1389	HPM	C13-C12	3.09	1.42	1.36
2	B	1388	HPM	C13-C12	3.07	1.42	1.36
2	B	1388	HPM	C7-N6	3.04	1.53	1.47
2	A	1389	HPM	C27-C22	3.02	1.44	1.39
2	B	1388	HPM	C16-C11	2.94	1.42	1.36
2	B	1388	HPM	C35-C30	2.91	1.43	1.39
2	A	1389	HPM	C35-C30	2.84	1.43	1.39
2	B	1388	HPM	C5-N6	2.70	1.54	1.46
2	B	1388	HPM	C23-C22	2.55	1.43	1.39
2	A	1389	HPM	C7-N6	2.48	1.52	1.47
2	B	1388	HPM	C1-N6	2.36	1.53	1.46
2	B	1388	HPM	O37-C12	2.36	1.41	1.37
2	B	1388	HPM	C7-C8	2.10	1.57	1.52
2	A	1389	HPM	C20-N19	2.04	1.36	1.34

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1389	HPM	O37-C12-C13	-5.25	118.51	125.16
2	B	1388	HPM	O37-C12-C13	-4.68	119.23	125.16
2	B	1388	HPM	C9-O10-C11	4.41	128.00	118.21
2	B	1388	HPM	C8-C7-N6	4.37	120.16	112.23
2	A	1389	HPM	O37-C12-C11	4.28	121.21	115.40
2	A	1389	HPM	C22-C23-N24	-4.24	120.56	124.07
2	A	1389	HPM	C18-N19-C20	4.10	119.78	116.60
2	B	1388	HPM	O37-C12-C11	3.78	120.53	115.40
2	A	1389	HPM	O10-C11-C16	-3.77	119.02	125.20
2	B	1388	HPM	C22-C23-N24	-3.73	120.98	124.07
2	B	1388	HPM	C18-N19-C20	3.61	119.40	116.60
2	B	1388	HPM	C2-C1-N6	3.39	116.56	111.30
2	B	1388	HPM	O10-C11-C16	-3.02	120.23	125.20
2	A	1389	HPM	C22-C27-N26	-2.72	121.82	124.07
2	A	1389	HPM	O10-C11-C12	2.61	121.15	115.75
2	A	1389	HPM	C9-O10-C11	2.53	123.84	118.21
2	A	1389	HPM	C18-N17-C15	2.47	118.23	115.43
2	B	1388	HPM	C18-N17-C15	2.39	118.15	115.43
2	B	1388	HPM	C5-N6-C1	2.37	113.94	108.84
2	B	1388	HPM	C22-C27-N26	-2.33	122.15	124.07
2	B	1388	HPM	C4-C5-N6	2.29	114.86	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1388	HPM	C7-C8-C9	2.25	116.60	111.59
2	A	1389	HPM	C14-C20-N21	2.11	121.90	119.77
2	A	1389	HPM	N17-C18-N19	-2.07	125.86	128.67
2	B	1388	HPM	C38-O37-C12	2.02	120.47	117.51

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1388	HPM	C7-C8-C9-O10
2	A	1389	HPM	C11-C12-O37-C38
2	B	1388	HPM	O39-C8-C9-O10
2	A	1389	HPM	C13-C12-O37-C38
2	B	1388	HPM	C11-C12-O37-C38
2	B	1388	HPM	C13-C12-O37-C38
2	A	1389	HPM	N6-C7-C8-C9

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1389	HPM	2	0
2	B	1388	HPM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/283 (88%)	0.50	20 (7%) 18 19	13, 28, 51, 69	0
1	B	246/283 (86%)	0.64	30 (12%) 8 8	15, 28, 53, 72	0
All	All	497/566 (87%)	0.57	50 (10%) 12 13	13, 28, 53, 72	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	255	ASP	7.5
1	B	255	ASP	5.8
1	B	173	VAL	5.7
1	B	387	SER	5.2
1	A	173	VAL	4.7
1	A	254	ARG	4.7
1	A	291	THR	4.4
1	B	333	TYR	4.3
1	A	290	GLY	4.3
1	B	252	ILE	4.2
1	B	155	LYS	4.0
1	B	253	HIS	4.0
1	A	175	HIS	3.8
1	A	253	HIS	3.8
1	B	256	ILE	3.7
1	B	290	GLY	3.7
1	B	386	SER	3.4
1	A	256	ILE	3.3
1	B	254	ARG	3.2
1	B	172	GLY	3.2
1	A	252	ILE	2.8
1	B	291	THR	2.8
1	B	175	HIS	2.7
1	A	174	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	171	ALA	2.6
1	B	332	THR	2.5
1	B	154	SER	2.4
1	A	337	TYR	2.4
1	A	292	LEU	2.4
1	A	333	TYR	2.3
1	A	302	GLY	2.3
1	B	292	LEU	2.3
1	B	303	ARG	2.3
1	B	304	MET	2.3
1	B	140	LYS	2.3
1	A	182	GLU	2.2
1	A	304	MET	2.2
1	A	169	GLU	2.2
1	B	182	GLU	2.2
1	B	171	ALA	2.2
1	B	273	ASP	2.2
1	A	334	GLN	2.1
1	B	126	GLN	2.1
1	B	186	HIS	2.1
1	B	337	TYR	2.1
1	B	156	PHE	2.1
1	A	155	LYS	2.1
1	B	317	LEU	2.0
1	B	230	GLN	2.0
1	B	330	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HPM	B	1388	39/39	0.91	0.09	16,19,37,37	0
2	HPM	A	1389	39/39	0.94	0.07	15,19,28,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HPM B 1388:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.