



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 01:40 PM UTC

PDB ID : 2C76 / pdb_00002c76
Title : Functional Role of the Aromatic Cage in Human Monoamine Oxidase B: Structures and Catalytic Properties of Tyr435 Mutant Proteins
Authors : Li, M.; Binda, C.; Mattevi, A.; Edmondson, D.E.
Deposited on : 2005-11-17
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	6% 91% 5%
1	B	520	7% 88% 6% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	RSA	A	601	X	-	-	-
3	RSA	B	601	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called AMINE OXIDASE (FLAVIN-CONTAINING) B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	1	0
			3974	2540	682	727	25			
1	B	494	Total	C	N	O	S	0	1	0
			3943	2521	677	720	25			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	435	TRP	TYR	engineered mutation	UNP P27338
B	435	TRP	TYR	engineered mutation	UNP P27338

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is N-PROPARGYL-1(S)-AMINOINDAN (CCD ID: RSA) (formula: C₁₂H₁₃N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			13	12	1		
3	B	1	Total	C	N	0	0
			13	12	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	425	Total	O	0	0
			425	425		
4	B	485	Total	O	0	0
			485	485		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AMINE OXIDASE (FLAVIN-CONTAINING) B



• Molecule 1: AMINE OXIDASE (FLAVIN-CONTAINING) B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.24Å 223.06Å 86.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.70 15.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.5 (15.00-1.70) 98.3 (15.00-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.191 , 0.213 0.191 , 0.212	Depositor DCC
R_{free} test set	3489 reflections (2.55%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.014 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8959	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, RSA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/4077	0.73	0/5535
1	B	0.50	0/4046	0.73	0/5492
All	All	0.50	0/8123	0.73	0/11027

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3974	0	3969	25	0
1	B	3943	0	3939	26	0
2	A	53	0	29	2	0
2	B	53	0	29	2	0
3	A	13	0	13	1	0
3	B	13	0	13	0	0
4	A	425	0	0	1	1
4	B	485	0	0	4	0
All	All	8959	0	7992	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:LEU:HD11	1:B:454:MET:HE1	1.52	0.90
1:A:28:LEU:HD11	1:A:454:MET:HE1	1.53	0.89
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.30	0.80
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.32	0.76
1:B:414:LEU:HD12	4:B:2415:HOH:O	1.94	0.66
1:B:251:ASN:HD22	1:B:251:ASN:H	1.43	0.65
1:A:251:ASN:HD22	1:A:251:ASN:H	1.44	0.63
1:B:321:GLU:H	1:B:321:GLU:CD	2.11	0.59
1:A:451:LEU:HA	1:A:454:MET:HE2	1.86	0.57
1:B:280:MET:HE3	1:B:281:MET:SD	2.46	0.56
1:A:248:GLU:OE2	1:B:252:HIS:HE1	1.88	0.55
1:A:143:TRP:HA	1:A:146:MET:HE3	1.87	0.55
1:A:280:MET:HE3	1:A:281:MET:SD	2.47	0.55
1:A:321:GLU:H	1:A:321:GLU:CD	2.14	0.55
1:A:252:HIS:HE1	1:B:248:GLU:OE2	1.91	0.53
1:B:412:ARG:HG2	4:B:2413:HOH:O	2.10	0.52
1:B:451:LEU:HA	1:B:454:MET:HE2	1.91	0.51
1:B:233:ARG:HG3	1:B:251:ASN:HD21	1.76	0.50
1:A:233:ARG:HG3	1:A:251:ASN:HD21	1.76	0.50
1:A:270:MET:HE1	1:B:268:LEU:HD23	1.94	0.50
1:A:168:PHE:HE1	1:A:199:ILE:CG2	2.25	0.50
1:B:143:TRP:HA	1:B:146:MET:HE3	1.95	0.48
1:B:122:MET:HE1	1:B:185:PHE:HE2	1.80	0.46
1:A:280:MET:HE1	1:B:353:ALA:HB3	1.97	0.46
1:A:451:LEU:HD23	1:A:454:MET:CE	2.45	0.46
1:B:168:PHE:HE1	1:B:199:ILE:CG2	2.28	0.45
1:A:268:LEU:HD23	1:B:270:MET:HE1	1.98	0.45
1:A:353:ALA:HB3	1:B:280:MET:HE1	1.99	0.45
1:A:434:GLY:O	2:A:600:FAD:H1'2	2.17	0.45
1:A:168:PHE:CE1	1:A:199:ILE:CG2	3.00	0.45
1:B:387:ASN:O	1:B:390:GLU:HG2	2.17	0.45
1:A:136:LYS:HE3	4:A:2153:HOH:O	2.17	0.44
1:A:171:LEU:HD21	3:A:601:RSA:H21	2.00	0.44
1:A:168:PHE:HE1	1:A:199:ILE:HG22	1.82	0.44
1:A:58:GLY:HA2	2:A:600:FAD:C4X	2.49	0.43
1:B:168:PHE:CE1	1:B:199:ILE:CG2	3.02	0.42
1:B:451:LEU:HD23	1:B:454:MET:CE	2.49	0.42
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.09	0.42
1:B:58:GLY:HA2	2:B:600:FAD:C4X	2.50	0.42
1:A:451:LEU:HD23	1:A:454:MET:HE1	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:GLN:HB2	4:B:2229:HOH:O	2.20	0.41
1:B:434:GLY:O	2:B:600:FAD:H1'2	2.20	0.41
1:A:49:GLN:OE1	1:A:49:GLN:O	2.39	0.41
1:B:309:LYS:NZ	1:B:373:GLY:O	2.54	0.41
1:B:343:PHE:HB2	4:B:2356:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2049:HOH:O	4:A:2049:HOH:O[3_655]	1.57	0.63

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/520 (96%)	486 (98%)	12 (2%)	0	100	100
1	B	493/520 (95%)	481 (98%)	12 (2%)	0	100	100
All	All	991/1040 (95%)	967 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	428/444 (96%)	423 (99%)	5 (1%)	63	51
1	B	425/444 (96%)	421 (99%)	4 (1%)	70	62
All	All	853/888 (96%)	844 (99%)	9 (1%)	65	54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	106	VAL
1	A	155	LEU
1	A	251	ASN
1	A	397	CYS
1	B	49	GLN
1	B	155	LEU
1	B	251	ASN
1	B	397	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	49	GLN
1	A	117	ASN
1	A	251	ASN
1	A	252	HIS
1	B	3	ASN
1	B	116	ASN
1	B	117	ASN
1	B	251	ASN
1	B	252	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	600	1,3	58,58,58	1.13	5 (8%)	85,89,89	1.59	13 (15%)
3	RSA	A	601	2	13,14,14	1.61	2 (15%)	16,18,18	2.32	4 (25%)
2	FAD	A	600	1,3	58,58,58	1.17	7 (12%)	85,89,89	1.61	16 (18%)
3	RSA	B	601	2	13,14,14	1.55	2 (15%)	16,18,18	2.40	5 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	600	1,3	-	1/34/50/50	0/6/6/6
3	RSA	A	601	2	1/1/1/2	2/3/13/13	0/2/2/2
2	FAD	A	600	1,3	-	2/34/50/50	0/6/6/6
3	RSA	B	601	2	1/1/1/2	2/3/13/13	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	RSA	C12-C13	4.76	1.32	1.18
3	B	601	RSA	C12-C13	4.55	1.31	1.18
2	A	600	FAD	C4X-N5	4.22	1.39	1.30
2	B	600	FAD	C4X-N5	4.11	1.39	1.30
2	A	600	FAD	C10-N1	3.04	1.39	1.33
2	B	600	FAD	C10-N1	2.92	1.39	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	FAD	C2A-N1A	2.71	1.38	1.33
2	B	600	FAD	C2A-N1A	2.65	1.38	1.33
2	A	600	FAD	C8A-N7A	2.41	1.36	1.31
2	A	600	FAD	C2A-N3A	2.41	1.38	1.33
3	B	601	RSA	C11-C12	2.40	1.54	1.46
3	A	601	RSA	C11-C12	2.39	1.54	1.46
2	A	600	FAD	P-O3P	2.25	1.61	1.59
2	A	600	FAD	PA-O3P	2.24	1.61	1.59
2	B	600	FAD	C2A-N3A	2.07	1.37	1.33
2	B	600	FAD	C8A-N7A	2.03	1.35	1.31

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	RSA	C11-C12-C13	-7.10	125.46	176.85
3	B	601	RSA	C11-C12-C13	-7.06	125.74	176.85
2	A	600	FAD	N3A-C2A-N1A	-5.73	119.90	128.58
2	B	600	FAD	N3A-C2A-N1A	-5.58	120.14	128.58
2	B	600	FAD	N9A-C8A-N7A	-4.03	108.22	113.94
2	A	600	FAD	N9A-C8A-N7A	-4.01	108.25	113.94
2	B	600	FAD	C9A-C5X-N5	-3.92	118.30	122.45
2	A	600	FAD	C9A-C5X-N5	-3.86	118.36	122.45
2	B	600	FAD	C5A-N7A-C8A	3.73	109.32	103.45
2	B	600	FAD	C5A-C4A-N3A	-3.60	121.76	126.72
2	A	600	FAD	C9A-N10-C10	-3.49	115.44	120.75
2	A	600	FAD	C5A-C4A-N3A	-3.21	122.30	126.72
2	A	600	FAD	C5A-N7A-C8A	3.19	108.47	103.45
3	A	601	RSA	C11-N10-C9	3.17	120.09	113.80
2	A	600	FAD	C4-N3-C2	-3.15	120.05	125.64
2	B	600	FAD	C9A-N10-C10	-3.11	116.01	120.75
2	B	600	FAD	C2A-N3A-C4A	2.99	119.13	111.83
2	B	600	FAD	C4A-C5A-N7A	-2.97	107.19	110.58
3	A	601	RSA	C12-C11-N10	2.97	121.88	112.51
2	B	600	FAD	C4-N3-C2	-2.95	120.41	125.64
2	A	600	FAD	C4X-C4-N3	2.86	120.52	113.25
2	A	600	FAD	C2A-N3A-C4A	2.85	118.79	111.83
3	B	601	RSA	C12-C11-N10	2.84	121.49	112.51
2	B	600	FAD	C4X-C4-N3	2.68	120.08	113.25
3	B	601	RSA	C6-C7-C8	-2.62	117.86	120.99
2	A	600	FAD	C10-C4X-N5	-2.59	119.52	124.81
2	B	600	FAD	O4-C4-C4X	-2.56	119.78	126.53
3	B	601	RSA	C11-N10-C9	2.49	118.75	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	FAD	C4A-C5A-N7A	-2.46	107.77	110.58
2	A	600	FAD	C4-C4X-N5	2.45	121.59	118.21
2	B	600	FAD	C5'-C4'-C3'	-2.44	107.62	112.22
2	B	600	FAD	C10-C4X-N5	-2.35	120.01	124.81
2	A	600	FAD	C5'-C4'-C3'	-2.30	107.89	112.22
2	A	600	FAD	O4-C4-C4X	-2.26	120.57	126.53
2	A	600	FAD	C4A-N9A-C1B	-2.20	121.48	126.63
2	A	600	FAD	C4A-N9A-C8A	2.17	108.01	105.74
3	A	601	RSA	C6-C7-C8	-2.11	118.46	120.99
3	B	601	RSA	C5-C4-C3	-2.10	117.81	120.88

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	601	RSA	C9
3	B	601	RSA	C9

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	RSA	C1-C9-N10-C11
3	A	601	RSA	C8-C9-N10-C11
3	B	601	RSA	C1-C9-N10-C11
3	B	601	RSA	C8-C9-N10-C11
2	A	600	FAD	C2'-C1'-N10-C10
2	B	600	FAD	C2'-C1'-N10-C10
2	A	600	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	600	FAD	2	0
3	A	601	RSA	1	0
2	A	600	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	499/520 (95%)	0.03	32 (6%)	25 27	8, 14, 31, 55	1 (0%)
1	B	494/520 (95%)	0.04	35 (7%)	22 24	8, 14, 28, 48	1 (0%)
All	All	993/1040 (95%)	0.03	67 (6%)	24 26	8, 14, 30, 55	2 (0%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	495	LEU	5.8
1	B	496	ILE	5.6
1	A	501	ILE	5.4
1	B	493	LEU	4.1
1	A	480	THR	4.0
1	A	92	VAL	4.0
1	B	494	ARG	3.9
1	B	107	TRP	3.8
1	B	103	PHE	3.8
1	B	481	PHE	3.7
1	B	95	LYS	3.5
1	A	498	LEU	3.5
1	B	491	GLY	3.5
1	A	107	TRP	3.3
1	A	495	LEU	3.2
1	B	3	ASN	3.2
1	A	103	PHE	3.2
1	B	492	LEU	3.1
1	A	481	PHE	3.1
1	B	412	ARG	3.1
1	A	49	GLN	2.9
1	A	492	LEU	2.9
1	A	493	LEU	2.9
1	A	321	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	93	LYS	2.8
1	A	496	ILE	2.8
1	A	482	LEU	2.7
1	B	112	TYR	2.7
1	B	482	LEU	2.7
1	B	163	GLN	2.7
1	A	500	THR	2.7
1	B	110	ILE	2.6
1	A	479	THR	2.6
1	B	157	TRP	2.6
1	A	484	ARG	2.6
1	A	489	VAL	2.6
1	A	499	THR	2.5
1	B	252	HIS	2.5
1	A	3	ASN	2.5
1	B	244	ASN	2.5
1	B	119	TRP	2.5
1	B	321	GLU	2.5
1	B	243	GLU	2.4
1	A	413	VAL	2.4
1	A	252	HIS	2.4
1	A	101	GLY	2.4
1	B	370	LYS	2.3
1	A	111	THR	2.3
1	A	486	LEU	2.3
1	B	246	LEU	2.3
1	A	412	ARG	2.3
1	A	494	ARG	2.3
1	A	490	PRO	2.2
1	B	476	PRO	2.2
1	B	490	PRO	2.2
1	B	242	ARG	2.2
1	A	227	ASP	2.2
1	A	475	GLN	2.2
1	B	489	VAL	2.1
1	B	226	GLY	2.1
1	B	254	MET	2.1
1	B	92	VAL	2.1
1	B	475	GLN	2.1
1	A	102	PRO	2.1
1	A	483	GLU	2.0
1	B	486	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	479	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	RSA	A	601	13/13	0.94	0.07	10,13,14,14	0
3	RSA	B	601	13/13	0.95	0.06	10,13,13,14	0
2	FAD	A	600	53/53	0.97	0.04	8,10,11,11	0
2	FAD	B	600	53/53	0.97	0.04	7,10,11,11	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.