



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:11 AM UTC

PDB ID : 4C74 / pdb_00004c74
Title : Phenylacetone monooxygenase: Reduced enzyme in complex with APADP
Authors : Martinoli, C.; Fraaije, M.W.; Mattevi, A.
Deposited on : 2013-09-19
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	 4% 79% 17% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4569 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLACETONE MONOOXYGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	531	4240	2706	729	795	10	0	3	0


- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	22	6	17	3		

- Molecule 4 is HEXAETHYLENE GLYCOL (CCD ID: P6G) (formula: C₁₂H₂₆O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	218	Total 218	O 218	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHENYLACETONE MONOOXYGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	107.62Å 107.62Å 106.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	70.22 – 1.97 70.22 – 1.97	Depositor EDS
% Data completeness (in resolution range)	98.7 (70.22-1.97) 98.7 (70.22-1.97)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.235 , 0.278 (Not available) , 0.296	Depositor DCC
R_{free} test set	2571 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.035 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4569	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: N01, P6G, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.28	11/4362 (0.3%)	1.30	19/5935 (0.3%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	525	ARG	CZ-NH1	6.37	1.41	1.32
1	A	516	LEU	N-CA	6.36	1.53	1.45
1	A	489	ILE	N-CA	6.19	1.54	1.46
1	A	507	VAL	N-CA	6.17	1.50	1.45
1	A	202	SER	N-CA	6.10	1.52	1.46
1	A	248	GLU	CA-C	5.80	1.60	1.52
1	A	506	ASN	CA-C	5.70	1.60	1.52
1	A	453	HIS	CA-C	-5.29	1.46	1.52
1	A	95	GLU	C-O	5.10	1.30	1.24
1	A	153	LEU	CA-C	-5.03	1.47	1.53
1	A	268	VAL	C-O	-5.02	1.17	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	LEU	CA-C-N	-14.51	106.16	120.52
1	A	158	LEU	C-N-CA	-14.51	106.16	120.52
1	A	336	LYS	N-CA-C	-7.72	97.74	110.17
1	A	177	TRP	CA-C-N	-6.45	113.64	120.03
1	A	177	TRP	C-N-CA	-6.45	113.64	120.03
1	A	438	GLY	N-CA-C	-6.25	106.46	115.27
1	A	231	ASP	CA-C-N	-6.16	113.46	119.87
1	A	231	ASP	C-N-CA	-6.16	113.46	119.87
1	A	12	GLN	CA-C-N	-6.07	114.13	120.38
1	A	12	GLN	C-N-CA	-6.07	114.13	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	311	ASN	N-CA-C	5.74	117.53	111.28
1	A	462	ALA	N-CA-C	-5.54	105.14	111.07
1	A	538	GLY	N-CA-C	-5.54	107.68	115.32
1	A	237	ASP	CB-CA-C	-5.36	101.74	110.85
1	A	506	ASN	N-CA-CB	-5.28	101.34	110.32
1	A	351	ASP	N-CA-C	5.25	117.68	111.33
1	A	499	ALA	N-CA-C	-5.20	100.62	108.67
1	A	159	PRO	CA-C-N	-5.06	115.82	122.85
1	A	159	PRO	C-N-CA	-5.06	115.82	122.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4240	0	4096	65	0
2	A	53	0	31	2	0
3	A	48	0	26	1	0
4	A	10	0	13	6	1
5	A	218	0	0	17	0
All	All	4569	0	4166	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:311:ASN:HB3	5:A:2135:HOH:O	1.31	1.27
4:A:1543:P6G:H92	5:A:2218:HOH:O	1.36	1.26
1:A:530:GLU:HB3	5:A:2216:HOH:O	1.51	1.10
1:A:12:GLN:HB3	5:A:2012:HOH:O	1.59	1.00
1:A:506:ASN:OD1	1:A:506:ASN:N	2.04	0.90
1:A:402:ARG:HD2	1:A:406:ASN:HD21	1.41	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:ARG:HD2	1:A:357:ASP:HB2	1.60	0.82
1:A:523:ARG:HD3	5:A:2212:HOH:O	1.79	0.82
1:A:402:ARG:HD2	1:A:406:ASN:ND2	1.99	0.77
1:A:279:ARG:HB3	4:A:1543:P6G:H51	1.66	0.76
1:A:200:GLN:NE2	3:A:701:N01:HAAB	2.02	0.75
1:A:313:ILE:O	1:A:316:THR:HG22	1.87	0.74
1:A:281:TRP:CD2	1:A:312[B]:LYS:HD3	2.23	0.73
1:A:335:THR:OG1	1:A:498:THR:HG21	1.89	0.73
1:A:369:ARG:HD3	1:A:379:GLU:OE2	1.92	0.70
1:A:65[A]:CYS:SG	2:A:700:FAD:O4	2.49	0.70
1:A:325:ARG:HD3	5:A:2138:HOH:O	1.90	0.70
1:A:311:ASN:CB	5:A:2135:HOH:O	2.05	0.69
1:A:281:TRP:CE2	1:A:312[B]:LYS:HD3	2.31	0.66
1:A:152:GLN:HE22	2:A:700:FAD:H5'2	1.60	0.66
1:A:512:ARG:HH11	1:A:512:ARG:HB3	1.60	0.66
1:A:332:PRO:HD2	1:A:498:THR:CG2	2.27	0.65
1:A:199:ILE:HD12	1:A:339:ILE:HD13	1.80	0.63
1:A:316:THR:HG23	1:A:317:VAL:HG23	1.80	0.63
1:A:260:GLN:HG2	5:A:2113:HOH:O	1.97	0.63
1:A:311:ASN:CG	5:A:2135:HOH:O	2.37	0.62
1:A:294:ASP:HA	1:A:297:ARG:HG3	1.82	0.61
1:A:144:ARG:HG2	1:A:145:TYR:CE2	2.36	0.61
1:A:12:GLN:N	5:A:2001:HOH:O	2.33	0.60
1:A:299:ARG:HH12	1:A:497:MET:HG3	1.68	0.57
1:A:332:PRO:HD2	1:A:498:THR:HG21	1.86	0.57
1:A:316:THR:CG2	1:A:345:TYR:HB3	2.35	0.57
1:A:206:ALA:O	1:A:350:ARG:NH2	2.37	0.57
1:A:501:TRP:HA	1:A:506:ASN:OD1	2.05	0.56
1:A:12:GLN:CB	5:A:2012:HOH:O	2.33	0.56
1:A:202:SER:HB2	1:A:203:PRO:HD3	1.86	0.55
1:A:316:THR:HG21	1:A:345:TYR:HB3	1.89	0.55
1:A:128:THR:HG21	1:A:142:ARG:HH22	1.73	0.54
1:A:332:PRO:HD2	1:A:498:THR:HG22	1.91	0.52
4:A:1543:P6G:C5	5:A:2119:HOH:O	2.56	0.52
1:A:23:GLY:O	1:A:28:GLY:HA3	2.10	0.51
1:A:523:ARG:HA	1:A:526:GLN:HE21	1.76	0.51
1:A:128:THR:O	1:A:130:THR:HG23	2.12	0.50
1:A:495:TYR:HB3	1:A:514:PHE:CE1	2.47	0.50
1:A:316:THR:CG2	1:A:317:VAL:HG23	2.43	0.49
1:A:504:GLY:C	1:A:506:ASN:OD1	2.56	0.48
1:A:241:ARG:HD2	5:A:2106:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:ALA:O	1:A:304:ARG:NH1	2.45	0.48
1:A:512:ARG:HB3	1:A:512:ARG:NH1	2.25	0.48
1:A:446:MET:HE2	1:A:446:MET:HB3	1.59	0.47
1:A:164:LEU:HD13	1:A:366:ILE:HG13	1.97	0.47
1:A:512:ARG:HD3	5:A:2205:HOH:O	2.13	0.47
1:A:12:GLN:HG3	1:A:114:THR:HG23	1.97	0.45
1:A:402:ARG:CD	1:A:406:ASN:ND2	2.75	0.45
1:A:429:PRO:O	1:A:430:ASN:HB2	2.16	0.45
4:A:1543:P6G:H52	5:A:2120:HOH:O	2.17	0.45
1:A:279:ARG:CZ	4:A:1543:P6G:H21	2.47	0.45
1:A:48:ALA:HB1	1:A:56:TYR:CE1	2.54	0.43
1:A:478:LYS:HE2	1:A:478:LYS:HB3	1.67	0.42
1:A:17:VAL:O	1:A:143:ALA:HA	2.19	0.42
1:A:18:ASP:HB2	1:A:41:SER:O	2.19	0.42
1:A:316:THR:HG23	1:A:345:TYR:HB3	2.02	0.42
1:A:344:TYR:O	1:A:347:MET:HB3	2.20	0.41
1:A:350:ARG:H	1:A:350:ARG:HG3	1.72	0.41
1:A:20:LEU:HD13	1:A:141:ILE:HG21	2.02	0.41
4:A:1543:P6G:C9	5:A:2218:HOH:O	2.20	0.41
1:A:298:ASP:OD1	1:A:298:ASP:C	2.63	0.41
1:A:525:ARG:NH1	5:A:2214:HOH:O	2.54	0.40
1:A:188:ARG:NH1	1:A:211:GLU:HG3	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1543:P6G:O10	4:A:1543:P6G:O10[6_555]	1.92	0.28

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	532/542 (98%)	516 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	444/453 (98%)	421 (95%)	23 (5%)	21	9

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	42	VAL
1	A	150	SER
1	A	160	ASN
1	A	164	LEU
1	A	196	SER
1	A	230	LEU
1	A	235	LEU
1	A	273	LEU
1	A	297	ARG
1	A	298	ASP
1	A	316	THR
1	A	329	LYS
1	A	360	SER
1	A	377	GLU
1	A	444	SER
1	A	447	LEU
1	A	478	LYS
1	A	484	GLU
1	A	498	THR
1	A	506	ASN
1	A	512	ARG
1	A	540	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	129	ASN
1	A	152	GLN
1	A	160	ASN
1	A	170	ASN
1	A	187	GLN
1	A	315	ASN
1	A	406	ASN
1	A	526	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	P6G	A	1543	-	9,9,18	0.78	0	8,8,17	0.96	0
2	FAD	A	700	-	58,58,58	1.24	7 (12%)	85,89,89	1.57	18 (21%)
3	N01	A	701	-	50,52,52	1.74	8 (16%)	71,80,80	2.12	23 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	P6G	A	1543	-	-	5/7/7/16	-
2	FAD	A	700	-	-	2/34/50/50	0/6/6/6
3	N01	A	701	-	-	6/35/67/67	0/5/5/5

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	N01	O4D-C1D	5.46	1.48	1.40
3	A	701	N01	P2B-O2B	4.38	1.67	1.59
3	A	701	N01	CAA-C7N	-3.65	1.39	1.49
3	A	701	N01	C4N-C3N	3.58	1.44	1.39
3	A	701	N01	C5N-C4N	3.47	1.44	1.38
2	A	700	FAD	C5A-N7A	-3.17	1.33	1.39
3	A	701	N01	C2N-C3N	3.10	1.43	1.39
2	A	700	FAD	C4X-N5	2.90	1.37	1.30
2	A	700	FAD	PA-O3P	2.80	1.62	1.59
3	A	701	N01	C6A-N6A	2.55	1.40	1.34
2	A	700	FAD	C8A-N7A	2.41	1.36	1.31
3	A	701	N01	PN-O3	2.40	1.62	1.59
2	A	700	FAD	C2A-N1A	2.37	1.38	1.33
2	A	700	FAD	O2-C2	-2.28	1.19	1.24
2	A	700	FAD	C8M-C8	2.06	1.54	1.51

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	N01	C2B-C1B-N9A	-5.85	104.12	113.75
3	A	701	N01	O3-PA-O2A	-5.18	95.11	110.70
3	A	701	N01	N3A-C2A-N1A	-4.76	121.37	128.58
2	A	700	FAD	C5A-C4A-N3A	-4.41	120.64	126.72
3	A	701	N01	CAA-C7N-C3N	4.29	127.33	119.21
2	A	700	FAD	N3A-C2A-N1A	-4.27	122.11	128.58
3	A	701	N01	C4A-N9A-C8A	4.20	110.15	105.74
3	A	701	N01	N9A-C8A-N7A	-4.10	108.12	113.94
3	A	701	N01	C3B-C2B-C1B	-3.60	95.91	102.81
2	A	700	FAD	C4X-C10-N10	3.56	121.58	116.48
3	A	701	N01	O3-PN-O2N	-3.47	100.26	110.70
2	A	700	FAD	C5A-N7A-C8A	3.28	108.60	103.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	700	FAD	C9A-N10-C10	-3.11	116.01	120.75
3	A	701	N01	O2B-P2B-O3X	-3.10	98.28	109.33
3	A	701	N01	O2X-P2B-O1X	3.01	119.10	107.80
2	A	700	FAD	C9-C9A-N10	-2.96	117.88	121.85
3	A	701	N01	O5B-C5B-C4B	-2.95	98.95	108.99
3	A	701	N01	C2A-N1A-C6A	2.92	123.53	118.73
3	A	701	N01	C4A-N9A-C1B	-2.91	119.83	126.63
3	A	701	N01	N6A-C6A-N1A	2.89	124.81	118.38
3	A	701	N01	O1N-PN-O3	2.83	114.93	107.27
2	A	700	FAD	C5A-C4A-N9A	2.83	108.89	105.81
3	A	701	N01	O3B-C3B-C4B	-2.77	103.12	111.08
3	A	701	N01	P2B-O2B-C2B	-2.74	116.11	123.43
2	A	700	FAD	C4-N3-C2	-2.74	120.77	125.64
2	A	700	FAD	C2A-N3A-C4A	2.74	118.52	111.83
3	A	701	N01	C5A-N7A-C8A	2.72	107.73	103.45
3	A	701	N01	C2A-N3A-C4A	2.70	118.42	111.83
2	A	700	FAD	C4A-C5A-N7A	-2.69	107.51	110.58
2	A	700	FAD	N9A-C8A-N7A	-2.65	110.17	113.94
2	A	700	FAD	C5X-C9A-N10	2.59	120.31	117.97
2	A	700	FAD	C6A-C5A-C4A	2.57	120.68	117.18
3	A	701	N01	N3A-C4A-N9A	2.53	131.47	127.17
3	A	701	N01	C5A-C4A-N3A	-2.44	123.35	126.72
2	A	700	FAD	C4'-C3'-C2'	2.26	117.32	113.57
2	A	700	FAD	O4B-C1B-N9A	2.25	112.41	108.09
2	A	700	FAD	O2A-PA-O3P	2.20	113.22	107.27
3	A	701	N01	O7N-C7N-C3N	-2.20	115.60	119.92
3	A	701	N01	C5N-C4N-C3N	-2.14	118.26	120.36
2	A	700	FAD	C5X-N5-C4X	-2.12	114.67	118.09
2	A	700	FAD	C4-C4X-C10	2.02	120.39	116.93

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	700	FAD	N10-C1'-C2'-O2'
2	A	700	FAD	N10-C1'-C2'-C3'
3	A	701	N01	C5D-O5D-PN-O3
3	A	701	N01	C5D-O5D-PN-O2N
3	A	701	N01	C2N-C3N-C7N-O7N
3	A	701	N01	C2N-C3N-C7N-CAA
3	A	701	N01	C4N-C3N-C7N-O7N
4	A	1543	P6G	O7-C8-C9-O10

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	701	N01	C4N-C3N-C7N-CAA
4	A	1543	P6G	O4-C5-C6-O7
4	A	1543	P6G	C6-C5-O4-C3
4	A	1543	P6G	C2-C3-O4-C5
4	A	1543	P6G	C5-C6-O7-C8

There are no ring outliers.


3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1543	P6G	6	1
2	A	700	FAD	2	0
3	A	701	N01	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand FAD A 700



Ligand N01 A 701



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	531/542 (97%)	0.58	20 (3%) 44 54	17, 31, 51, 71	3 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	161	PHE	3.9
1	A	235	LEU	3.6
1	A	160	ASN	3.4
1	A	506	ASN	3.4
1	A	233	GLU	2.8
1	A	497	MET	2.7
1	A	165	LYS	2.6
1	A	127	ALA	2.5
1	A	142	ARG	2.4
1	A	491	ASP	2.4
1	A	350	ARG	2.4
1	A	232	PRO	2.4
1	A	283	GLU	2.2
1	A	179	HIS	2.2
1	A	534	LYS	2.2
1	A	512	ARG	2.1
1	A	375	GLU	2.1
1	A	231	ASP	2.1
1	A	128	THR	2.0
1	A	343	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	P6G	A	1543	10/19	0.87	0.17	40,44,46,53	0
2	FAD	A	700	53/53	0.97	0.06	18,22,29,34	0
3	N01	A	701	48/48	0.98	0.06	18,24,31,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around FAD A 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around N01 A 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.