



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 01:45 PM UTC

PDB ID : 8C7F / pdb_00008c7f
Title : Crystal structure of beta-xylosidase mutant (D281N, E517Q) from *Thermotoga maritima* in complex with xylopentaose
Authors : Gloster, T.M.; Foltanyi, F.
Deposited on : 2023-01-15
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	780	2% 94% ..
1	B	780	2% 90% .. 5%
2	X	3	100%
3	C	4	25% 75%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MPD	B	801	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 25248 atoms, of which 11908 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Xylosidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	772	Total	C	H	N	O	S	0	10	0
			12068	3864	6007	1037	1137	23			
1	B	739	Total	C	H	N	O	S	0	11	0
			11705	3740	5832	1003	1108	22			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	281	ASN	ASP	engineered mutation	UNP Q9WXT1
A	517	GLN	GLU	engineered mutation	UNP Q9WXT1
A	779	ALA	-	expression tag	UNP Q9WXT1
A	780	ALA	-	expression tag	UNP Q9WXT1
B	281	ASN	ASP	engineered mutation	UNP Q9WXT1
B	517	GLN	GLU	engineered mutation	UNP Q9WXT1
B	779	ALA	-	expression tag	UNP Q9WXT1
B	780	ALA	-	expression tag	UNP Q9WXT1

- Molecule 2 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	X	3	Total	C	H	O	0	0	0
			46	15	18	13			

- Molecule 3 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	4	Total	C	H	O	0	0	0
			62	20	25	17			

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			21	6	13	2		
4	B	1	Total	C	H	O	0	0
			21	6	13	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	695	Total	O	0	1
			696	696		
5	B	627	Total	O	0	2
			629	629		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Xylosidase



• Molecule 1: Xylosidase



• Molecule 2: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose



• Molecule 3: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	98.35Å 100.89Å 179.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.45 – 1.52 50.45 – 1.52	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.45-1.52) 99.8 (50.45-1.52)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.164 , 0.189 0.164 , 0.188	Depositor DCC
R_{free} test set	13593 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.431	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	25248	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	1/6173 (0.0%)	0.94	3/8343 (0.0%)
1	B	0.66	0/5981	0.94	3/8078 (0.0%)
All	All	0.66	1/12154 (0.0%)	0.94	6/16421 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	318	GLU	CD-OE1	5.75	1.36	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	223	ARG	CG-CD-NE	-6.79	97.07	112.00
1	A	223	ARG	CG-CD-NE	-6.24	98.27	112.00
1	A	371	ASP	CA-CB-CG	5.97	118.57	112.60
1	B	721	ASP	CA-CB-CG	5.89	118.49	112.60
1	B	371	ASP	CA-CB-CG	5.46	118.06	112.60
1	A	721	ASP	CA-CB-CG	5.30	117.90	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	346	ARG	Sidechain
1	A	600	ARG	Sidechain
1	A	674	ARG	Sidechain
1	B	769	ARG	Sidechain
1	B	85[A]	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6061	6007	6091	22	0
1	B	5873	5832	5924	26	0
2	X	28	18	0	0	0
3	C	37	25	0	0	0
4	A	8	13	14	5	0
4	B	8	13	14	6	0
5	A	696	0	0	9	0
5	B	629	0	0	7	0
All	All	13340	11908	12043	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:ASN:HD21	4:A:801:MPD:H12	1.13	1.09
1:B:260:ASN:HD21	4:B:801:MPD:HM2	1.11	1.09
1:B:321:GLN:H	1:B:321:GLN:HE21	1.12	0.92
1:B:260:ASN:ND2	4:B:801:MPD:HM2	1.90	0.85
1:A:260:ASN:ND2	4:A:801:MPD:H12	1.93	0.80
1:A:653:PRO:HG2	1:A:656:VAL:HG12	1.66	0.77
1:B:768:MET:HE3	5:B:1300:HOH:O	1.89	0.73
1:B:50:LYS:O	1:B:53:GLU:HG3	1.90	0.70
1:B:318:GLU:C	1:B:321:GLN:HE22	1.98	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:321:GLN:HE21	1:B:321:GLN:N	1.90	0.69
1:B:768:MET:CE	5:B:1300:HOH:O	2.43	0.65
4:B:801:MPD:H52	5:B:1230:HOH:O	2.00	0.62
4:A:801:MPD:H11	5:A:1253:HOH:O	2.01	0.61
1:B:226:LYS:NZ	5:B:901:HOH:O	2.28	0.60
4:B:801:MPD:HM1	5:B:1203:HOH:O	2.00	0.60
4:A:801:MPD:H52	5:A:1273:HOH:O	2.03	0.59
4:B:801:MPD:O4	4:B:801:MPD:H13	2.03	0.57
1:A:769[B]:ARG:NH2	5:A:902:HOH:O	2.38	0.57
1:A:12:GLU:HG3	5:A:1536:HOH:O	2.05	0.57
1:A:768:MET:CE	5:A:1344:HOH:O	2.54	0.55
1:A:656:VAL:HG11	1:A:662[B]:VAL:HG22	1.88	0.54
1:B:653:PRO:HG2	1:B:656:VAL:HG12	1.88	0.53
1:B:653:PRO:HG2	1:B:656:VAL:CG1	2.38	0.53
1:A:448:SER:HA	1:A:451:GLU:HG2	1.91	0.52
1:A:142:ARG:HG2	5:A:1390:HOH:O	2.09	0.52
1:A:656:VAL:HG11	1:A:662[B]:VAL:CG2	2.41	0.50
1:B:185:LYS:HE3	1:B:240:ALA:HA	1.93	0.50
1:A:768:MET:HE3	5:A:1344:HOH:O	2.10	0.50
1:B:506:LYS:HE2	5:B:1225:HOH:O	2.12	0.50
1:A:653:PRO:HG2	1:A:656:VAL:CG1	2.39	0.48
1:A:52:LYS:O	1:A:56:LYS:HB3	2.13	0.48
1:B:82:GLU:OE1	1:B:85[B]:ARG:NH2	2.46	0.48
1:A:72:GLU:HB2	1:A:75:GLU:OE1	2.15	0.46
1:B:223:ARG:NH1	1:B:223:ARG:HG3	2.30	0.46
1:A:327:VAL:HG21	1:A:337:ILE:HD12	1.98	0.45
1:A:397:LYS:HG2	5:A:1304:HOH:O	2.17	0.45
1:B:139[B]:GLU:HB3	5:B:909:HOH:O	2.16	0.45
1:A:239:GLU:HG3	1:A:771:PHE:CD2	2.52	0.44
1:A:374[B]:LEU:HD23	1:A:575:ALA:HB1	1.98	0.44
1:A:443:GLU:N	5:A:918:HOH:O	2.51	0.44
1:A:223:ARG:NH1	1:A:223:ARG:HG3	2.33	0.43
1:B:490:ILE:HD11	1:B:536:GLU:HB3	2.00	0.43
1:B:251:GLU:HA	1:B:256:PRO:HA	2.01	0.42
1:B:185:LYS:CE	1:B:185:LYS:HA	2.48	0.42
1:B:49:GLU:H	1:B:49:GLU:CD	2.27	0.42
1:B:401:ILE:HA	1:B:473:ALA:O	2.19	0.42
1:B:223:ARG:HG2	1:B:730:MET:SD	2.60	0.42
1:B:570:TRP:O	1:B:571:LEU:C	2.62	0.41
4:A:801:MPD:HM1	4:A:801:MPD:H4	1.82	0.41
1:A:87:LEU:O	1:A:94:GLY:HA2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:327:VAL:HG21	1:B:337:ILE:HD12	2.02	0.41
4:B:801:MPD:O4	4:B:801:MPD:C1	2.68	0.40
1:B:185:LYS:HE3	1:B:240:ALA:CA	2.51	0.40
1:A:570:TRP:O	1:A:571:LEU:C	2.63	0.40
1:B:100:HIS:CD2	1:B:150:LEU:HD11	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	778/780 (100%)	763 (98%)	15 (2%)	0	100	100
1	B	746/780 (96%)	733 (98%)	13 (2%)	0	100	100
All	All	1524/1560 (98%)	1496 (98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	643/660 (97%)	638 (99%)	5 (1%)	73	53
1	B	630/660 (96%)	622 (99%)	8 (1%)	61	34
All	All	1273/1320 (96%)	1260 (99%)	13 (1%)	68	44

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	ILE
1	A	408	VAL
1	A	421	ILE
1	A	448	SER
1	A	643	LYS
1	B	12	GLU
1	B	53	GLU
1	B	114	ASN
1	B	130	ILE
1	B	185	LYS
1	B	321	GLN
1	B	407	GLU
1	B	707	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	568	GLN
1	B	57	ASN
1	B	321	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	XYP	C	1	3	10,10,10	0.74	0	14,14,14	1.57	3 (21%)
3	XYP	C	2	3	9,9,10	0.87	0	10,12,14	2.43	3 (30%)
3	XYP	C	3	3	9,9,10	0.69	0	10,12,14	0.91	0
3	XYP	C	4	3	9,9,10	0.37	0	10,12,14	1.10	1 (10%)
2	XYP	X	1	2	10,10,10	0.36	0	14,14,14	1.00	1 (7%)
2	XYP	X	2	2	9,9,10	0.65	0	10,12,14	1.84	1 (10%)
2	XYP	X	3	2	9,9,10	0.67	0	10,12,14	1.47	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XYP	C	1	3	-	-	0/1/1/1
3	XYP	C	2	3	-	-	0/1/1/1
3	XYP	C	3	3	-	-	0/1/1/1
3	XYP	C	4	3	-	-	0/1/1/1
2	XYP	X	1	2	-	-	0/1/1/1
2	XYP	X	2	2	-	-	0/1/1/1
2	XYP	X	3	2	-	-	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	XYP	C1-C2-C3	5.80	118.08	109.64
2	X	2	XYP	C4-C3-C2	5.31	117.23	110.92
3	C	1	XYP	O4-C4-C5	-3.86	100.38	109.22
2	X	3	XYP	C1-C2-C3	3.31	114.46	109.64
3	C	1	XYP	C5-C4-C3	2.73	113.62	109.64
3	C	1	XYP	C4-C3-C2	2.31	114.93	110.86
3	C	2	XYP	C4-C3-C2	-2.28	108.21	110.92
2	X	1	XYP	O4-C4-C3	-2.20	105.60	110.15
3	C	2	XYP	O4-C4-C5	-2.15	104.29	109.22
3	C	4	XYP	O2-C2-C3	-2.13	105.74	110.15


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MPD	B	801	-	7,7,7	0.35	0	9,10,10	0.92	0
4	MPD	A	801	-	7,7,7	0.18	0	9,10,10	0.95	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	B	801	-	-	0/5/5/5	-
4	MPD	A	801	-	-	0/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	801	MPD	C5-C4-C3	2.05	121.19	111.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	801	MPD	6	0
4	A	801	MPD	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	772/780 (98%)	-0.39	18 (2%) 61 67	8, 20, 38, 63	10 (1%)
1	B	739/780 (94%)	-0.42	13 (1%) 67 73	8, 19, 37, 52	11 (1%)
All	All	1511/1560 (96%)	-0.40	31 (2%) 63 69	8, 20, 38, 63	21 (1%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	459	VAL	6.4
1	B	460	LEU	4.9
1	A	424	LEU	4.6
1	A	432	PHE	3.9
1	B	576	GLY	3.7
1	B	418	LEU	3.6
1	A	443	GLU	3.3
1	A	445	LEU	3.0
1	A	780	ALA	3.0
1	A	427	ASN	3.0
1	B	407	GLU	2.8
1	A	425	LEU	2.8
1	A	430	ASP	2.8
1	A	431	VAL	2.8
1	A	428	ILE	2.8
1	B	490	ILE	2.7
1	B	419	ALA	2.6
1	A	421	ILE	2.6
1	A	446	LYS	2.6
1	A	429	ASP	2.5
1	B	468	ILE	2.4
1	B	575	ALA	2.4
1	A	449	ILE	2.4
1	A	444	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	434	ASN	2.3
1	A	460	LEU	2.3
1	B	70	ASN	2.2
1	B	472	TYR	2.1
1	A	454	LYS	2.1
1	B	463	PHE	2.0
1	B	462	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYP	X	1	10/10	-	-	33,40,42,44	0
2	XYP	X	2	9/10	-	-	24,27,30,31	0
2	XYP	X	3	9/10	-	-	15,18,24,29	0
3	XYP	C	1	10/10	0.85	0.11	15,18,27,27	0
3	XYP	C	2	9/10	0.89	0.12	27,30,34,38	0
3	XYP	C	3	9/10	0.97	0.06	32,35,38,38	0
3	XYP	C	4	9/10	-	-	33,40,44,47	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain X:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain C:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPD	A	801	8/8	0.90	0.13	24,31,40,46	0
4	MPD	B	801	8/8	0.93	0.11	22,31,39,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.