



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:52 PM UTC

PDB ID : 3C8K / pdb_00003c8k
Title : The crystal structure of Ly49C bound to H-2Kb
Authors : Deng, L.; Mariuzza, R.A.
Deposited on : 2008-02-12
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	8% 68% 27% . .
2	B	99	2% 70% 19% 9% .
3	P	8	12% 62% 25% 12%
4	D	125	% 66% 24% 9% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	0	0
			2232	1408	393	422	9			

- Molecule 2 is a protein called beta-2 microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			818	523	138	150	7			

- Molecule 3 is a protein called Ovalbumin peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	8	Total	C	N	O	0	0	0
			68	45	10	13			

- Molecule 4 is a protein called Natural killer cell receptor Ly-49C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	125	Total	C	N	O	S	0	0	0
			1046	682	173	180	11			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	171	GLY	SER	engineered mutation	UNP Q61198
D	193	GLY	GLU	engineered mutation	UNP Q61198
D	223	LYS	ARG	engineered mutation	UNP Q61198


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	26	Total 26	O 26	0	0
5	B	21	Total 21	O 21	0	0
5	P	2	Total 2	O 2	0	0
5	D	16	Total 16	O 16	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 2: beta-2 microglobulin



- Molecule 3: Ovalbumin peptide



- Molecule 4: Natural killer cell receptor Ly-49C



C251	K255	D258	K259	F260	P261	D262
------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	152.01Å 152.01Å 64.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.90 30.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.6 (30.00-2.90) 95.5 (30.00-2.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.42 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.199 , 0.263 0.258 , 0.297	Depositor DCC
R_{free} test set	665 reflections (3.84%)	wwPDB-VP
Wilson B-factor (Å ²)	68.9	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4229	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	8/2293 (0.3%)	1.06	9/3113 (0.3%)
2	B	0.71	0/844	1.03	5/1144 (0.4%)
3	P	0.61	0/68	1.01	0/88
4	D	0.86	3/1077 (0.3%)	0.99	2/1445 (0.1%)
All	All	0.89	11/4282 (0.3%)	1.04	16/5790 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	PRO	C-N	19.01	1.60	1.33
1	A	58	GLU	CD-OE1	16.08	1.55	1.25
1	A	58	GLU	CG-CD	12.55	1.83	1.52
4	D	205	LYS	CD-CE	12.25	1.89	1.52
1	A	57	PRO	C-O	-8.10	1.13	1.24
1	A	55	GLU	CD-OE2	7.92	1.40	1.25
1	A	58	GLU	CD-OE2	7.82	1.40	1.25
1	A	58	GLU	CB-CG	7.18	1.74	1.52
4	D	205	LYS	CG-CD	5.73	1.69	1.52
1	A	55	GLU	CD-OE1	5.25	1.35	1.25
4	D	262	ASP	C-OXT	5.10	1.33	1.23

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	TYR	CA-C-N	16.78	140.81	119.84
1	A	209	TYR	C-N-CA	16.78	140.81	119.84
2	B	31	HIS	CA-C-N	9.27	129.93	120.38
2	B	31	HIS	C-N-CA	9.27	129.93	120.38
1	A	58	GLU	CG-CD-OE2	9.04	139.19	118.40
2	B	32	PRO	N-CA-C	-8.31	100.56	110.70
4	D	205	LYS	CG-CD-CE	-6.98	95.25	111.30
1	A	58	GLU	OE1-CD-OE2	-6.53	107.23	122.90
1	A	58	GLU	CB-CG-CD	-6.27	101.94	112.60
1	A	210	PRO	N-CA-C	-5.85	100.42	112.47
2	B	32	PRO	CA-C-N	-5.76	113.88	119.87
2	B	32	PRO	C-N-CA	-5.76	113.88	119.87
4	D	240	ILE	N-CA-C	-5.39	100.43	108.46
1	A	209	TYR	C-N-CD	-5.19	103.71	125.00
1	A	113	TYR	N-CA-C	5.05	117.64	109.06
1	A	209	TYR	N-CA-C	5.01	120.87	109.81

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	209	TYR	Peptide
2	B	31	HIS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2232	0	2123	58	0
2	B	818	0	797	26	0
3	P	68	0	74	4	0
4	D	1046	0	1033	33	0
5	A	26	0	0	0	0
5	B	21	0	0	0	0
5	D	16	0	0	0	0
5	P	2	0	0	0	0
All	All	4229	0	4027	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:GLU:CD	1:A:58:GLU:CG	1.83	1.50
4:D:205:LYS:CD	4:D:205:LYS:CE	1.89	1.48
4:D:153:PHE:HB3	4:D:155:MET:HE1	1.46	0.95
2:B:32:PRO:HD2	2:B:84:HIS:NE2	1.94	0.83
4:D:188:ARG:HG3	4:D:188:ARG:HH11	1.43	0.82
4:D:190:VAL:O	4:D:237:LYS:HE2	1.81	0.81
1:A:126:LEU:HD22	1:A:156:LEU:HD23	1.63	0.81
4:D:167:CYS:HB3	4:D:172:VAL:O	1.83	0.78
2:B:3:LYS:HD3	2:B:31:HIS:HB3	1.66	0.77
1:A:191:HIS:NE2	1:A:199:VAL:HG11	2.00	0.76
1:A:213:ILE:H	4:D:230:ARG:HH21	1.30	0.76
4:D:205:LYS:CE	4:D:205:LYS:CG	2.65	0.75
4:D:188:ARG:HH11	4:D:188:ARG:CG	2.00	0.75
4:D:240:ILE:HD13	4:D:240:ILE:N	2.04	0.72
2:B:32:PRO:HD2	2:B:84:HIS:HE2	1.56	0.70
2:B:32:PRO:HD3	2:B:84:HIS:CE1	2.27	0.69
1:A:210:PRO:HD2	1:A:263:HIS:NE2	2.08	0.69
2:B:74:GLU:HG2	2:B:75:THR:HG23	1.75	0.68
2:B:32:PRO:CD	2:B:84:HIS:CE1	2.79	0.65
4:D:172:VAL:HG13	4:D:255:LYS:HB2	1.79	0.65
1:A:35:ARG:HB3	1:A:48:ARG:HD2	1.79	0.64
1:A:23:MET:CE	1:A:35:ARG:HE	2.12	0.62
1:A:155:ARG:NH2	3:P:4:ASN:O	2.32	0.61
4:D:154:ILE:HB	4:D:251:CYS:HB3	1.82	0.61
4:D:164:LYS:HE3	4:D:211:ILE:HD11	1.82	0.60
1:A:110:LEU:HD13	4:D:226:ASN:HD21	1.66	0.60
4:D:205:LYS:CD	4:D:205:LYS:NZ	2.63	0.60
1:A:202:ARG:NH1	1:A:244:TRP:CZ3	2.70	0.59
1:A:234:ARG:HE	1:A:242:GLN:HG3	1.68	0.59
1:A:58:GLU:CD	1:A:58:GLU:CB	2.77	0.58
2:B:73:THR:HB	2:B:76:ASP:OD2	2.04	0.57
4:D:153:PHE:HB3	4:D:155:MET:CE	2.29	0.57
1:A:114:GLN:HG3	1:A:156:LEU:HD21	1.87	0.56
2:B:58:LYS:HZ3	4:D:241:GLU:HG3	1.71	0.56
1:A:52:MET:C	1:A:54:GLN:N	2.63	0.55
2:B:58:LYS:NZ	4:D:241:GLU:HG3	2.20	0.55
1:A:202:ARG:NH2	2:B:99:MET:HE2	2.21	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:GLU:OE2	1:A:61:GLU:HA	2.06	0.55
2:B:10:TYR:CD1	2:B:10:TYR:N	2.73	0.55
1:A:261:VAL:HB	1:A:270:LEU:HB2	1.89	0.55
2:B:2:GLN:HA	2:B:31:HIS:O	2.06	0.55
1:A:127:ASN:OD1	1:A:134:THR:HG23	2.08	0.54
1:A:47:PRO:O	1:A:48:ARG:HG2	2.09	0.53
1:A:229:GLU:HG2	1:A:230:LEU:H	1.75	0.52
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.44	0.51
1:A:23:MET:HE1	1:A:35:ARG:HE	1.74	0.51
1:A:52:MET:C	1:A:54:GLN:H	2.16	0.51
1:A:226:GLN:HG2	1:A:227:ASP:N	2.24	0.51
1:A:196:GLU:O	1:A:198:LYS:HG3	2.10	0.51
1:A:226:GLN:O	1:A:228:MET:N	2.43	0.51
2:B:32:PRO:CD	2:B:84:HIS:NE2	2.68	0.51
1:A:177:ALA:C	1:A:179:LEU:H	2.18	0.51
4:D:260:PHE:O	4:D:262:ASP:N	2.42	0.51
1:A:47:PRO:C	1:A:48:ARG:HG2	2.36	0.50
1:A:271:THR:C	1:A:272:LEU:HD12	2.36	0.50
2:B:21:ASN:HB3	2:B:70:PHE:CE1	2.46	0.50
4:D:221:LYS:HG2	4:D:225:MET:HE2	1.92	0.50
2:B:32:PRO:HD2	2:B:84:HIS:CE1	2.44	0.50
4:D:244:ASP:HB3	4:D:247:ILE:HG13	1.92	0.50
1:A:264:GLN:C	1:A:264:GLN:HE21	2.20	0.49
1:A:3:HIS:ND1	1:A:29:ASP:OD2	2.45	0.49
1:A:255:GLN:OE1	1:A:273:ARG:NH1	2.44	0.49
4:D:154:ILE:HD13	4:D:166:ASN:HD22	1.77	0.49
1:A:112:GLY:HA3	1:A:160:LEU:HD13	1.95	0.49
4:D:225:MET:HB3	4:D:227:PHE:CE1	2.47	0.48
1:A:219:LEU:HD13	1:A:257:TYR:CZ	2.49	0.48
1:A:128:GLU:H	1:A:128:GLU:CD	2.21	0.48
1:A:210:PRO:CD	1:A:263:HIS:NE2	2.77	0.48
2:B:59:ASP:OD1	2:B:59:ASP:C	2.57	0.48
2:B:69:GLU:CD	2:B:69:GLU:H	2.22	0.47
4:D:258:ASP:C	4:D:259:LYS:HG2	2.39	0.47
4:D:158:THR:O	4:D:248:PRO:HA	2.15	0.47
1:A:147:TRP:HB3	1:A:152:GLU:HB2	1.97	0.47
4:D:219:ASP:O	4:D:223:LYS:HB2	2.15	0.47
4:D:201:TYR:CZ	4:D:229:SER:HB3	2.50	0.46
1:A:202:ARG:HD2	1:A:204:TRP:CZ2	2.51	0.45
1:A:226:GLN:HG2	1:A:227:ASP:HB2	1.98	0.45
1:A:255:GLN:NE2	1:A:274:TRP:O	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ALA:HA	1:A:204:TRP:O	2.18	0.44
4:D:174:ILE:HG12	4:D:198:GLY:N	2.32	0.44
4:D:164:LYS:O	4:D:168:GLN:HG2	2.17	0.44
1:A:102:GLU:O	1:A:109:LEU:HD12	2.18	0.43
4:D:240:ILE:HD13	4:D:240:ILE:H	1.83	0.43
1:A:84:TYR:OH	3:P:8:LEU:OXT	2.36	0.43
4:D:188:ARG:HG3	4:D:188:ARG:NH1	2.23	0.43
1:A:71:GLU:HG2	1:A:75:ARG:NH2	2.34	0.43
1:A:101:CYS:HB2	1:A:109:LEU:HD13	2.01	0.43
1:A:173:LYS:HA	1:A:176:ASN:HD21	1.83	0.43
1:A:194:ARG:HB3	1:A:195:PRO:HD2	2.00	0.43
1:A:233:THR:OG1	1:A:243:LYS:HD2	2.19	0.43
4:D:190:VAL:HG13	4:D:191:ILE:O	2.19	0.43
1:A:113:TYR:CD1	1:A:113:TYR:N	2.86	0.42
1:A:106:ASP:OD1	1:A:106:ASP:N	2.53	0.42
2:B:29:GLN:HA	2:B:61:SER:HB2	2.02	0.42
1:A:117:ALA:HB2	2:B:60:TRP:CZ2	2.54	0.42
2:B:41:LYS:C	2:B:43:GLY:H	2.27	0.42
4:D:247:ILE:HA	4:D:248:PRO:HD3	1.88	0.42
1:A:163:THR:HG22	1:A:167:TRP:HD1	1.85	0.41
1:A:50:ARG:O	1:A:53:GLU:HG2	2.21	0.41
2:B:32:PRO:HD3	2:B:84:HIS:HE1	1.82	0.41
1:A:191:HIS:CD2	1:A:199:VAL:HG11	2.54	0.41
4:D:190:VAL:O	4:D:237:LYS:CE	2.60	0.41
1:A:32:GLU:OE2	1:A:48:ARG:HD3	2.21	0.41
2:B:76:ASP:OD2	2:B:76:ASP:N	2.53	0.41
1:A:159:TYR:CG	3:P:3:ILE:HD12	2.56	0.41
2:B:19:LYS:HE2	2:B:21:ASN:ND2	2.36	0.41
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.90	0.41
3:P:3:ILE:CG1	3:P:4:ASN:N	2.83	0.41
1:A:226:GLN:C	1:A:228:MET:H	2.29	0.40
2:B:14:PRO:HA	2:B:15:PRO:HD3	1.92	0.40
2:B:39:MET:HG3	2:B:49:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/274 (99%)	243 (89%)	24 (9%)	5 (2%)	6	25
2	B	97/99 (98%)	87 (90%)	5 (5%)	5 (5%)	1	5
3	P	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
4	D	123/125 (98%)	113 (92%)	9 (7%)	1 (1%)	16	44
All	All	498/506 (98%)	448 (90%)	39 (8%)	11 (2%)	5	20

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	58	LYS
1	A	197	ASP
1	A	267	PRO
2	B	60	TRP
4	D	261	PRO
1	A	210	PRO
1	A	227	ASP
2	B	32	PRO
1	A	226	GLN
2	B	48	LYS
2	B	43	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	210 (90%)	22 (10%)	8	26
2	B	93/93 (100%)	84 (90%)	9 (10%)	8	25
3	P	8/8 (100%)	7 (88%)	1 (12%)	4	15
4	D	114/114 (100%)	100 (88%)	14 (12%)	4	15
All	All	447/447 (100%)	401 (90%)	46 (10%)	7	23

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	32	GLU
1	A	54	GLN
1	A	58	GLU
1	A	61	GLU
1	A	78	LEU
1	A	111	ARG
1	A	113	TYR
1	A	114	GLN
1	A	128	GLU
1	A	141	LEU
1	A	146	LYS
1	A	152	GLU
1	A	169	ARG
1	A	170	ARG
1	A	182	THR
1	A	199	VAL
1	A	200	THR
1	A	248	VAL
1	A	251	LEU
1	A	264	GLN
1	A	273	ARG
2	B	1	ILE
2	B	2	GLN
2	B	3	LYS
2	B	16	GLU
2	B	19	LYS
2	B	48	LYS
2	B	73	THR
2	B	74	GLU
2	B	77	THR
3	P	8	LEU
4	D	155	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	158	THR
4	D	168	GLN
4	D	178	GLU
4	D	188	ARG
4	D	190	VAL
4	D	203	LYS
4	D	211	ILE
4	D	213	ASN
4	D	218	LEU
4	D	223	LYS
4	D	226	ASN
4	D	240	ILE
4	D	259	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	176	ASN
1	A	242	GLN
1	A	264	GLN
2	B	6	GLN
2	B	29	GLN
2	B	31	HIS
4	D	166	ASN
4	D	194	ASN
4	D	226	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/274 (100%)	1.03	22 (8%) 18 15	72, 86, 103, 106	0
2	B	99/99 (100%)	0.62	2 (2%) 65 56	58, 69, 78, 81	0
3	P	8/8 (100%)	0.75	1 (12%) 8 7	75, 77, 80, 81	0
4	D	125/125 (100%)	0.16	1 (0%) 82 77	44, 53, 62, 72	0
All	All	506/506 (100%)	0.73	26 (5%) 33 26	44, 78, 100, 106	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	36	PHE	3.6
1	A	225	ILE	3.2
3	P	1	SER	2.8
1	A	52	MET	2.7
1	A	193	SER	2.6
2	B	17	ASN	2.5
1	A	252	GLY	2.4
1	A	45	TYR	2.4
1	A	172	LEU	2.4
1	A	72	GLN	2.4
1	A	231	VAL	2.4
1	A	113	TYR	2.3
1	A	224	LEU	2.3
1	A	42	ASN	2.3
1	A	95	ILE	2.3
1	A	43	PRO	2.3
1	A	9	VAL	2.2
1	A	217	TRP	2.2
1	A	171	TYR	2.2
1	A	115	GLN	2.1
1	A	197	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	168	LEU	2.1
1	A	211	ALA	2.1
1	A	256	TYR	2.1
4	D	220	MET	2.1
2	B	36	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.