



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:44 AM UTC

PDB ID : 3C9F / pdb_00003c9f
Title : Crystal structure of 5'-nucleotidase from *Candida albicans* SC5314
Authors : Patskovsky, Y.; Romero, R.; Gilmore, M.; Eberle, M.; Bain, K.; Smith, D.; Wasserman, S.R.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2008-02-15
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	5% 88% 6% • 5%
1	B	557	3% 91% 5% • •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9842 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-nucleotidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	531	Total	C	N	O	S	0	18	0
			4457	2886	748	817	6			
1	B	540	Total	C	N	O	S	0	19	0
			4538	2932	767	832	7			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	MET	-	expression tag	UNP Q5A5Q7
A	13	SER	-	expression tag	UNP Q5A5Q7
A	14	LEU	-	expression tag	UNP Q5A5Q7
A	62	MET	LEU	engineered mutation	UNP Q5A5Q7
A	151	LEU	SER	engineered mutation	UNP Q5A5Q7
A	157	LEU	SER	engineered mutation	UNP Q5A5Q7
A	172	MET	LEU	engineered mutation	UNP Q5A5Q7
A	323	LEU	SER	engineered mutation	UNP Q5A5Q7
A	367	LEU	SER	engineered mutation	UNP Q5A5Q7
A	561	GLU	-	expression tag	UNP Q5A5Q7
A	562	GLY	-	expression tag	UNP Q5A5Q7
A	563	HIS	-	expression tag	UNP Q5A5Q7
A	564	HIS	-	expression tag	UNP Q5A5Q7
A	565	HIS	-	expression tag	UNP Q5A5Q7
A	566	HIS	-	expression tag	UNP Q5A5Q7
A	567	HIS	-	expression tag	UNP Q5A5Q7
A	568	HIS	-	expression tag	UNP Q5A5Q7
B	12	MET	-	expression tag	UNP Q5A5Q7
B	13	SER	-	expression tag	UNP Q5A5Q7
B	14	LEU	-	expression tag	UNP Q5A5Q7
B	62	MET	LEU	engineered mutation	UNP Q5A5Q7
B	151	LEU	SER	engineered mutation	UNP Q5A5Q7
B	157	LEU	SER	engineered mutation	UNP Q5A5Q7
B	172	MET	LEU	engineered mutation	UNP Q5A5Q7
B	323	LEU	SER	engineered mutation	UNP Q5A5Q7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	367	LEU	SER	engineered mutation	UNP Q5A5Q7
B	561	GLU	-	expression tag	UNP Q5A5Q7
B	562	GLY	-	expression tag	UNP Q5A5Q7
B	563	HIS	-	expression tag	UNP Q5A5Q7
B	564	HIS	-	expression tag	UNP Q5A5Q7
B	565	HIS	-	expression tag	UNP Q5A5Q7
B	566	HIS	-	expression tag	UNP Q5A5Q7
B	567	HIS	-	expression tag	UNP Q5A5Q7
B	568	HIS	-	expression tag	UNP Q5A5Q7


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	B	3	Total	Na	0	0
			3	3		

- Molecule 4 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	380	Total	O	0	1
			381	381		
5	B	453	Total	O	0	1
			454	454		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 5'-nucleotidase



• Molecule 1: 5'-nucleotidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.95Å 106.74Å 150.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 20.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-1.90) 99.3 (20.00-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, R_{free}	0.186 , 0.225 0.188 , 0.226	Depositor DCC
R_{free} test set	4265 reflections (2.98%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9842	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NA, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/4629	0.86	4/6294 (0.1%)
1	B	0.67	0/4718	0.84	4/6414 (0.1%)
All	All	0.65	0/9347	0.85	8/12708 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	466[A]	MET	CA-C-O	7.08	130.92	122.36
1	B	466[B]	MET	CA-C-O	7.08	130.92	122.36
1	A	198	GLU	CA-C-N	5.42	126.61	119.84
1	A	198	GLU	C-N-CA	5.42	126.61	119.84
1	A	353	VAL	N-CA-C	-5.29	104.85	109.19
1	B	198	GLU	CA-C-N	5.28	124.74	119.24
1	B	198	GLU	C-N-CA	5.28	124.74	119.24
1	A	262	SER	N-CA-C	5.07	116.86	108.13

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4457	0	4399	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4538	0	4459	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	3	0	0	0	0
4	A	3	0	1	0	0
4	B	3	0	1	0	0
5	A	381	0	0	2	0
5	B	454	0	0	2	0
All	All	9842	0	8860	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188[A]:ARG:HH11	1:B:467:GLN:HB2	1.37	0.85
1:A:188[B]:ARG:HH22	1:B:467:GLN:HA	1.42	0.84
1:A:140[B]:CYS:SG	1:A:143:VAL:HB	2.42	0.59
1:A:188[A]:ARG:NH1	1:B:467:GLN:HB2	2.15	0.57
1:B:82[A]:HIS:CD2	1:B:94:ASN:HD22	2.24	0.55
1:B:203:GLU:OE1	1:B:206:LYS:HE2	2.10	0.51
1:B:376:SER:HB2	1:B:379:GLU:HB2	1.92	0.50
1:A:373:LEU:HD13	1:A:552:LEU:HD11	1.94	0.50
1:B:437:ILE:O	1:B:437:ILE:HG13	2.08	0.50
1:A:188[B]:ARG:HH12	1:B:467:GLN:HB2	1.75	0.50
1:A:289:PRO:HD2	1:A:292:GLN:HE21	1.78	0.48
1:B:160:LYS:HE2	5:B:668:HOH:O	2.13	0.48
1:B:317:ASP:HB3	1:B:322:LYS:HD2	1.96	0.48
1:B:291:ARG:NH1	5:B:789:HOH:O	2.46	0.47
1:A:36:HIS:NE2	5:A:975[B]:HOH:O	2.36	0.46
1:A:371:LYS:HG3	5:A:659:HOH:O	2.16	0.46
1:B:82[A]:HIS:NE2	1:B:115:HIS:O	2.49	0.45
1:A:531:LYS:HA	1:A:531:LYS:HD3	1.81	0.45
1:B:67:HIS:CD2	1:B:166:ILE:HG13	2.53	0.44
1:B:466[B]:MET:HB2	1:B:466[B]:MET:HE2	1.74	0.44
1:A:220:ILE:HG23	1:A:247:GLY:HA3	1.99	0.43
1:A:188[A]:ARG:HH11	1:B:467:GLN:CB	2.19	0.42
1:A:198:GLU:HA	1:A:199:PRO:HD2	1.84	0.42
1:A:417[B]:ILE:HD11	1:A:541:PRO:HG3	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:THR:HA	1:B:517:VAL:O	2.21	0.41
1:B:469:LYS:H	1:B:469:LYS:HG2	1.57	0.41
1:A:305:SER:OG	1:A:486:ASP:OD2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	543/557 (98%)	523 (96%)	19 (4%)	1 (0%)	43	36
1	B	553/557 (99%)	531 (96%)	20 (4%)	2 (0%)	30	22
All	All	1096/1114 (98%)	1054 (96%)	39 (4%)	3 (0%)	36	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	543	PRO
1	A	84	GLY
1	B	84	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	505/509 (99%)	481 (95%)	24 (5%)	23	15
1	B	514/509 (101%)	500 (97%)	14 (3%)	39	34
All	All	1019/1018 (100%)	981 (96%)	38 (4%)	30	22

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	SER
1	A	76	ILE
1	A	92	SER
1	A	125	GLN
1	A	135	GLN
1	A	136	ASP
1	A	171	VAL
1	A	208[A]	GLU
1	A	208[B]	GLU
1	A	240	ASP
1	A	256[A]	THR
1	A	256[B]	THR
1	A	283	LYS
1	A	292	GLN
1	A	305	SER
1	A	314	LYS
1	A	416	ASN
1	A	453	GLN
1	A	467	GLN
1	A	469	LYS
1	A	531	LYS
1	A	536	SER
1	A	538	GLU
1	A	561	GLU
1	B	92	SER
1	B	283	LYS
1	B	311	ASN
1	B	377	LYS
1	B	413	ILE
1	B	416	ASN
1	B	437	ILE
1	B	450	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	455	LEU
1	B	461	GLN
1	B	467	GLN
1	B	469	LYS
1	B	531	LYS
1	B	540	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	125	GLN
1	A	287	ASN
1	A	292	GLN
1	A	451	GLN
1	A	477	HIS
1	A	554	ASN
1	B	21	ASN
1	B	461	GLN
1	B	525	ASN
1	B	559	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FMT	B	607	2	2,2,2	0.75	0	1,1,1	0.47	0
4	FMT	A	607	2	2,2,2	0.47	0	1,1,1	0.54	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	531/557 (95%)	0.06	28 (5%) 32 34	15, 36, 69, 115	18 (3%)
1	B	540/557 (96%)	-0.22	17 (3%) 51 55	15, 32, 62, 121	19 (3%)
All	All	1071/1114 (96%)	-0.08	45 (4%) 40 43	15, 34, 68, 121	37 (3%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	455	LEU	8.4
1	B	466[A]	MET	6.3
1	A	436	TYR	5.2
1	A	17	PHE	5.1
1	B	437	ILE	5.0
1	B	452	VAL	4.1
1	B	467	GLN	3.8
1	A	452	VAL	3.6
1	B	440[A]	SER	3.6
1	A	560	ASN	3.3
1	A	18	PRO	3.2
1	A	433	ASP	3.2
1	B	457	THR	3.2
1	B	453	GLN	3.1
1	A	440[A]	SER	3.0
1	A	43[A]	ILE	3.0
1	A	16	SER	2.9
1	B	531	LYS	2.9
1	B	378	HIS	2.8
1	A	228	TYR	2.7
1	A	286	LEU	2.7
1	A	450	LEU	2.7
1	A	437	ILE	2.7
1	A	453	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	467	GLN	2.7
1	B	417	ILE	2.6
1	B	439	ALA	2.6
1	A	561	GLU	2.5
1	A	438[A]	SER	2.5
1	B	460	HIS	2.4
1	A	439	ALA	2.4
1	A	469	LYS	2.4
1	A	451	GLN	2.3
1	A	279	VAL	2.3
1	B	456	SER	2.2
1	B	559	ASN	2.2
1	A	281	LEU	2.2
1	A	240	ASP	2.2
1	B	543	PRO	2.2
1	A	161	TYR	2.2
1	A	284	ALA	2.2
1	A	166	ILE	2.1
1	A	470	LEU	2.1
1	B	43[A]	ILE	2.0
1	A	537	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NA	B	603	1/1	0.88	0.12	59,59,59,59	0
3	NA	B	606	1/1	0.88	0.10	54,54,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NA	B	604	1/1	0.94	0.07	44,44,44,44	0
3	NA	A	605	1/1	0.94	0.05	44,44,44,44	0
4	FMT	B	607	3/3	0.95	0.08	33,33,37,38	0
4	FMT	A	607	3/3	0.97	0.06	26,26,27,36	0
2	ZN	A	601	1/1	1.00	0.03	32,32,32,32	0
2	ZN	B	601	1/1	1.00	0.03	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.