



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 08:05 PM EDT

PDB ID : 4C9B / pdb_00004c9b
Title : Crystal structure of eIF4AIII-CWC22 complex
Authors : Buchwald, G.; Schuessler, S.; Basquin, C.; LeHir, H.; Conti, E.
Deposited on : 2013-10-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	
2	B	291	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1413	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5751 atoms, of which 56 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called EUKARYOTIC INITIATION FACTOR 4A-III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	389	3124	1973	547	584	20	0	7	0

- Molecule 2 is a protein called PRE-MRNA-SPLICING FACTOR CWC22 HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	277	2214	1419	386	400	9	0	2	0

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O P		
3	A	1	5	4 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 6	C 3	O 3		0	0
4	A	1	Total 6	C 3	O 3		0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0
4	B	1	Total 14	C 3	H 8	O 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	169	Total	O	0	0
			169	169		
5	B	129	Total	O	0	0
			129	129		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: EUKARYOTIC INITIATION FACTOR 4A-III



- Molecule 2: PRE-MRNA-SPLICING FACTOR CWC22 HOMOLOG



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.29Å 108.72Å 61.37Å 90.00° 99.94° 90.00°	Depositor
Resolution (Å)	55.45 – 2.00 55.45 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (55.45-2.00) 99.8 (55.45-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.00Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.202 , 0.236 0.216 , 0.252	Depositor DCC
R_{free} test set	2451 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.975	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5751	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/3192	0.77	0/4310
2	B	0.41	0/2255	0.73	2/3044 (0.1%)
All	All	0.40	0/5447	0.75	2/7354 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	135	PRO	CA-C-N	5.58	125.68	119.32
2	B	135	PRO	C-N-CA	5.58	125.68	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3124	0	3151	63	1
2	B	2214	0	2284	48	1
3	A	5	0	0	0	0
4	A	48	48	64	10	0
4	B	6	8	8	0	0
5	A	169	0	0	2	0
5	B	129	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5695	56	5507	108	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354:LEU:HD11	1:A:381:VAL:HG12	1.34	1.08
1:A:307:MET:HE1	1:A:320:MET:HE2	1.48	0.95
1:A:59:GLU:HB2	4:A:1413:GOL:H32	1.47	0.95
1:A:148:GLU:HG3	1:A:151:ARG:HH12	1.40	0.87
1:A:383:ASN:HA	1:A:386:ILE:HG13	1.57	0.84
2:B:152:LEU:HA	2:B:387:MET:HE1	1.62	0.79
1:A:99:CYS:SG	2:B:330:LYS:NZ	2.55	0.79
2:B:264:ILE:HG13	2:B:274:CYS:SG	2.25	0.76
2:B:124:PRO:HA	2:B:128:ARG:HH21	1.56	0.70
2:B:279:THR:O	2:B:283:GLU:HG3	1.96	0.66
1:A:59:GLU:HB2	4:A:1413:GOL:C3	2.25	0.65
1:A:79:ILE:HD12	1:A:238[B]:ARG:HG2	1.79	0.65
2:B:264:ILE:HD13	2:B:269:ALA:HB3	1.79	0.63
1:A:251:ILE:HD11	1:A:367:ARG:HG3	1.80	0.62
1:A:59:GLU:CB	4:A:1413:GOL:H32	2.28	0.62
1:A:142:GLY:HA2	1:A:146:VAL:HG22	1.81	0.62
2:B:152:LEU:HA	2:B:387:MET:CE	2.30	0.62
2:B:225:LYS:HG2	2:B:405:ILE:HG12	1.82	0.60
1:A:28:PHE:HE2	1:A:82:SER:HB3	1.66	0.60
1:A:59:GLU:H	4:A:1413:GOL:H12	1.65	0.60
1:A:99:CYS:HB3	5:B:2097:HOH:O	2.02	0.60
1:A:28:PHE:CE2	1:A:82:SER:HB3	2.38	0.59
1:A:79:ILE:HD12	1:A:238[A]:ARG:HG2	1.85	0.57
2:B:225:LYS:HE3	2:B:405:ILE:CG1	2.33	0.57
1:A:114:PRO:HD2	1:A:118:LEU:HD23	1.85	0.57
1:A:170:MET:HA	1:A:170:MET:HE2	1.87	0.57
2:B:225:LYS:HE3	2:B:405:ILE:HG12	1.86	0.57
1:A:354:LEU:HD21	1:A:379:ASN:HB3	1.86	0.56
2:B:287:ASP:HB3	2:B:331:ARG:CZ	2.35	0.56
2:B:218:LEU:HD23	2:B:218:LEU:C	2.31	0.56
2:B:316:PHE:CE2	2:B:343:ARG:HD3	2.40	0.56
1:A:317:GLU:O	1:A:321:LYS:HG3	2.06	0.55
1:A:88:LYS:HD2	1:A:217:ILE:HG23	1.88	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:GLU:CG	1:A:151:ARG:HH12	2.17	0.55
2:B:374:ASP:HB2	2:B:376:TYR:CZ	2.41	0.54
1:A:390:ARG:O	1:A:394:GLN:HG3	2.08	0.54
1:A:59:GLU:H	4:A:1413:GOL:C1	2.21	0.54
1:A:266:ASP:OD2	2:B:136:PRO:HB2	2.08	0.54
4:A:1413:GOL:H11	5:A:2168:HOH:O	2.08	0.54
1:A:405:MET:SD	2:B:129:THR:HG22	2.48	0.53
1:A:138:HIS:CE1	1:A:152:LYS:HD2	2.44	0.52
1:A:225:ILE:O	1:A:229:THR:HG23	2.10	0.52
2:B:234:LEU:HB3	2:B:370:LEU:HD12	1.91	0.52
1:A:249:GLU:O	1:A:367:ARG:HD3	2.10	0.52
2:B:139:LEU:HD23	2:B:139:LEU:O	2.10	0.52
1:A:308:HIS:CE1	1:A:310:ASP:HB2	2.46	0.51
1:A:354:LEU:HD12	1:A:385:ASP:HB3	1.92	0.51
1:A:22:ASP:OD1	1:A:24:THR:HG22	2.11	0.51
1:A:336:VAL:HG23	1:A:337:TRP:CD1	2.45	0.51
1:A:348:LEU:HD11	1:A:378:ILE:HG13	1.93	0.50
2:B:166:ILE:O	2:B:170:ILE:HG12	2.11	0.50
2:B:251[B]:GLN:NE2	5:B:2040:HOH:O	2.45	0.50
1:A:385:ASP:O	1:A:388:ILE:HG22	2.12	0.50
4:A:1420:GOL:H11	2:B:172:LYS:HE3	1.92	0.50
2:B:294:ILE:HD13	2:B:335:MET:C	2.37	0.49
2:B:369:MET:O	2:B:369:MET:HE3	2.13	0.49
1:A:388:ILE:O	1:A:392:ILE:HG12	2.11	0.49
2:B:151:SER:O	2:B:155:GLN:HG3	2.13	0.49
2:B:152:LEU:HA	2:B:387:MET:SD	2.53	0.48
1:A:256:VAL:HG22	1:A:403:MET:HE2	1.95	0.48
2:B:352:ILE:HG22	2:B:353:ILE:HG13	1.96	0.48
2:B:294:ILE:HG22	2:B:298:LYS:HE3	1.94	0.48
2:B:234:LEU:HD21	2:B:263:LEU:HD13	1.96	0.48
1:A:168:PHE:CE1	1:A:203:ASP:HB3	2.49	0.48
2:B:134:ILE:HG23	2:B:138:LYS:HD3	1.96	0.48
2:B:260:VAL:O	2:B:264:ILE:HG12	2.13	0.48
1:A:311:MET:O	1:A:316:ARG:NH1	2.47	0.47
2:B:294:ILE:HD13	2:B:335:MET:O	2.13	0.47
1:A:33:GLU:OE2	1:A:373:ARG:NE	2.47	0.47
1:A:107:THR:HA	1:A:157:GLN:O	2.13	0.47
2:B:228:GLN:HG2	2:B:229:ILE:N	2.29	0.47
1:A:143:GLY:O	1:A:146:VAL:HG23	2.15	0.47
1:A:148:GLU:HG3	1:A:151:ARG:NH1	2.19	0.47
1:A:361:TYR:CE2	1:A:365:ILE:HD11	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:ARG:HB3	5:A:2164:HOH:O	2.15	0.46
1:A:149:ASP:O	1:A:153:LEU:HG	2.15	0.46
2:B:124:PRO:HA	2:B:128:ARG:NH2	2.28	0.46
1:A:383:ASN:HA	1:A:386:ILE:CG1	2.38	0.46
2:B:139:LEU:HD23	2:B:139:LEU:C	2.41	0.46
2:B:304:LEU:HA	2:B:304:LEU:HD23	1.77	0.45
1:A:39:THR:OG1	1:A:42:THR:HG23	2.16	0.45
2:B:399:LYS:O	2:B:403:LYS:HG3	2.17	0.45
2:B:234:LEU:CD2	2:B:263:LEU:HD13	2.46	0.45
2:B:294:ILE:HD12	2:B:335:MET:HB3	1.98	0.45
1:A:246:LEU:HA	1:A:367:ARG:O	2.17	0.45
2:B:294:ILE:CD1	2:B:335:MET:HB3	2.47	0.44
1:A:22:ASP:CG	1:A:24:THR:HG22	2.43	0.44
1:A:269:CYS:HB3	4:A:1420:GOL:H2	1.98	0.44
1:A:383:ASN:CA	1:A:386:ILE:HG13	2.40	0.44
1:A:60:LYS:HB3	4:A:1413:GOL:O3	2.17	0.44
2:B:387:MET:HE3	2:B:387:MET:HA	2.00	0.43
2:B:182:ILE:O	2:B:186:LEU:HG	2.18	0.43
1:A:82:SER:O	1:A:88:LYS:HE3	2.18	0.43
1:A:203:ASP:OD1	1:A:206:ARG:NH2	2.49	0.43
1:A:320:MET:HE1	1:A:331:LEU:HD21	2.01	0.43
2:B:231:GLU:O	2:B:235:LYS:HG3	2.19	0.42
2:B:322:ILE:O	2:B:326:SER:HB2	2.19	0.42
1:A:398:THR:OG1	1:A:399:GLN:N	2.52	0.42
2:B:387:MET:CE	2:B:387:MET:HA	2.50	0.42
1:A:273:ASP:HB3	2:B:164:LYS:HB3	2.02	0.42
1:A:188:GLU:OE2	4:A:1419:GOL:O2	2.34	0.42
2:B:284:ARG:O	2:B:284:ARG:HG3	2.19	0.41
1:A:359:GLU:HA	1:A:396:TYR:OH	2.20	0.41
2:B:155:GLN:CD	2:B:387:MET:HE2	2.45	0.41
1:A:151:ARG:HH22	1:A:152:LYS:HE2	1.85	0.41
1:A:307:MET:HE1	1:A:320:MET:CE	2.33	0.41
1:A:260:ARG:HB3	1:A:262:GLU:CD	2.46	0.41
2:B:215:TYR:O	2:B:219:VAL:HG23	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288[A]:ARG:NH1	2:B:276:GLU:OE2[1_554]	2.06	0.14

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/411 (95%)	385 (98%)	7 (2%)	0	100	100
2	B	275/291 (94%)	270 (98%)	5 (2%)	0	100	100
All	All	667/702 (95%)	655 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/361 (94%)	338 (99%)	3 (1%)	70	78
2	B	244/263 (93%)	243 (100%)	1 (0%)	84	89
All	All	585/624 (94%)	581 (99%)	4 (1%)	76	82

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	LYS
1	A	105	ARG
1	A	110	LEU
2	B	387	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	253	GLN
1	A	383	ASN
1	A	406	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	1412	-	4,4,4	1.01	0	6,6,6	0.53	0
4	GOL	B	1407	-	5,5,5	0.34	0	5,5,5	0.51	0
4	GOL	A	1415	-	5,5,5	0.41	0	5,5,5	0.29	0
4	GOL	A	1414	-	5,5,5	0.32	0	5,5,5	0.55	0
4	GOL	A	1416	-	5,5,5	0.37	0	5,5,5	0.27	0
4	GOL	A	1420	-	5,5,5	0.39	0	5,5,5	0.53	0
4	GOL	A	1419	-	5,5,5	0.36	0	5,5,5	0.35	0
4	GOL	A	1418	-	5,5,5	0.37	0	5,5,5	0.21	0
4	GOL	A	1417	-	5,5,5	0.35	0	5,5,5	0.34	0
4	GOL	A	1413	-	5,5,5	0.36	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	1407	-	-	2/4/4/4	-
4	GOL	A	1415	-	-	0/4/4/4	-
4	GOL	A	1414	-	-	2/4/4/4	-
4	GOL	A	1416	-	-	2/4/4/4	-
4	GOL	A	1420	-	-	4/4/4/4	-
4	GOL	A	1419	-	-	2/4/4/4	-
4	GOL	A	1418	-	-	0/4/4/4	-
4	GOL	A	1417	-	-	0/4/4/4	-
4	GOL	A	1413	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1414	GOL	C1-C2-C3-O3
4	A	1414	GOL	O2-C2-C3-O3
4	A	1416	GOL	C1-C2-C3-O3
4	A	1420	GOL	O1-C1-C2-C3
4	A	1413	GOL	C1-C2-C3-O3
4	A	1419	GOL	C1-C2-C3-O3
4	A	1420	GOL	C1-C2-C3-O3
4	A	1416	GOL	O2-C2-C3-O3
4	A	1420	GOL	O1-C1-C2-O2
4	A	1413	GOL	O2-C2-C3-O3
4	A	1419	GOL	O2-C2-C3-O3
4	B	1407	GOL	O2-C2-C3-O3
4	A	1420	GOL	O2-C2-C3-O3
4	B	1407	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1420	GOL	2	0
4	A	1419	GOL	1	0
4	A	1413	GOL	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	389/411 (94%)	0.89	38 (9%) 13 12	23, 40, 66, 94	7 (1%)
2	B	277/291 (95%)	0.81	28 (10%) 12 11	20, 37, 77, 95	2 (0%)
All	All	666/702 (94%)	0.86	66 (9%) 13 11	20, 39, 68, 95	9 (1%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	TRP	7.8
1	A	411	ILE	6.0
2	B	405	ILE	4.3
1	A	341	LEU	4.2
1	A	23	MET	4.1
1	A	410	LEU	4.1
1	A	340	GLY	4.0
2	B	123	ASP	3.7
2	B	406	LEU	3.6
1	A	246	LEU	3.5
2	B	274	CYS	3.5
2	B	325	GLU	3.3
2	B	129	THR	3.2
1	A	24	THR	3.2
2	B	125	LEU	3.1
1	A	155	TYR	3.1
2	B	124	PRO	3.1
2	B	139	LEU	3.0
1	A	35	ASP	3.0
1	A	248	LEU	3.0
1	A	313	GLN	2.9
2	B	127	THR	2.9
2	B	326	SER	2.8
1	A	258	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	211	ALA	2.8
1	A	336	VAL	2.7
2	B	134	ILE	2.6
1	A	104	VAL	2.5
1	A	225	ILE	2.5
2	B	128	ARG	2.5
2	B	307	VAL	2.5
1	A	371	TYR	2.5
2	B	150	ASN	2.5
2	B	136	PRO	2.5
2	B	141	MET	2.5
1	A	396	TYR	2.5
1	A	207	TYR	2.4
1	A	172[A]	ARG	2.4
1	A	343	VAL	2.4
1	A	146	VAL	2.3
1	A	388	ILE	2.3
1	A	392	ILE	2.3
2	B	328	ILE	2.3
1	A	229	THR	2.3
2	B	345	ASP	2.3
2	B	185	LEU	2.3
1	A	386	ILE	2.2
1	A	148	GLU	2.2
2	B	404	GLU	2.2
1	A	398	THR	2.2
1	A	26	VAL	2.2
2	B	392	MET	2.2
1	A	354	LEU	2.1
1	A	84	SER	2.1
2	B	387	MET	2.1
2	B	390	ASN	2.1
2	B	137	ALA	2.1
2	B	149	LYS	2.1
2	B	389	PRO	2.1
1	A	223	HIS	2.1
1	A	150	ILE	2.1
1	A	22	ASP	2.1
1	A	141	ILE	2.1
2	B	130	GLY	2.0
1	A	219	ALA	2.0
1	A	404	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	1417	6/6	0.75	0.14	59,71,74,74	0
4	GOL	B	1407	6/6	0.75	0.19	61,73,77,78	0
4	GOL	A	1414	6/6	0.76	0.21	50,53,54,55	0
4	GOL	A	1415	6/6	0.77	0.13	62,74,76,76	0
4	GOL	A	1420	6/6	0.78	0.16	53,64,67,68	0
3	PO4	A	1412	5/5	0.79	0.16	56,57,57,58	0
4	GOL	A	1419	6/6	0.79	0.14	71,85,86,86	0
4	GOL	A	1413	6/6	0.81	0.14	47,49,50,51	0
4	GOL	A	1418	6/6	0.82	0.15	66,80,81,81	0
4	GOL	A	1416	6/6	0.82	0.12	60,72,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.