



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 08:05 PM UTC

PDB ID : 4C9D / pdb_00004c9d
Title : Cas6 (TTHB231) product complex
Authors : Jinek, M.; Niewoehner, O.; Doudna, J.A.
Deposited on : 2013-10-02
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)
RNA backbone	3983	1109 (3.20-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	14% 73% 22% • •
1	B	268	11% 69% 26% • •
2	C	29	17% 14% 21% 10% 55%
2	D	29	21% 14% 17% 17% 52%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9032 atoms, of which 4407 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CAS6B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	262	Total	C	H	N	O	S	0	0	0
			4097	1322	2065	353	355	2			
1	B	260	Total	C	H	N	O	S	0	0	0
			4103	1317	2073	358	353	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q53VU8
A	-2	ALA	-	expression tag	UNP Q53VU8
A	-1	ALA	-	expression tag	UNP Q53VU8
A	0	SER	-	expression tag	UNP Q53VU8
B	-3	GLY	-	expression tag	UNP Q53VU8
B	-2	ALA	-	expression tag	UNP Q53VU8
B	-1	ALA	-	expression tag	UNP Q53VU8
B	0	SER	-	expression tag	UNP Q53VU8

- Molecule 2 is a RNA chain called R3 REPEAT RNA CLEAVAGE PRODUCT.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	13	Total	C	H	N	O	P	0	0	1
			392	115	129	48	87	13			
2	D	14	Total	C	H	N	O	P	0	0	1
			425	125	140	53	93	14			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	O	0	0
			9	9		
3	B	5	Total	O	0	0
			5	5		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CAS6B




• Molecule 1: CAS6B



• Molecule 2: R3 REPEAT RNA CLEAVAGE PRODUCT



• Molecule 2: R3 REPEAT RNA CLEAVAGE PRODUCT



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	75.22Å 75.22Å 308.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.08 – 3.00 73.08 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (73.08-3.00) 99.9 (73.08-3.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 3.01Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.244 , 0.291 0.243 , 0.281	Depositor DCC
R_{free} test set	937 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	57.1	Xtriage
Anisotropy	0.670	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 68.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	9032	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 23G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2091	0.66	0/2839
1	B	0.36	0/2088	0.69	0/2835
2	C	0.25	0/264	0.39	0/411
2	D	0.24	0/289	0.36	0/450
All	All	0.34	0/4732	0.64	0/6535

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2032	2065	2059	43	0
1	B	2030	2073	2067	46	0
2	C	263	129	129	9	0
2	D	285	140	140	10	0
3	A	9	0	0	1	0
3	B	5	0	0	0	0
3	C	1	0	0	0	0
All	All	4625	4407	4395	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:ARG:NH1	3:A:2003:HOH:O	2.18	0.75
1:A:141:PRO:HG2	1:A:209:VAL:HG21	1.69	0.72
1:B:32:ARG:NH1	1:B:89:GLU:OE1	2.22	0.72
2:C:24:A:H1'	2:C:25:G:C8	2.31	0.66
1:B:141:PRO:HG2	1:B:209:VAL:HG11	1.80	0.64
2:D:20:U:H2'	2:D:21:C:H5'	1.79	0.62
1:A:21:TYR:HB3	1:A:97:LEU:HB2	1.82	0.60
1:A:20:ARG:NE	1:A:212:GLY:O	2.35	0.60
1:B:114:HIS:CD2	1:B:115:PRO:HD2	2.37	0.59
1:B:48:PRO:HG2	1:B:79:LEU:HD11	1.84	0.58
1:B:68:LEU:HD23	1:B:70:LEU:HD11	1.86	0.58
1:B:213:LYS:HB2	2:D:16:A:C6	2.39	0.57
1:A:194:LEU:HB3	1:B:152:ARG:NH1	2.19	0.57
1:B:254:THR:N	1:B:255:PRO:CD	2.68	0.56
2:D:20:U:C2'	2:D:21:C:H5'	2.36	0.55
1:A:209:VAL:HG12	1:A:210:GLN:N	2.21	0.55
1:A:254:THR:N	1:A:255:PRO:CD	2.70	0.55
1:A:6:VAL:HG21	1:A:114:HIS:CD2	2.43	0.54
1:A:74:THR:OG1	1:A:79:LEU:HB3	2.07	0.54
1:B:81:ALA:HB3	1:B:82:PRO:HD3	1.91	0.53
1:B:213:LYS:HD2	2:D:16:A:N6	2.24	0.53
1:B:11:GLY:HA3	1:B:102:TYR:CD2	2.43	0.53
2:C:22:G:H1'	2:C:24:A:N6	2.23	0.53
1:A:20:ARG:NH2	1:A:210:GLN:HG2	2.24	0.52
1:A:33:VAL:HG21	1:A:86:ALA:HB2	1.91	0.52
1:A:2:PRO:HA	1:A:74:THR:O	2.09	0.52
1:A:208:ARG:NH2	2:C:17:G:O6	2.43	0.51
1:B:156:THR:HG22	1:B:158:LEU:HD23	1.93	0.50
1:A:83:PHE:O	1:A:84:LEU:C	2.54	0.50
1:B:68:LEU:CD2	1:B:70:LEU:HD11	2.40	0.50
2:D:24:A:O2'	2:D:25:G:OP2	2.27	0.49
1:A:209:VAL:CG1	1:A:210:GLN:N	2.75	0.49
1:A:30:LEU:CD2	1:A:41:LEU:HD12	2.42	0.49
1:B:104:LEU:HD21	1:B:107:VAL:CG2	2.43	0.48
2:C:21:C:C2'	2:C:22:G:O5'	2.61	0.48
1:B:158:LEU:HB2	1:B:159:PRO:HD2	1.96	0.48
1:A:75:LEU:HD13	1:A:175:HIS:CD2	2.49	0.48
1:B:55:PRO:O	1:B:56:ARG:CB	2.61	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:PRO:HG2	1:B:68:LEU:HD22	1.95	0.48
1:B:52:ALA:HA	1:B:257:GLY:HA2	1.94	0.48
1:A:81:ALA:HB3	1:A:82:PRO:CD	2.43	0.48
1:B:86:ALA:O	1:B:90:ALA:N	2.41	0.48
2:C:20:U:C2'	2:C:21:C:H5'	2.45	0.47
2:D:16:A:H4'	2:D:17:G:OP1	2.14	0.47
1:A:254:THR:HA	1:A:258:MET:O	2.14	0.47
1:A:160:ASP:HB3	1:A:163:LEU:HB2	1.96	0.46
1:B:71:ARG:NH2	1:B:246:PHE:O	2.45	0.46
2:C:20:U:H2'	2:C:21:C:H5'	1.96	0.46
1:A:253:LYS:NZ	2:C:28:23G:N7	2.64	0.46
1:B:103:ARG:O	1:B:105:ALA:N	2.48	0.46
1:A:120:THR:HG22	1:A:121:SER:N	2.30	0.46
1:A:192:PHE:O	1:A:196:LEU:HB2	2.16	0.46
1:B:146:THR:O	1:B:154:ARG:HB2	2.15	0.46
1:A:20:ARG:CZ	1:A:210:GLN:HG2	2.46	0.46
1:A:208:ARG:NH2	2:C:17:G:N7	2.57	0.45
1:B:242:HIS:CE1	1:B:263:PRO:HB3	2.51	0.45
1:B:3:GLN:OE1	1:B:80:PHE:CE1	2.69	0.45
1:B:45:PRO:O	2:D:27:G:H5'	2.16	0.45
1:B:242:HIS:O	1:B:263:PRO:HG3	2.16	0.45
2:D:22:G:H1'	2:D:23:U:C5	2.51	0.45
1:B:0:SER:HB3	1:B:77:ASP:HB3	1.99	0.45
1:A:161:PRO:HD3	1:A:198:VAL:HG23	1.99	0.45
1:B:3:GLN:OE1	1:B:80:PHE:CD1	2.70	0.44
1:A:256:TYR:OH	2:C:28:23G:H2'	2.17	0.44
1:B:71:ARG:HD2	1:B:116:LEU:HB3	1.97	0.44
1:B:250:VAL:HG23	1:B:261:ALA:HB3	1.99	0.44
1:A:11:GLY:HA3	1:A:102:TYR:CD2	2.52	0.44
1:B:63:THR:HG23	1:B:64:LEU:N	2.32	0.44
1:B:119:ALA:O	1:B:120:THR:HG23	2.18	0.44
1:A:204:LEU:O	1:A:205:ARG:HG3	2.18	0.44
1:A:44:ALA:O	1:A:47:LYS:HE2	2.17	0.44
1:A:72:LEU:HD22	1:A:83:PHE:CE2	2.53	0.44
1:B:130:LYS:HA	1:B:236:GLU:HA	2.00	0.44
1:A:64:LEU:C	1:A:64:LEU:HD23	2.43	0.44
1:B:214:GLY:HA3	2:D:16:A:C2	2.52	0.44
1:A:227:LEU:HD11	1:A:234:ALA:HB1	2.01	0.43
1:B:64:LEU:HD23	1:B:65:LYS:O	2.17	0.43
2:D:22:G:H4'	2:D:23:U:OP1	2.19	0.43
1:B:133:LYS:HE2	1:B:226:ARG:HD2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:PRO:HA	1:A:15:PRO:HD3	1.94	0.42
1:A:97:LEU:HD12	1:A:102:TYR:CE1	2.55	0.42
1:B:134:ALA:HB1	1:B:242:HIS:CD2	2.55	0.41
1:A:244:LEU:HG	1:A:248:SER:HB3	2.03	0.41
1:A:111:ARG:HG2	1:A:117:ALA:O	2.19	0.41
1:B:197:GLU:HG2	1:B:198:VAL:N	2.35	0.41
1:A:120:THR:CG2	1:A:121:SER:N	2.84	0.41
1:B:140:THR:OG1	1:B:260:LEU:HB3	2.21	0.41
1:B:146:THR:CG2	1:B:163:LEU:HD11	2.51	0.41
1:B:2:PRO:HA	1:B:74:THR:O	2.21	0.41
1:B:192:PHE:CD1	1:B:196:LEU:HD11	2.56	0.40
1:A:91:ALA:HB3	1:A:92:PRO:CD	2.52	0.40
1:A:96:PRO:HA	1:A:101:SER:HA	2.03	0.40
1:A:141:PRO:HG2	1:A:209:VAL:CG2	2.45	0.40
1:A:171:LYS:HD3	1:A:248:SER:HB2	2.02	0.40
1:B:49:PHE:CD1	1:B:49:PHE:C	2.98	0.40
1:B:77:ASP:OD1	1:B:78:GLY:N	2.54	0.40
1:B:140:THR:O	1:B:141:PRO:C	2.64	0.40
1:A:204:LEU:HA	1:A:220:THR:O	2.22	0.40
1:B:177:PRO:C	1:B:179:PRO:HD3	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/268 (96%)	242 (94%)	16 (6%)	0	100	100
1	B	256/268 (96%)	244 (95%)	11 (4%)	1 (0%)	30	65
All	All	514/536 (96%)	486 (95%)	27 (5%)	1 (0%)	43	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	104	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/209 (99%)	193 (93%)	14 (7%)	14	45
1	B	207/209 (99%)	199 (96%)	8 (4%)	28	62
All	All	414/418 (99%)	392 (95%)	22 (5%)	20	54

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LYS
1	A	70	LEU
1	A	74	THR
1	A	77	ASP
1	A	79	LEU
1	A	85	ARG
1	A	97	LEU
1	A	121	SER
1	A	190	GLU
1	A	197	GLU
1	A	204	LEU
1	A	233	GLU
1	A	254	THR
1	A	262	VAL
1	B	3	GLN
1	B	37	LEU
1	B	64	LEU
1	B	95	LEU
1	B	198	VAL
1	B	204	LEU
1	B	232	LEU
1	B	262	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	114	HIS
1	A	175	HIS
1	B	42	HIS
1	B	114	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	11/29 (37%)	4 (36%)	0
2	D	13/29 (44%)	4 (30%)	1 (7%)
All	All	24/58 (41%)	8 (33%)	1 (4%)

All (8) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	18	C
2	C	22	G
2	C	23	U
2	C	24	A
2	D	17	G
2	D	22	G
2	D	23	U
2	D	24	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	D	16	A

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	23G	C	28	2	24,29,30	0.50	0	34,45,48	1.34	3 (8%)
2	23G	D	28	2	24,29,30	0.52	0	34,45,48	1.20	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	23G	C	28	2	-	0/7/35/36	0/4/4/4
2	23G	D	28	2	-	0/7/35/36	0/4/4/4

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	C	28	23G	O2'-PC-OC2	-5.64	100.86	115.76
2	D	28	23G	O2'-PC-OC2	-4.40	104.14	115.76
2	D	28	23G	O3'-PC-OC2	-2.97	107.91	115.76
2	C	28	23G	O3'-PC-OC2	-2.93	108.01	115.76
2	C	28	23G	OC1-PC-OC2	2.90	119.24	109.89
2	D	28	23G	OC1-PC-OC2	2.85	119.11	109.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	28	23G	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/268 (97%)	0.72	37 (14%) 6 4	24, 54, 90, 130	0
1	B	260/268 (97%)	0.66	29 (11%) 10 6	25, 51, 85, 116	0
2	C	12/29 (41%)	1.62	5 (41%) 0 1	56, 81, 97, 114	0
2	D	13/29 (44%)	2.07	6 (46%) 0 0	51, 68, 89, 98	0
All	All	547/594 (92%)	0.75	77 (14%) 6 4	24, 54, 90, 130	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	46	ARG	6.2
1	B	46	ARG	5.5
1	B	63	THR	5.0
1	B	39	GLN	4.6
2	D	20	U	4.4
1	B	77	ASP	4.4
1	A	20	ARG	4.3
1	A	39	GLN	4.2
1	A	213	LYS	4.0
1	A	33	VAL	3.9
1	B	213	LYS	3.8
1	B	0	SER	3.7
1	A	2	PRO	3.7
1	B	236	GLU	3.7
1	A	111	ARG	3.7
1	B	56	ARG	3.6
1	A	212	GLY	3.5
1	B	32	ARG	3.5
2	C	16	A	3.4
1	A	59	PRO	3.4
1	A	106	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	153	THR	3.3
1	B	65	LYS	3.2
1	A	181	ASN	3.2
1	B	36	GLU	3.1
1	B	106	ARG	3.1
1	B	214	GLY	3.1
2	D	19	C	3.1
1	A	36	GLU	3.0
1	B	-1	ALA	3.0
2	D	17	G	2.9
1	A	69	ARG	2.8
1	A	43	GLU	2.8
1	A	84	LEU	2.8
1	B	40	LYS	2.8
1	B	99	ASP	2.7
1	A	99	ASP	2.7
1	A	60	GLU	2.7
1	A	58	GLY	2.7
1	A	40	LYS	2.7
2	D	16	A	2.6
1	A	153	THR	2.6
1	A	32	ARG	2.6
1	A	34	SER	2.6
1	A	75	LEU	2.6
1	A	145	ALA	2.6
1	A	152	ARG	2.6
1	A	205	ARG	2.5
1	B	132	GLU	2.5
1	A	214	GLY	2.5
1	B	151	GLY	2.5
1	A	211	ALA	2.5
1	B	76	ASP	2.5
2	C	20	U	2.5
1	A	150	GLY	2.5
1	B	43	GLU	2.4
1	A	17	TYR	2.4
1	A	1	MET	2.4
1	A	146	THR	2.4
2	C	24	A	2.4
1	B	145	ALA	2.4
1	B	146	THR	2.3
1	A	95	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	148	LYS	2.3
2	D	21	C	2.3
1	B	152	ARG	2.3
1	B	147	SER	2.2
1	B	20	ARG	2.2
1	A	77	ASP	2.2
2	C	17	G	2.2
2	C	25	G	2.2
1	B	111	ARG	2.1
1	B	110	THR	2.1
1	B	155	TYR	2.1
1	A	122	TRP	2.1
2	D	23	U	2.1
1	A	110	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	23G	D	28	26/27	0.89	0.15	54,59,61,62	0
2	23G	C	28	26/27	0.91	0.13	59,69,71,72	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.