



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:06 PM UTC

PDB ID : 6C9B / pdb_00006c9b
Title : The structure of MppP soaked with the products 4HKA and 2KA
Authors : Han, L.; Silvaggi, N.R.
Deposited on : 2018-01-26
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	
1	B	376	
1	C	376	
1	D	376	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 23097 atoms, of which 10937 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PLP-Dependent L-Arginine Hydroxylase MppP.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	353	Total	C	H	N	O	P	S	0	0	0
			5435	1726	2690	487	524	1	7			
1	B	369	Total	C	H	N	O	P	S	0	4	0
			5733	1822	2838	510	555	1	7			
1	C	353	Total	C	H	N	O	P	S	0	1	0
			5436	1726	2691	487	524	1	7			
1	D	353	Total	C	H	N	O	P	S	0	4	0
			5470	1735	2708	492	527	1	7			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		

- Molecule 3 is (4S)-5-carbamimidamido-4-hydroxy-2-oxopentanoic acid (CCD ID: EGV) (formula: C₆H₁₁N₃O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	H	N	O	0	0
			23	6	10	3	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	239	Total	O	0	0
			239	239		
4	B	256	Total	O	0	0
			256	256		
4	C	245	Total	O	0	0
			245	245		
4	D	257	Total	O	0	0
			257	257		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP




- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP



- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP



- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.11Å 108.83Å 195.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.79 – 1.69 48.79 – 1.69	Depositor EDS
% Data completeness (in resolution range)	97.0 (48.79-1.69) 97.0 (48.79-1.69)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.69Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.151 , 0.165 0.152 , 0.166	Depositor DCC
R_{free} test set	2000 reflections (0.97%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	23097	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, LLP, EGV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2778	0.56	0/3782
1	B	0.38	0/2949	0.58	0/4015
1	C	0.40	0/2787	0.58	0/3794
1	D	0.39	0/2813	0.60	0/3828
All	All	0.39	0/11327	0.58	0/15419

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	2690	2690	14	0
1	B	2895	2838	2823	11	0
1	C	2745	2691	2685	1	0
1	D	2762	2708	2696	10	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	B	13	10	0	0	0
4	A	239	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	256	0	0	4	0
4	C	245	0	0	1	0
4	D	257	0	0	2	0
All	All	12160	10937	10894	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:ASP:OD2	4:A:501:HOH:O	2.13	0.66
1:B:90:SER:HB3	1:B:221:LLP:H5'2	1.81	0.62
1:C:145:GLU:OE2	4:C:501:HOH:O	2.17	0.59
1:A:99:ARG:HG2	1:A:125:ASN:OD1	2.07	0.55
1:B:169:GLU:HG2	4:B:663:HOH:O	2.08	0.54
1:D:164:ARG:HH11	1:D:164:ARG:HG3	1.76	0.50
1:D:370:ARG:O	1:D:374:GLU:HG3	2.13	0.49
1:D:363:GLU:O	1:D:367:GLN:HG2	2.14	0.47
1:D:116:ASP:OD2	4:D:501:HOH:O	2.20	0.47
1:D:30:ALA:HA	1:D:356:SER:HB3	1.96	0.47
1:A:30:ALA:HA	1:A:356:SER:HB3	1.97	0.46
1:D:190:SER:HA	1:D:218:ASP:HB3	1.98	0.46
1:A:157:THR:HA	1:A:158:PRO:C	2.41	0.45
1:A:297:GLY:O	1:A:314:PHE:HA	2.16	0.45
1:A:318:THR:HA	1:A:348:ASP:O	2.16	0.44
1:B:220:GLY:O	4:B:502:HOH:O	2.21	0.44
1:D:157:THR:HA	1:D:158:PRO:C	2.43	0.44
1:B:30:ALA:HA	1:B:356:SER:HB3	1.99	0.44
1:D:123:ARG:NH2	4:D:515:HOH:O	2.51	0.42
1:A:80:GLU:H	1:A:80:GLU:CD	2.26	0.42
1:B:169:GLU:CG	4:B:663:HOH:O	2.64	0.42
1:A:27:ASP:HB3	4:A:501:HOH:O	2.18	0.42
1:B:288:VAL:HG22	1:B:366:VAL:HG11	2.02	0.42
1:B:7:LEU:N	4:B:512:HOH:O	2.52	0.42
1:B:190:SER:HA	1:B:218:ASP:HB3	2.02	0.42
1:D:80:GLU:CD	1:D:80:GLU:H	2.28	0.42
1:A:207:GLN:NE2	4:A:504:HOH:O	2.41	0.41
1:B:66:GLN:O	1:B:70:THR:HG23	2.20	0.41
1:A:195:ASP:OD1	1:A:306:ARG:NH1	2.53	0.41
1:A:288:VAL:HG22	1:A:366:VAL:HG21	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:323:TRP:CD1	1:A:335:PRO:HD3	2.56	0.41
1:A:190:SER:HA	1:A:218:ASP:HB3	2.03	0.41
1:B:157:THR:HA	1:B:158:PRO:C	2.44	0.41
1:A:23:LEU:HB3	1:A:24:ASN:H	1.78	0.40
1:D:164:ARG:HG3	1:D:164:ARG:NH1	2.36	0.40
1:B:80:GLU:H	1:B:80:GLU:CD	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/376 (93%)	344 (98%)	6 (2%)	0	100	100
1	B	370/376 (98%)	360 (97%)	10 (3%)	0	100	100
1	C	351/376 (93%)	345 (98%)	6 (2%)	0	100	100
1	D	354/376 (94%)	349 (99%)	5 (1%)	0	100	100
All	All	1425/1504 (95%)	1398 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/314 (93%)	287 (98%)	5 (2%)	53	29
1	B	311/314 (99%)	307 (99%)	4 (1%)	61	39
1	C	293/314 (93%)	291 (99%)	2 (1%)	76	62
1	D	296/314 (94%)	293 (99%)	3 (1%)	68	52
All	All	1192/1256 (95%)	1178 (99%)	14 (1%)	63	43

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	ARG
1	A	227	ASP
1	A	317	ARG
1	A	363	GLU
1	A	364	LYS
1	B	287	VAL
1	B	296	GLU
1	B	314	PHE
1	B	348	ASP
1	C	314	PHE
1	C	317	ARG
1	D	31	ARG
1	D	164	ARG
1	D	307	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	HIS
1	A	58	GLN
1	A	75	HIS
1	A	338	GLN
1	B	39	GLN
1	B	75	HIS
1	B	125	ASN
1	B	330	HIS
1	C	58	GLN
1	C	75	HIS
1	C	125	ASN
1	D	39	GLN
1	D	75	HIS
1	D	207	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	LLP	C	221	1	23,24,25	1.09	1 (4%)	25,32,34	1.26	2 (8%)
1	LLP	A	221	1	23,24,25	1.20	2 (8%)	25,32,34	1.16	3 (12%)
1	LLP	D	221	1	23,24,25	1.13	2 (8%)	25,32,34	1.35	3 (12%)
1	LLP	B	221	1	23,24,25	0.94	1 (4%)	25,32,34	1.85	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	C	221	1	-	4/16/17/19	0/1/1/1
1	LLP	A	221	1	-	4/16/17/19	0/1/1/1
1	LLP	D	221	1	-	3/16/17/19	0/1/1/1
1	LLP	B	221	1	-	6/16/17/19	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	LLP	O-C	3.98	1.35	1.20
1	D	221	LLP	O-C	3.95	1.35	1.20
1	C	221	LLP	O-C	3.75	1.34	1.20
1	B	221	LLP	OP4-C5'	-3.11	1.33	1.44
1	A	221	LLP	C3-C2	2.48	1.43	1.41
1	D	221	LLP	OP4-C5'	-2.43	1.35	1.44

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	221	LLP	C5'-C5-C6	-5.69	110.08	119.36
1	B	221	LLP	OP3-P-OP4	-3.73	96.94	106.67
1	B	221	LLP	C3-C4-C5	-3.51	115.46	118.28
1	D	221	LLP	C5'-C5-C6	-3.42	113.78	119.36
1	A	221	LLP	C5'-C5-C6	-3.10	114.31	119.36
1	D	221	LLP	C3-C4-C5	-3.06	115.82	118.28
1	C	221	LLP	OP2-P-OP4	-2.95	98.98	106.67
1	C	221	LLP	C5'-C5-C6	-2.54	115.22	119.36
1	B	221	LLP	OP4-C5'-C5	2.50	114.04	109.36
1	A	221	LLP	OP4-C5'-C5	2.44	113.92	109.36
1	A	221	LLP	OP2-P-OP4	-2.15	101.07	106.67
1	D	221	LLP	OP4-P-OP1	-2.03	100.96	106.44

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	221	LLP	C4-C5-C5'-OP4
1	A	221	LLP	C6-C5-C5'-OP4
1	B	221	LLP	C4-C5-C5'-OP4
1	B	221	LLP	C6-C5-C5'-OP4
1	B	221	LLP	C5'-OP4-P-OP2
1	B	221	LLP	C5'-OP4-P-OP3
1	C	221	LLP	C4-C5-C5'-OP4
1	C	221	LLP	C6-C5-C5'-OP4
1	D	221	LLP	C4-C5-C5'-OP4
1	D	221	LLP	C6-C5-C5'-OP4
1	C	221	LLP	C4-C4'-NZ-CE
1	B	221	LLP	CG-CD-CE-NZ
1	D	221	LLP	C4-C4'-NZ-CE
1	C	221	LLP	CA-CB-CG-CD
1	A	221	LLP	CA-CB-CG-CD
1	B	221	LLP	C5'-OP4-P-OP1
1	A	221	LLP	C4-C4'-NZ-CE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	221	LLP	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EGV	B	401	-	12,12,12	2.12	5 (41%)	13,15,15	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EGV	B	401	-	-	0/13/13/13	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	EGV	C03-N01	5.17	1.43	1.33
3	B	401	EGV	O11-C10	-3.12	1.22	1.30
3	B	401	EGV	C07-C09	2.56	1.55	1.51
3	B	401	EGV	C03-N04	2.13	1.42	1.34
3	B	401	EGV	O08-C06	-2.04	1.37	1.43

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/376 (93%)	-0.27	8 (2%) 61 68	18, 31, 62, 81	0
1	B	368/376 (97%)	-0.20	14 (3%) 44 52	13, 29, 64, 99	2 (0%)
1	C	352/376 (93%)	-0.33	9 (2%) 57 65	15, 30, 56, 89	0
1	D	352/376 (93%)	-0.28	9 (2%) 57 65	15, 29, 67, 88	2 (0%)
All	All	1424/1504 (94%)	-0.27	40 (2%) 55 62	13, 30, 63, 99	4 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	7	LEU	5.7
1	B	375	THR	4.7
1	A	23	LEU	4.4
1	B	115	PHE	4.2
1	B	294	GLY	4.2
1	C	375	THR	4.0
1	B	293	ALA	4.0
1	D	293	ALA	3.9
1	D	375	THR	3.6
1	B	314	PHE	3.5
1	A	375	THR	3.5
1	A	315	ALA	3.4
1	A	115	PHE	3.4
1	D	332	PHE	3.4
1	D	294	GLY	3.2
1	C	115	PHE	3.2
1	B	11	LEU	3.0
1	A	323	TRP	2.9
1	C	316	GLY	2.9
1	C	23	LEU	2.9
1	A	314	PHE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	293	ALA	2.8
1	D	23	LEU	2.8
1	D	373	LEU	2.8
1	C	315	ALA	2.7
1	B	373	LEU	2.7
1	D	252	LEU	2.6
1	D	366	VAL	2.6
1	C	314	PHE	2.5
1	C	330	HIS	2.4
1	B	19	LEU	2.4
1	B	372	VAL	2.4
1	C	293	ALA	2.2
1	D	295	VAL	2.2
1	B	296	GLU	2.2
1	A	117	ASN	2.1
1	B	295	VAL	2.1
1	C	323	TRP	2.1
1	B	16	TYR	2.1
1	B	21	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	221	24/25	0.98	0.06	18,29,37,43	0
1	LLP	B	221	24/25	0.98	0.05	17,26,41,45	0
1	LLP	C	221	24/25	0.98	0.06	19,27,41,44	0
1	LLP	D	221	24/25	0.98	0.05	16,25,38,40	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	C	401	1/1	0.90	0.15	33,33,33,33	0
2	CL	A	401	1/1	0.94	0.10	36,36,36,36	0
3	EGV	B	401	13/13	0.94	0.11	19,49,62,66	0
2	CL	D	401	1/1	0.99	0.06	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.