



wwPDB EM Validation Summary Report ⓘ

Mar 24, 2026 – 02:52 AM UTC

PDB ID : 8C9M / pdb_00008c9m
EMDB ID : EMD-16511
Title : HERV-K Gag immature lattice
Authors : Krebs, A.-S.; Liu, H.-F.; Zhou, Y.; Rey, J.S.; Levintov, L.; Perilla, J.R.; Bartesaghi, A.; Zhang, P.
Deposited on : 2023-01-23
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	666	 27% 7% 66%
1	B	666	 27% 7% 66%
1	C	666	 27% 8% 66%
1	D	666	 27% 7% 66%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	666	 27% 7% 66%
1	F	666	 27% 7% 66%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 21480 atoms, of which 10752 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gag protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	228	Total	C	H	N	O	S	0	0
			3580	1127	1792	315	334	12		
1	B	228	Total	C	H	N	O	S	0	0
			3580	1127	1792	315	334	12		
1	C	228	Total	C	H	N	O	S	0	0
			3580	1127	1792	315	334	12		
1	D	228	Total	C	H	N	O	S	0	0
			3580	1127	1792	315	334	12		
1	E	228	Total	C	H	N	O	S	0	0
			3580	1127	1792	315	334	12		
1	F	228	Total	C	H	N	O	S	0	0
			3580	1127	1792	315	334	12		

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-191	ILE	LEU	conflict	UNP Q96897
A	-185	ASN	LYS	conflict	UNP Q96897
A	-174	GLY	SER	conflict	UNP Q96897
A	-57	PRO	SER	conflict	UNP Q96897
A	38	HIS	TYR	conflict	UNP Q96897
A	120	CYS	VAL	conflict	UNP Q96897
A	133	THR	ALA	conflict	UNP Q96897
A	160	GLN	GLU	conflict	UNP Q96897
A	191	PRO	SER	conflict	UNP Q96897
A	209	ALA	VAL	conflict	UNP Q96897
A	255	ASN	SER	conflict	UNP Q96897
A	291	GLN	LEU	conflict	UNP Q96897
A	356	PRO	LEU	conflict	UNP Q96897
B	-191	ILE	LEU	conflict	UNP Q96897
B	-185	ASN	LYS	conflict	UNP Q96897
B	-174	GLY	SER	conflict	UNP Q96897
B	-57	PRO	SER	conflict	UNP Q96897
B	38	HIS	TYR	conflict	UNP Q96897

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	120	CYS	VAL	conflict	UNP Q96897
B	133	THR	ALA	conflict	UNP Q96897
B	160	GLN	GLU	conflict	UNP Q96897
B	191	PRO	SER	conflict	UNP Q96897
B	209	ALA	VAL	conflict	UNP Q96897
B	255	ASN	SER	conflict	UNP Q96897
B	291	GLN	LEU	conflict	UNP Q96897
B	356	PRO	LEU	conflict	UNP Q96897
C	-191	ILE	LEU	conflict	UNP Q96897
C	-185	ASN	LYS	conflict	UNP Q96897
C	-174	GLY	SER	conflict	UNP Q96897
C	-57	PRO	SER	conflict	UNP Q96897
C	38	HIS	TYR	conflict	UNP Q96897
C	120	CYS	VAL	conflict	UNP Q96897
C	133	THR	ALA	conflict	UNP Q96897
C	160	GLN	GLU	conflict	UNP Q96897
C	191	PRO	SER	conflict	UNP Q96897
C	209	ALA	VAL	conflict	UNP Q96897
C	255	ASN	SER	conflict	UNP Q96897
C	291	GLN	LEU	conflict	UNP Q96897
C	356	PRO	LEU	conflict	UNP Q96897
D	-191	ILE	LEU	conflict	UNP Q96897
D	-185	ASN	LYS	conflict	UNP Q96897
D	-174	GLY	SER	conflict	UNP Q96897
D	-57	PRO	SER	conflict	UNP Q96897
D	38	HIS	TYR	conflict	UNP Q96897
D	120	CYS	VAL	conflict	UNP Q96897
D	133	THR	ALA	conflict	UNP Q96897
D	160	GLN	GLU	conflict	UNP Q96897
D	191	PRO	SER	conflict	UNP Q96897
D	209	ALA	VAL	conflict	UNP Q96897
D	255	ASN	SER	conflict	UNP Q96897
D	291	GLN	LEU	conflict	UNP Q96897
D	356	PRO	LEU	conflict	UNP Q96897
E	-191	ILE	LEU	conflict	UNP Q96897
E	-185	ASN	LYS	conflict	UNP Q96897
E	-174	GLY	SER	conflict	UNP Q96897
E	-57	PRO	SER	conflict	UNP Q96897
E	38	HIS	TYR	conflict	UNP Q96897
E	120	CYS	VAL	conflict	UNP Q96897
E	133	THR	ALA	conflict	UNP Q96897
E	160	GLN	GLU	conflict	UNP Q96897

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	191	PRO	SER	conflict	UNP Q96897
E	209	ALA	VAL	conflict	UNP Q96897
E	255	ASN	SER	conflict	UNP Q96897
E	291	GLN	LEU	conflict	UNP Q96897
E	356	PRO	LEU	conflict	UNP Q96897
F	-191	ILE	LEU	conflict	UNP Q96897
F	-185	ASN	LYS	conflict	UNP Q96897
F	-174	GLY	SER	conflict	UNP Q96897
F	-57	PRO	SER	conflict	UNP Q96897
F	38	HIS	TYR	conflict	UNP Q96897
F	120	CYS	VAL	conflict	UNP Q96897
F	133	THR	ALA	conflict	UNP Q96897
F	160	GLN	GLU	conflict	UNP Q96897
F	191	PRO	SER	conflict	UNP Q96897
F	209	ALA	VAL	conflict	UNP Q96897
F	255	ASN	SER	conflict	UNP Q96897
F	291	GLN	LEU	conflict	UNP Q96897
F	356	PRO	LEU	conflict	UNP Q96897


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gag protein



- Molecule 1: Gag protein





GLY
LYS
ALA
PHE
PRO
ILE
GLN
PRO
PHE
VAL
GLN
GLN
PHE
GLY
GLN
GLN
PRO
PRO
LEU
SER
GLN
VAL
PHE
GLN
GLY
ILE
SER
GLN
LEU
PRO
GLN
TYR
ASN
ASN
CYS
PRO
PRO
PRO
GLN
ALA
ALA
VAL
GLN
GLN

● Molecule 1: Gag protein

Chain F: 27% 7% 66%

MET
GLY
THR
ILE
THR
LEU
LYS
SER
LYS
ILE
LYS
VAL
SER
GLY
TVR
LYS
SER
ALA
TYR
ILE
SER
PHE
PRO
ILE
LEU
LYS
ILE
LEU
TRP
GLY
GLN
ASP
ARG
GLY
ILE
VAL
LYS
VAL
SER
THR
LYS
ASN
PHE
LEU
PHE
LEU
ILE
LYS
THR
LEU
GLU
SER
PHE
GLY
GLN
SER
CYS
PRO
TRP
PHE
PRO
PRO
GLU
GLN
ILE
GLY
THR
ASP
CYS
LEU
ASP
GLY
TRP
ARG

LYS
ARG
ILE
GLY
LYS
GLU
LEU
LYS
THR
GLU
GLN
ALA
GLY
CYS
ARG
LYS
GLY
ASN
ILE
ILE
PRO
PRO
LEU
THR
VAL
TRP
ASN
SER
LEU
THR
TRP
ALA
ILE
ILE
LYS
LYS
ALA
VAL
SER
ALA
GLY
LEU
GLU
PHE
LEU
THR
GLU
SER
VAL
SER
SER
ASP
ALA
PRO
PRO
GLY
GLY
SER
GLY
CYS
SER
CYS
ILE
GLN
GLY
THR
ASP
CYS
GLU
ASN
GLY
THR
ARG

LYS
LYS
SER
GLN
LYS
GLU
THR
GLY
GLU
GLY
HIS
CYS
GLY
TYR
VAL
ALA
GLU
PRO
LYS
GLN
VAL
MET
ALA
GLN
SER
THR
GLN
ASN
GLN
VAL
LEU
GLN
GLY
VAL
PHE
LEU
GLU
GLY
LYS
GLY
PRO
VAL
GLY
GLY
PRO
SER
SER
GLU
SER
LYS
PRO
GLY
ARG
GLY

THR
SER
PRO
LEU
PRO
ALA
GLY
GLN
VAL
ARG
VAL
THR
LEU
ASN
THR
GLN
PRO
ALA
PRO
GLN
LYS
GLN
VAL
LYS
GLY
GLY
THR
THR
GLN
PRO
ALA
VAL
ILE
VAL
ASP
ALA
TYR
LYS
GLN
TYR
TRP
TRP
LYS
GLY
GLU
PRO
GLU
THR
THR
PHE
PRO
PRO
GLU
SER
GLN
TYR
TYR
MET
PRO
PRO
GLY
GLY
PRO
MET
GLY
GLY
PRO
ALA
GLN
PRO
GLN
GLY
GLY
ARG

ALA
PRO
TYR
PRO
GLN
PRO
PRO
THR
THR
ARG
ARG
VAL
LEU
ASN
PRO
THR
ALA
PRO
GLN
SER
SER
ARG
GLN
GLY
SER
GLY
GLU
LEU
HIS
GLU
ILE
ILE
ASP
LYS
SER
ARG
LYS
GLY
GLU
GLY
THR
ALA
GLN
THR
VAL
THR
LEU
GLU
PRO
MET
PRO
PRO
GLY
GLY
GLY
GLY
GLY
ALA
GLN
GLU
GLY
GLY
PRO
GLY
ARG

PRO
THR
VAL
E1
A2
R3
Y4
I9
G18
V19
G23
S26
P27
D34
S35
I36
H40
R41
L42
I43
D46
K52
S56
F63
W67
Q72
E73
Q74
P83
D90
S100
T101
E114
Q115
A118
T119
C120
W124
D129
T133

C134
T139
V140
F151
V152
A153
Q156
D157
N182
Q186
K193
V196
V202
E206
Y206
V207
K208
D211
K219
T223
GLY
VAL
VAL
LEU
GLY
GLY
GLN
VAL
ARG
THR
PHE
GLY
GLY
LYS
CYS
TYR
ASN
CYS
GLY
GLN
LEU
PRO
SER
GLY
ASN
GLN
GLN
ILE
GLY
HIS
LEU
LYS
LYS
ASN
CYS

PRO
VAL
LEU
ASN
LYS
GLN
ASN
ILE
THR
THR
GLN
VAL
ALA
THR
THR
THR
GLY
GLY
ARG
GLU
GLN
PRO
PRO
ASP
LEU
CYS
PRO
GLN
ARG
CYS
LYS
LYS
GLY
LYS
HIS
TRP
ALA
SER
GLN
TYR
ASN
CYS
SER
CYS
PRO
PHE
PRO
GLN
ALA
GLY
VAL
GLN

GLN
THR
GLY
ALA
PHE
PRO
ILE
GLN
PRO
PHE
VAL
PRO
GLN
GLY
PHE
GLN
GLN
GLN
PRO
PRO
SER
SER
VAL
PHE
GLN
ILE
SER
GLN
LEU
PRO
GLN

4 Experimental information

Property	Value	Source
EM reconstruction method	SUBTOMOGRAM AVERAGING	Depositor
Imposed symmetry	POINT, C6	Depositor
Number of subtomograms used	188111	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	127.5	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.198	Depositor
Minimum map value	-0.932	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.031	Depositor
Recommended contour level	0.0736	Depositor
Map size (Å)	384.768, 384.768, 384.768	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.503, 1.503, 1.503	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.65	36/1825 (2.0%)	1.10	17/2467 (0.7%)
1	B	1.65	35/1825 (1.9%)	1.11	17/2467 (0.7%)
1	C	1.68	39/1825 (2.1%)	1.11	19/2467 (0.8%)
1	D	1.65	35/1825 (1.9%)	1.10	16/2467 (0.6%)
1	E	1.66	35/1825 (1.9%)	1.11	16/2467 (0.6%)
1	F	1.65	36/1825 (2.0%)	1.10	17/2467 (0.7%)
All	All	1.66	216/10950 (2.0%)	1.10	102/14802 (0.7%)

The worst 5 of 216 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	67	TRP	CA-C	-10.85	1.39	1.52
1	B	67	TRP	CA-C	-10.82	1.39	1.52
1	D	67	TRP	CA-C	-10.82	1.39	1.52
1	A	67	TRP	CA-C	-10.81	1.39	1.52
1	C	67	TRP	CA-C	-10.81	1.39	1.52

The worst 5 of 102 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	23	GLY	CA-C-N	8.02	128.46	119.32
1	E	23	GLY	C-N-CA	8.02	128.46	119.32
1	D	23	GLY	CA-C-N	8.00	128.44	119.32
1	D	23	GLY	C-N-CA	8.00	128.44	119.32
1	C	23	GLY	CA-C-N	8.00	128.44	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1788	1792	1794	4	0
1	B	1788	1792	1794	4	0
1	C	1788	1792	1794	4	0
1	D	1788	1792	1794	4	0
1	E	1788	1792	1794	4	0
1	F	1788	1792	1794	4	0
All	All	10728	10752	10764	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:100:SER:OG	1:E:101:THR:N	2.43	0.51
1:F:100:SER:OG	1:F:101:THR:N	2.43	0.51
1:A:100:SER:OG	1:A:101:THR:N	2.43	0.51
1:B:100:SER:OG	1:B:101:THR:N	2.43	0.50
1:F:133:THR:O	1:F:134:CYS:C	2.56	0.49

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	226/666 (34%)	218 (96%)	6 (3%)	2 (1%)	14 47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	226/666 (34%)	218 (96%)	6 (3%)	2 (1%)	14	47
1	C	226/666 (34%)	218 (96%)	6 (3%)	2 (1%)	14	47
1	D	226/666 (34%)	218 (96%)	6 (3%)	2 (1%)	14	47
1	E	226/666 (34%)	218 (96%)	6 (3%)	2 (1%)	14	47
1	F	226/666 (34%)	218 (96%)	6 (3%)	2 (1%)	14	47
All	All	1356/3996 (34%)	1308 (96%)	36 (3%)	12 (1%)	16	47

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	TYR
1	B	4	TYR
1	C	4	TYR
1	D	4	TYR
1	E	4	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/570 (34%)	192 (100%)	0	100	100
1	B	192/570 (34%)	192 (100%)	0	100	100
1	C	192/570 (34%)	192 (100%)	0	100	100
1	D	192/570 (34%)	192 (100%)	0	100	100
1	E	192/570 (34%)	192 (100%)	0	100	100
1	F	192/570 (34%)	192 (100%)	0	100	100
All	All	1152/3420 (34%)	1152 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-16511. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 110




Y Index: 141




Z Index: 147


6.3.2 Raw map



X Index: 110



Y Index: 136




Z Index: 127


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0736. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 579 nm³; this corresponds to an approximate mass of 523 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.38	4.06	3.40

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-16511 and PDB model 8C9M. Per-residue inclusion information can be found in section [3](#) on page [7](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0736 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0736).

9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 77% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0736) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7660	0.4140
A	0.7710	0.4170
B	0.7760	0.4160
C	0.7760	0.4100
D	0.7720	0.4090
E	0.7730	0.4150
F	0.7790	0.4150

