



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:22 AM UTC

PDB ID : 2CAR / pdb_00002car
Title : Crystal Structure Of Human Inosine Triphosphatase
Authors : Stenmark, P.; Kursula, P.; Arrowsmith, C.; Berglund, H.; Edwards, A.; Ehn, M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Hallberg, B.M.; Hogbom, M.; Holmberg Schiavone, L.; Kotenyova, T.; Nilsson-Ehle, P.; Nyman, T.; Ogg, D.; Persson, C.; Sagemark, J.; Schuler, H.; Sundstrom, M.; Thorsell, A.G.; van den Berg, S.; Weigelt, J.; Nordlund, P.
Deposited on : 2005-12-22
Resolution : 1.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1625 (1.12-1.08)
Clashscore	190562	1656 (1.12-1.08)
Ramachandran outliers	187476	1616 (1.12-1.08)
Sidechain outliers	187428	1612 (1.12-1.08)
RSRZ outliers	180081	1625 (1.12-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	 74% 21% . .
1	B	196	 78% 17% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INOSINE TRIPHOSPHATE PYROPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	20	0
			1646	1053	275	306	12			
1	B	194	Total	C	N	O	S	0	15	0
			1611	1032	269	301	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	319	Total	O	0	0
			319	319		
2	B	329	Total	O	0	0
			329	329		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INOSINE TRIPHOSPHATE PYROPHOSPHATASE



• Molecule 1: INOSINE TRIPHOSPHATE PYROPHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.23Å 105.00Å 50.14Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	40.00 – 1.09 40.00 – 1.09	Depositor EDS
% Data completeness (in resolution range)	90.7 (40.00-1.09) 91.8 (40.00-1.09)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 1.09Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.097 , 0.153 0.105 , 0.160	Depositor DCC
R_{free} test set	2636 reflections (2.13%)	wwPDB-VP
Wilson B-factor (Å ²)	7.7	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.488 for h,-k,-l	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	3905	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	0/1721	1.92	39/2323 (1.7%)
1	B	0.92	0/1679	1.76	18/2269 (0.8%)
All	All	0.93	0/3400	1.84	57/4592 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	ASP	CA-CB-CG	-24.37	88.23	112.60
1	A	125	PRO	CA-C-N	13.27	146.88	121.54
1	A	125	PRO	C-N-CA	13.27	146.88	121.54
1	B	41	ASP	CA-CB-CG	12.28	124.88	112.60
1	A	124	ASP	CB-CA-C	11.54	132.90	110.17
1	A	144	ARG	NE-CZ-NH2	10.56	128.71	119.20
1	A	124	ASP	CA-C-N	10.52	132.99	119.84
1	A	124	ASP	C-N-CA	10.52	132.99	119.84
1	A	126	SER	CA-C-N	10.16	137.69	121.83
1	A	126	SER	C-N-CA	10.16	137.69	121.83
1	B	178	ARG	CD-NE-CZ	9.33	137.47	124.40
1	B	16	ASN	OD1-CG-ND2	9.12	131.72	122.60
1	A	125	PRO	CA-C-O	-9.06	104.10	120.60
1	B	187	GLU	CB-CG-CD	9.00	127.90	112.60
1	A	124	ASP	CB-CG-OD2	8.28	137.45	118.40
1	A	144	ARG	NE-CZ-NH1	-7.89	113.61	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	GLU	CB-CG-CD	7.79	125.84	112.60
1	B	58	GLN	OE1-CD-NE2	7.37	129.97	122.60
1	B	135	ARG	NE-CZ-NH1	-7.26	114.24	121.50
1	B	98	GLU	CB-CG-CD	7.08	124.63	112.60
1	B	122	THR	CA-CB-CG2	6.99	122.38	110.50
1	A	77	PHE	CA-CB-CG	6.86	120.66	113.80
1	A	0	SER	CA-C-O	6.68	132.16	120.80
1	A	2	ALA	CA-C-O	6.33	127.47	120.82
1	A	39	LYS	CA-C-O	6.29	127.78	120.98
1	A	125	PRO	O-C-N	-6.26	114.19	122.64
1	B	135	ARG	NH1-CZ-NH2	6.22	127.38	119.30
1	A	147	GLN	CB-CG-CD	6.21	123.15	112.60
1	A	64	VAL	N-CA-C	6.13	116.91	110.72
1	A	124	ASP	CB-CG-OD1	-6.11	104.34	118.40
1	A	159	GLY	O-C-N	6.08	129.13	122.56
1	B	55	GLN	CG-CD-NE2	5.98	125.37	116.40
1	A	16[A]	ASN	CA-CB-CG	-5.97	106.63	112.60
1	A	16[B]	ASN	CA-CB-CG	-5.97	106.63	112.60
1	A	158	ASP	CA-C-O	5.92	127.66	121.15
1	A	33	CYS	CA-C-O	-5.79	114.86	121.58
1	A	126	SER	CA-CB-OG	-5.73	99.65	111.10
1	B	67	PRO	N-CA-CB	-5.72	98.22	103.31
1	B	62	ARG	NE-CZ-NH2	5.60	124.24	119.20
1	A	107	PHE	O-C-N	-5.60	116.76	123.31
1	A	58	GLN	CG-CD-NE2	-5.58	108.04	116.40
1	B	66	GLY	O-C-N	5.50	127.27	121.77
1	B	193	ALA	O-C-N	-5.47	115.74	122.25
1	A	64	VAL	O-C-N	-5.46	116.21	121.83
1	B	124	ASP	CA-CB-CG	-5.46	107.14	112.60
1	B	162	GLN	CA-CB-CG	5.42	124.94	114.10
1	A	108	GLU	CB-CG-CD	5.34	121.68	112.60
1	B	126	SER	N-CA-C	-5.31	106.80	113.28
1	A	127	GLN	N-CA-CB	5.26	117.69	110.01
1	B	51	GLU	O-C-N	5.26	127.69	122.12
1	A	2	ALA	N-CA-CB	-5.26	102.39	110.01
1	A	0	SER	O-C-N	-5.25	114.59	123.00
1	A	28	GLY	O-C-N	-5.11	117.24	122.60
1	A	15	GLY	CA-C-O	-5.07	114.43	119.04
1	A	124	ASP	N-CA-CB	-5.07	101.35	110.37
1	A	55[A]	GLN	O-C-N	5.05	127.48	122.12
1	A	55[B]	GLN	O-C-N	5.05	127.48	122.12

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	124	ASP	Peptide
1	A	125	PRO	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1646	0	1662	32	0
1	B	1611	0	1622	22	0
2	A	319	0	0	7	0
2	B	329	0	0	11	0
All	All	3905	0	3284	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71[B]:GLU:OE1	1:A:178[B]:ARG:NH1	1.76	1.15
1:B:51:GLU:O	1:B:55:GLN:HG3	1.82	0.80
1:A:52:ILE:HG23	2:A:2128:HOH:O	1.86	0.75
1:B:22[B]:GLU:HB2	2:B:2073:HOH:O	1.86	0.74
1:B:22[A]:GLU:HB2	2:B:2073:HOH:O	1.87	0.74
1:B:40:ILE:HG23	2:B:2172:HOH:O	1.98	0.64
1:A:124:ASP:HB3	1:A:125:PRO:O	1.98	0.63
1:A:124:ASP:CG	1:A:127:GLN:H	2.10	0.59
1:A:127:GLN:OE1	1:A:128:PRO:HD2	2.03	0.57
1:A:45:TYR:HD2	2:A:2128:HOH:O	1.87	0.57
1:B:56:LYS:HE3	2:B:2126:HOH:O	2.04	0.57
1:B:3:ALA:HB3	2:B:2009:HOH:O	2.05	0.56
1:A:71[B]:GLU:OE1	1:A:178[B]:ARG:CZ	2.53	0.53
1:A:143:PRO:HA	1:A:153[B]:PRO:O	2.09	0.53
1:B:38:GLN:HG3	2:B:2044:HOH:O	2.09	0.52
1:B:16:ASN:OD1	1:B:18:LYS:HB3	2.10	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:PRO:HG3	2:A:2216:HOH:O	2.08	0.52
1:A:162[A]:GLN:HG3	2:A:2261:HOH:O	2.11	0.49
1:B:93:GLU:HG2	2:B:2097:HOH:O	2.12	0.49
1:B:1:MET:HB3	1:B:188:TYR:OH	2.13	0.48
1:A:1[B]:MET:CE	1:A:130[B]:ARG:HG3	2.44	0.48
1:A:97:PRO:HB3	1:A:154[B]:CYS:SG	2.54	0.47
1:A:160:TYR:CD2	1:A:167[A]:MET:HE3	2.50	0.47
1:A:98[B]:GLU:OE2	1:A:147:GLN:NE2	2.46	0.47
1:A:125:PRO:HB2	1:A:126:SER:OG	2.15	0.46
1:B:63:GLN:NE2	2:B:2172:HOH:O	2.48	0.46
1:B:62:ARG:NH1	2:B:2167:HOH:O	2.49	0.46
1:B:4:SER:N	2:B:2009:HOH:O	2.48	0.45
1:B:130[B]:ARG:HG2	1:B:132:PHE:CZ	2.52	0.45
1:A:153[A]:PRO:O	1:A:163:THR:HG21	2.17	0.45
1:A:124:ASP:OD1	1:A:127:GLN:N	2.45	0.44
1:B:59:GLU:OE2	1:B:62:ARG:NH2	2.49	0.44
1:A:149:PHE:O	1:A:152:ASP:HB2	2.17	0.44
1:A:67:PRO:HA	1:A:121:SER:O	2.18	0.43
1:A:1[A]:MET:HE2	1:A:122:THR:HG23	2.00	0.43
1:A:178[A]:ARG:NH2	2:A:2290:HOH:O	2.45	0.43
1:B:130[B]:ARG:HD3	1:B:132:PHE:CZ	2.54	0.43
1:A:4:SER:N	2:A:2001:HOH:O	2.50	0.43
1:A:124:ASP:OD2	1:A:127:GLN:O	2.36	0.43
1:A:163:THR:O	1:A:167[A]:MET:HG3	2.19	0.43
1:B:137:SER:OG	1:B:158[A]:ASP:OD2	2.30	0.43
1:B:8:LYS:HD3	2:B:2124:HOH:O	2.18	0.42
1:B:1:MET:HE2	1:B:122:THR:HG23	2.01	0.42
1:A:125:PRO:HB2	1:A:126:SER:CB	2.49	0.42
1:B:168:PRO:HG2	1:B:171:GLU:OE2	2.20	0.42
1:A:50:ASP:OD1	1:A:133[B]:ARG:NH2	2.50	0.42
1:B:42:LEU:HD11	1:B:60:ALA:HB2	2.00	0.42
1:A:66:GLY:O	1:A:68[B]:VAL:HG23	2.20	0.41
1:A:5:LEU:HA	1:A:8:LYS:HG3	2.03	0.41
1:A:1[A]:MET:SD	1:A:122:THR:HG23	2.60	0.41
1:A:124:ASP:OD1	1:A:127:GLN:HB3	2.21	0.41
1:A:139:ARG:NH2	2:A:2234:HOH:O	2.53	0.40
1:A:163:THR:O	1:A:167[B]:MET:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/196 (109%)	208 (97%)	3 (1%)	3 (1%)	9	0
1	B	207/196 (106%)	206 (100%)	0	1 (0%)	24	5
All	All	421/392 (107%)	414 (98%)	3 (1%)	4 (1%)	12	1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	ASP
1	A	126	SER
1	A	125	PRO
1	B	96	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/161 (112%)	172 (95%)	9 (5%)	22	1
1	B	175/161 (109%)	171 (98%)	4 (2%)	44	9
All	All	356/322 (111%)	343 (96%)	13 (4%)	35	3

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	MET
1	A	1[B]	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	97	PRO
1	A	98[A]	GLU
1	A	98[B]	GLU
1	A	124	ASP
1	A	139	ARG
1	A	162[A]	GLN
1	A	162[B]	GLN
1	B	1	MET
1	B	16	ASN
1	B	126	SER
1	B	164	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	186	GLN
1	B	63	GLN
1	B	162	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/196 (100%)	-1.16	0 100 100	3, 10, 23, 39	20 (10%)
1	B	194/196 (98%)	-1.18	0 100 100	5, 9, 24, 39	15 (7%)
All	All	390/392 (99%)	-1.17	0 100 100	3, 9, 24, 39	35 (8%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.