



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:19 AM UTC

PDB ID : 5CA5 / pdb_00005ca5
Title : Structure of the *C. elegans* NONO-1 homodimer
Authors : Knott, G.J.; Bond, C.S.
Deposited on : 2015-06-29
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	 81% 13% ..
1	B	259	 80% 14% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4176 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NONO-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	248	Total 2010	C 1264	N 366	O 370	S 10	0	1	0
1	B	248	Total 2002	C 1259	N 363	O 370	S 10	0	0	0

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	74	Total 74	O 74	0	0
3	B	74	Total 74	O 74	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NONO-1



- Molecule 1: NONO-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.32Å 63.12Å 83.84Å 90.00° 101.03° 90.00°	Depositor
Resolution (Å)	19.67 – 2.40 19.67 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.5 (19.67-2.40) 97.3 (19.67-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.41Å)	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.243 , 0.301 0.256 , 0.318	Depositor DCC
R_{free} test set	1206 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtrriage
Anisotropy	0.379	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4176	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.88	0/2054	1.47	12/2762 (0.4%)
1	B	0.89	0/2043	1.44	8/2748 (0.3%)
All	All	0.88	0/4097	1.45	20/5510 (0.4%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	ASP	CA-CB-CG	6.31	118.91	112.60
1	A	134	GLU	CA-C-N	6.13	128.50	120.28
1	A	134	GLU	C-N-CA	6.13	128.50	120.28
1	A	302	ASN	CA-CB-CG	6.04	118.64	112.60
1	B	245	ALA	CA-C-N	5.88	128.09	120.44
1	B	245	ALA	C-N-CA	5.88	128.09	120.44
1	A	238	ARG	CA-C-N	5.77	127.99	120.26
1	A	238	ARG	C-N-CA	5.77	127.99	120.26
1	B	238	ARG	CA-C-N	5.68	127.87	120.26
1	B	238	ARG	C-N-CA	5.68	127.87	120.26
1	B	286	PRO	CA-C-N	5.29	125.85	119.98
1	B	286	PRO	C-N-CA	5.29	125.85	119.98
1	B	289	SER	CA-C-N	5.16	127.45	120.44
1	B	289	SER	C-N-CA	5.16	127.45	120.44
1	A	331	ARG	CA-C-N	5.15	127.44	120.44
1	A	331	ARG	C-N-CA	5.15	127.44	120.44
1	A	201	SER	CA-C-N	5.05	127.36	120.54
1	A	201	SER	C-N-CA	5.05	127.36	120.54
1	A	289	SER	CA-C-N	5.03	127.28	120.44
1	A	289	SER	C-N-CA	5.03	127.28	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2010	0	2005	18	0
1	B	2002	0	1992	23	0
2	A	8	0	12	1	0
2	B	8	0	12	1	0
3	A	74	0	0	1	0
3	B	74	0	0	2	0
All	All	4176	0	4021	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:LYS:HD3	1:B:227:ARG:HH11	1.62	0.64
1:B:225:LYS:HD3	1:B:227:ARG:NH1	2.13	0.63
1:A:197:SER:HB3	1:A:200:VAL:HG23	1.81	0.61
1:B:197:SER:HB3	1:B:200:VAL:HG23	1.84	0.60
1:A:319:GLU:HG2	1:B:330:LEU:HD22	1.85	0.59
1:A:299:PRO:HD2	1:B:253:VAL:O	2.01	0.59
1:B:270:PRO:HA	3:B:534:HOH:O	2.03	0.58
1:A:225:LYS:HD3	1:A:227:ARG:NH2	2.20	0.56
1:A:253:VAL:O	1:B:299:PRO:HD2	2.07	0.55
1:A:330:LEU:HD22	1:B:319:GLU:HG2	1.89	0.53
1:A:326:LEU:HD22	1:B:326:LEU:HD22	1.92	0.52
1:B:327:ASP:O	1:B:331:ARG:HB2	2.10	0.52
1:B:327:ASP:HB3	1:B:331:ARG:HH11	1.75	0.51
1:A:233:ILE:HG12	1:A:268:LEU:HD11	1.93	0.51
1:A:225:LYS:HD3	1:A:227:ARG:HH21	1.75	0.50
1:B:115:ARG:HD3	3:B:548:HOH:O	2.12	0.50
1:A:113:THR:HG22	1:A:115:ARG:H	1.79	0.48
1:B:194:LYS:HD3	1:B:195:GLU:HG2	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:VAL:HG22	1:B:313:LYS:HG3	1.97	0.47
1:B:149:GLY:H	2:B:401:EDO:H22	1.80	0.46
1:B:269:GLU:H	1:B:269:GLU:HG2	1.54	0.46
1:B:253:VAL:HG13	1:B:262:PRO:HB3	1.99	0.45
1:A:243:ASN:HB3	2:A:402:EDO:H22	1.98	0.45
1:B:157:LEU:HD13	1:B:163:ALA:HA	1.99	0.44
1:B:222:VAL:HG12	1:B:228:PRO:HA	1.99	0.44
1:A:253:VAL:HG13	1:A:262:PRO:HB3	1.99	0.43
1:A:211:HIS:HB2	3:A:554:HOH:O	2.19	0.43
1:A:222:VAL:HG12	1:A:228:PRO:HA	2.00	0.43
1:A:273:GLU:HG2	1:B:221:ILE:HG12	2.00	0.42
1:A:337:LEU:HG	1:A:341:MET:HE3	2.02	0.42
1:A:201:SER:HB3	1:B:282:ILE:HD11	2.02	0.41
1:B:288:LEU:O	1:B:291:GLU:HG2	2.20	0.41
1:B:337:LEU:HG	1:B:341:MET:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/259 (95%)	245 (99%)	2 (1%)	0	100	100
1	B	246/259 (95%)	245 (100%)	1 (0%)	0	100	100
All	All	493/518 (95%)	490 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/221 (97%)	203 (95%)	11 (5%)	21	37
1	B	213/221 (96%)	203 (95%)	10 (5%)	23	41
All	All	427/442 (97%)	406 (95%)	21 (5%)	22	39

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	GLU
1	A	113	THR
1	A	115	ARG
1	A	144	GLU
1	A	145	CYS
1	A	175	ILE
1	A	180	VAL
1	A	186	VAL
1	A	290	LYS
1	A	325	GLN
1	A	342	GLU
1	B	115	ARG
1	B	144	GLU
1	B	157	LEU
1	B	175	ILE
1	B	186	VAL
1	B	268	LEU
1	B	269	GLU
1	B	290	LYS
1	B	325	GLN
1	B	342	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	302	ASN
1	A	325	GLN
1	B	302	ASN
1	B	325	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	401	-	3,3,3	0.57	0	2,2,2	0.30	0
2	EDO	B	402	-	3,3,3	0.51	0	2,2,2	0.34	0
2	EDO	A	401	-	3,3,3	0.50	0	2,2,2	0.36	0
2	EDO	A	402	-	3,3,3	0.60	0	2,2,2	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	401	-	-	1/1/1/1	-
2	EDO	B	402	-	-	0/1/1/1	-
2	EDO	A	401	-	-	1/1/1/1	-
2	EDO	A	402	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	EDO	O1-C1-C2-O2
2	A	401	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	EDO	1	0
2	A	402	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/259 (95%)	0.41	3 (1%) 76 73	26, 46, 69, 84	1 (0%)
1	B	248/259 (95%)	0.35	1 (0%) 88 86	29, 44, 64, 76	0
All	All	496/518 (95%)	0.38	4 (0%) 82 80	26, 44, 68, 84	1 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	ARG	3.3
1	B	108	PRO	2.4
1	A	291	GLU	2.2
1	A	302	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	B	402	4/4	0.64	0.17	54,55,55,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	B	401	4/4	0.80	0.14	57,57,57,58	0
2	EDO	A	402	4/4	0.83	0.15	32,36,38,39	0
2	EDO	A	401	4/4	0.90	0.08	46,46,48,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.