



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:53 PM UTC

PDB ID : 1CB2 / pdb_00001cb2
Title : CELLOBIOHYDROLASE II, CATALYTIC DOMAIN, MUTANT Y169F
Authors : Kleywegt, G.J.; Szardenings, M.; Jones, T.A.
Deposited on : 1995-11-25
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	 4% 91% 8% ..
1	B	365	 4% 91% 8% ..

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CELLOBIOHYDROLASE II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	363	Total 2746	C 1743	N 463	O 530	S 10	0	0	0
1	B	363	Total 2746	C 1743	N 463	O 530	S 10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	169	PHE	TYR	engineered mutation	UNP P07987
B	169	PHE	TYR	engineered mutation	UNP P07987

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 14	C 8	N 1	O 5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is alpha-D-mannopyranose (CCD ID: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	196	Total	O	0	0
			196	196		
4	B	196	Total	O	0	0
			196	196		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CELLOBIOHYDROLASE II



- Molecule 1: CELLOBIOHYDROLASE II



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.10Å 75.80Å 92.90Å 90.00° 103.20° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00 8.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	91.7 (8.00-2.00) 90.9 (8.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 1.86Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.210 , 0.232 0.213 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6094	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.12 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3794e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2822	0.93	12/3868 (0.3%)
1	B	0.42	0/2822	0.93	12/3868 (0.3%)
All	All	0.42	0/5644	0.93	24/7736 (0.3%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	157	LYS	N-CA-C	-7.09	103.89	112.54
1	A	157	LYS	N-CA-C	-7.08	103.91	112.54
1	A	218	ILE	N-CA-C	6.21	117.05	107.99
1	B	218	ILE	N-CA-C	6.20	117.05	107.99
1	B	391	PHE	N-CA-C	-5.96	99.94	109.96
1	A	391	PHE	N-CA-C	-5.95	99.97	109.96
1	B	229	ASN	N-CA-C	5.75	119.27	112.72
1	A	400	CYS	N-CA-C	5.74	118.16	110.35
1	A	229	ASN	N-CA-C	5.72	119.25	112.72
1	B	400	CYS	N-CA-C	5.71	118.12	110.35
1	B	368	CYS	N-CA-C	5.64	117.87	109.25
1	A	368	CYS	N-CA-C	5.62	117.85	109.25
1	B	219	GLU	N-CA-C	5.58	122.14	109.81
1	A	219	GLU	N-CA-C	5.57	122.11	109.81
1	A	101	ASN	N-CA-C	5.55	117.89	110.35
1	B	101	ASN	N-CA-C	5.54	117.88	110.35
1	B	255	LEU	CA-C-N	5.21	124.83	119.05
1	B	255	LEU	C-N-CA	5.21	124.83	119.05
1	A	255	LEU	CA-C-N	5.19	124.82	119.05
1	A	255	LEU	C-N-CA	5.19	124.82	119.05
1	A	159	GLY	N-CA-C	5.13	122.18	115.36
1	B	159	GLY	N-CA-C	5.09	122.14	115.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	446	PHE	N-CA-C	-5.04	107.26	113.41
1	B	446	PHE	N-CA-C	-5.03	107.27	113.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2746	0	2606	14	0
1	B	2746	0	2606	14	0
2	A	28	0	26	0	0
2	B	28	0	26	0	0
3	A	77	0	70	2	0
3	B	77	0	70	2	0
4	A	196	0	0	0	0
4	B	196	0	0	0	0
All	All	6094	0	5404	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:ARG:HD3	1:A:389:ASP:OD2	1.98	0.63
1:B:297:ARG:HD3	1:B:389:ASP:OD2	1.98	0.62
1:A:361:GLN:HG3	1:A:370:VAL:HG11	1.84	0.59
1:B:361:GLN:HG3	1:B:370:VAL:HG11	1.84	0.58
1:B:116:LEU:HB2	1:B:121:ALA:HB2	1.86	0.58
1:A:116:LEU:HB2	1:A:121:ALA:HB2	1.86	0.57
1:B:281:GLN:HG2	1:B:340:HIS:CE1	2.43	0.54
1:A:281:GLN:HG2	1:A:340:HIS:CE1	2.43	0.53
1:B:157:LYS:O	1:B:157:LYS:HD3	2.10	0.52
1:A:157:LYS:O	1:A:157:LYS:HD3	2.10	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:TYR:CE2	1:A:297:ARG:HG2	2.47	0.50
1:A:97:THR:O	1:A:162:TYR:HA	2.12	0.49
1:B:261:TYR:CE2	1:B:297:ARG:HG2	2.47	0.49
1:B:97:THR:O	1:B:162:TYR:HA	2.12	0.49
1:A:374:GLY:HA2	1:A:438:LEU:HA	1.95	0.49
1:A:298:GLY:HA3	1:A:346:PHE:O	2.14	0.48
1:B:374:GLY:HA2	1:B:438:LEU:HA	1.95	0.48
1:B:298:GLY:HA3	1:B:346:PHE:O	2.14	0.47
1:A:297:ARG:O	1:A:297:ARG:HG3	2.18	0.44
1:B:277:ASP:O	1:B:281:GLN:HG3	2.19	0.43
1:B:87:THR:HG21	3:B:503:MAN:H5	2.01	0.43
1:A:277:ASP:O	1:A:281:GLN:HG3	2.19	0.42
1:B:297:ARG:O	1:B:297:ARG:HG3	2.18	0.42
1:B:277:ASP:HB3	1:B:278:PRO:HD3	2.01	0.42
1:A:277:ASP:HB3	1:A:278:PRO:HD3	2.01	0.42
1:A:87:THR:HG21	3:A:503:MAN:H5	2.01	0.41
1:A:161:ASN:HB3	3:A:504:MAN:O2	2.21	0.41
1:B:161:ASN:HB3	3:B:504:MAN:O2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/365 (99%)	345 (96%)	16 (4%)	0	100	100
1	B	361/365 (99%)	345 (96%)	16 (4%)	0	100	100
All	All	722/730 (99%)	690 (96%)	32 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/285 (100%)	282 (99%)	2 (1%)	76	82
1	B	284/285 (100%)	282 (99%)	2 (1%)	76	82
All	All	568/570 (100%)	564 (99%)	4 (1%)	76	82

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	213	ARG
1	A	297	ARG
1	B	213	ARG
1	B	297	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	158	ASN
1	A	165	GLN
1	A	252	GLN
1	A	339	ASN
1	B	158	ASN
1	B	165	GLN
1	B	252	GLN
1	B	339	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	B	506	1	11,11,12	0.46	0	15,15,17	0.54	0
3	MAN	A	507	1	11,11,12	0.43	0	15,15,17	0.56	0
3	MAN	B	508	1	11,11,12	0.41	0	15,15,17	0.51	0
3	MAN	A	503	1	11,11,12	0.43	0	15,15,17	0.70	1 (6%)
2	NAG	B	502	1	14,14,15	0.39	0	17,19,21	0.65	0
3	MAN	B	504	1	11,11,12	0.42	0	15,15,17	0.61	0
3	MAN	B	503	1	11,11,12	0.43	0	15,15,17	0.70	1 (6%)
3	MAN	B	507	1	11,11,12	0.42	0	15,15,17	0.57	0
3	MAN	B	509	1	11,11,12	0.47	0	15,15,17	0.59	0
3	MAN	A	509	1	11,11,12	0.47	0	15,15,17	0.59	0
3	MAN	A	504	1	11,11,12	0.42	0	15,15,17	0.61	0
3	MAN	A	506	1	11,11,12	0.46	0	15,15,17	0.55	0
2	NAG	B	501	1	14,14,15	0.42	0	17,19,21	0.78	0
2	NAG	A	501	1	14,14,15	0.43	0	17,19,21	0.79	0
3	MAN	A	508	1	11,11,12	0.41	0	15,15,17	0.52	0
3	MAN	A	505	1	11,11,12	0.42	0	15,15,17	0.69	1 (6%)
2	NAG	A	502	1	14,14,15	0.40	0	17,19,21	0.64	0
3	MAN	B	505	1	11,11,12	0.43	0	15,15,17	0.69	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	B	506	1	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	A	507	1	-	0/2/19/22	0/1/1/1
3	MAN	B	508	1	-	1/2/19/22	0/1/1/1
3	MAN	A	503	1	-	1/2/19/22	0/1/1/1
2	NAG	B	502	1	-	0/6/23/26	0/1/1/1
3	MAN	B	504	1	-	2/2/19/22	0/1/1/1
3	MAN	B	503	1	-	1/2/19/22	0/1/1/1
3	MAN	B	507	1	-	0/2/19/22	0/1/1/1
3	MAN	B	509	1	-	0/2/19/22	0/1/1/1
3	MAN	A	509	1	-	0/2/19/22	0/1/1/1
3	MAN	A	504	1	-	2/2/19/22	0/1/1/1
3	MAN	A	506	1	-	0/2/19/22	0/1/1/1
2	NAG	B	501	1	-	0/6/23/26	0/1/1/1
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
3	MAN	A	508	1	-	1/2/19/22	0/1/1/1
3	MAN	A	505	1	-	0/2/19/22	0/1/1/1
2	NAG	A	502	1	-	0/6/23/26	0/1/1/1
3	MAN	B	505	1	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	MAN	C1-O5-C5	2.37	115.36	112.19
3	B	503	MAN	C1-O5-C5	2.36	115.34	112.19
3	B	505	MAN	C1-O5-C5	2.15	115.07	112.19
3	A	505	MAN	C1-O5-C5	2.15	115.07	112.19

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	504	MAN	O5-C5-C6-O6
3	B	504	MAN	O5-C5-C6-O6
3	A	504	MAN	C4-C5-C6-O6
3	B	504	MAN	C4-C5-C6-O6
3	B	508	MAN	C4-C5-C6-O6
3	A	508	MAN	C4-C5-C6-O6
3	B	503	MAN	O5-C5-C6-O6
3	A	503	MAN	O5-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	503	MAN	1	0
3	B	504	MAN	1	0
3	B	503	MAN	1	0
3	A	504	MAN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/365 (99%)	0.04	15 (4%) 41 40	2, 10, 34, 100	0
1	B	363/365 (99%)	-0.02	13 (3%) 46 45	2, 10, 34, 100	0
All	All	726/730 (99%)	0.01	28 (3%) 43 42	2, 10, 35, 100	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	158	ASN	5.0
1	B	85	THR	4.6
1	A	159	GLY	4.5
1	B	204	GLN	4.2
1	A	407	SER	4.1
1	A	160	GLY	3.9
1	B	158	ASN	3.6
1	A	409	PRO	3.3
1	B	409	PRO	3.1
1	A	118	GLY	3.0
1	A	363	GLN	2.9
1	B	159	GLY	2.9
1	A	85	THR	2.8
1	A	319	GLN	2.7
1	A	121	ALA	2.7
1	A	408	ALA	2.7
1	B	407	SER	2.7
1	B	426	GLN	2.6
1	B	140	ASP	2.4
1	B	89	SER	2.4
1	A	157	LYS	2.4
1	A	294	ARG	2.4
1	A	117	THR	2.4
1	A	426	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	408	ALA	2.3
1	B	208	GLU	2.3
1	B	157	LYS	2.1
1	B	118	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	A	503	11/12	0.26	0.26	97,97,97,97	0
3	MAN	B	509	11/12	0.47	0.17	64,64,64,64	0
3	MAN	B	503	11/12	0.48	0.23	97,97,97,97	0
3	MAN	B	504	11/12	0.54	0.23	77,77,77,77	0
3	MAN	A	508	11/12	0.55	0.17	75,75,75,75	0
3	MAN	A	509	11/12	0.69	0.17	64,64,64,64	0
3	MAN	A	504	11/12	0.69	0.23	77,77,77,77	0
3	MAN	B	508	11/12	0.72	0.16	75,75,75,75	0
2	NAG	A	502	14/15	0.73	0.11	20,20,20,20	0
2	NAG	B	502	14/15	0.77	0.11	20,20,20,20	0
3	MAN	A	505	11/12	0.77	0.10	40,40,40,40	0
3	MAN	B	507	11/12	0.78	0.09	37,37,37,37	0
3	MAN	A	507	11/12	0.78	0.10	37,37,37,37	0
3	MAN	B	505	11/12	0.78	0.14	40,40,40,40	0
3	MAN	B	506	11/12	0.81	0.09	20,20,20,20	0
2	NAG	B	501	14/15	0.82	0.08	23,23,23,23	0
3	MAN	A	506	11/12	0.85	0.08	20,20,20,20	0
2	NAG	A	501	14/15	0.86	0.08	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.