



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:15 AM UTC

PDB ID : 3CBL / pdb_00003cbl
Title : Crystal structure of human feline sarcoma viral oncogene homologue (v-FES) in complex with staurosporine and a consensus peptide
Authors : Filippakopoulos, P.; Salah, E.; Cooper, C.; Picaud, S.S.; Elkins, J.M.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.M.; Weigelt, J.; Bountra, C.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2008-02-22
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	37% 86%8%6%
2	B	6	17% 100%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Fes/Fps.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2824	1815	479	515	15	0	13	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	446	SER	-	expression tag	UNP P07332
A	447	MET	-	expression tag	UNP P07332

- Molecule 2 is a protein called Synthetic peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	47	31	5	11	0	0	0

- Molecule 3 is STAUROSPORINE (CCD ID: STU) (formula: $C_{28}H_{26}N_4O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			35	28	4	3		
3	A	1	Total	C	N	O	0	0
			35	28	4	3		
3	A	1	Total	C	N	O	0	0
			35	28	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	320	Total	O	0	0
			320	320		
4	B	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Fes/Fps



- Molecule 2: Synthetic peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	35.68Å 77.18Å 149.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.58 – 1.75 38.58 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.1 (38.58-1.75) 99.1 (38.58-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 1.75Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.180 , 0.227 0.189 , 0.229	Depositor DCC
R_{free} test set	2141 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	21.1	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3304	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, STU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/2935	0.87	2/3977 (0.1%)
2	B	0.77	0/45	0.75	0/59
All	All	0.83	0/2980	0.86	2/4036 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	804	ARG	CA-C-N	-5.51	114.91	120.31
1	A	804	ARG	C-N-CA	-5.51	114.91	120.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2824	0	2778	23	0
2	B	47	0	45	0	0
3	A	105	0	78	6	0
4	A	320	0	0	2	0
4	B	8	0	0	1	0
All	All	3304	0	2901	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:902:STU:H261	3:A:902:STU:H16	1.63	0.81
1:A:627:THR:HA	1:A:632:ILE:HD12	1.62	0.80
1:A:704[A]:MET:HE2	4:A:321:HOH:O	1.87	0.74
3:A:903:STU:H261	3:A:903:STU:H16	1.74	0.68
1:A:704[A]:MET:CE	4:A:321:HOH:O	2.44	0.63
1:A:480:PHE:CG	1:A:545:LEU:HD22	2.34	0.63
3:A:901:STU:H261	3:A:901:STU:H16	1.80	0.63
1:A:625:VAL:HG12	1:A:627:THR:HG23	1.81	0.62
1:A:721:GLN:HG2	4:B:169:HOH:O	2.03	0.59
1:A:493:LEU:HD23	1:A:515:LEU:HD11	1.84	0.59
1:A:629:LYS:O	1:A:632:ILE:HD13	2.03	0.58
1:A:464:ALA:HB3	1:A:632:ILE:HD11	1.88	0.55
1:A:469:GLU:OE1	1:A:609:ARG:NH2	2.40	0.54
3:A:902:STU:H16	3:A:902:STU:C26	2.38	0.53
1:A:493:LEU:CD2	1:A:515:LEU:HD11	2.40	0.51
1:A:512:LEU:HD23	1:A:520:PHE:C	2.36	0.51
1:A:504:PHE:HB3	1:A:515:LEU:HD22	1.92	0.50
1:A:480:PHE:CD1	1:A:545:LEU:HD22	2.47	0.49
1:A:579:ARG:HG2	1:A:586:LEU:HD23	1.95	0.49
1:A:656:ARG:HD3	1:A:658:LYS:HZ3	1.79	0.47
1:A:571[B]:ASN:ND2	1:A:721:GLN:HE22	2.13	0.46
1:A:613:GLN:NE2	1:A:707[B]:GLU:OE2	2.38	0.46
1:A:520:PHE:CZ	1:A:529:HIS:CB	2.99	0.45
3:A:903:STU:H16	3:A:903:STU:C26	2.45	0.42
1:A:697:LEU:C	1:A:697:LEU:HD23	2.44	0.41
3:A:902:STU:C26	3:A:902:STU:C16	2.98	0.41
1:A:610:ILE:CD1	1:A:707[B]:GLU:HG3	2.51	0.41
1:A:596:LEU:HD22	1:A:600:LEU:HD23	2.02	0.41
1:A:464:ALA:CB	1:A:632:ILE:HD11	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/377 (95%)	349 (98%)	7 (2%)	1 (0%)	36	21
2	B	4/6 (67%)	4 (100%)	0	0	100	100
All	All	361/383 (94%)	353 (98%)	7 (2%)	1 (0%)	36	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	701	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/327 (93%)	303 (99%)	2 (1%)	76	67
2	B	5/5 (100%)	5 (100%)	0	100	100
All	All	310/332 (93%)	308 (99%)	2 (1%)	76	71

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	594	GLU
1	A	821	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	721	GLN
1	A	821	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	STU	A	902	-	39,42,42	1.73	10 (25%)	50,68,68	1.96	12 (24%)
3	STU	A	903	-	39,42,42	2.47	12 (30%)	50,68,68	1.86	12 (24%)
3	STU	A	901	-	39,42,42	1.98	11 (28%)	50,68,68	1.47	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	STU	A	902	-	-	0/4/42/42	-
3	STU	A	903	-	-	1/4/42/42	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	STU	A	901	-	-	0/4/42/42	-

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	903	STU	C25-N3	6.69	1.51	1.45
3	A	903	STU	C20-N3	-5.78	1.32	1.40
3	A	903	STU	C17-N2	-5.28	1.34	1.40
3	A	903	STU	C8-N1	-5.14	1.31	1.35
3	A	901	STU	C20-N3	-4.55	1.34	1.40
3	A	902	STU	C20-N3	-4.55	1.34	1.40
3	A	903	STU	C18-N2	-4.39	1.35	1.40
3	A	901	STU	C25-N3	-4.35	1.42	1.45
3	A	901	STU	C18-N2	-4.19	1.35	1.40
3	A	902	STU	C17-N2	-3.84	1.36	1.40
3	A	901	STU	C12-C11	-3.80	1.37	1.46
3	A	901	STU	C17-N2	-3.77	1.36	1.40
3	A	903	STU	C26-C21	3.68	1.55	1.51
3	A	903	STU	C24-C25	3.65	1.58	1.51
3	A	902	STU	C12-C11	-3.38	1.38	1.46
3	A	903	STU	C23-N4	3.31	1.51	1.47
3	A	902	STU	C18-N2	-3.31	1.36	1.40
3	A	903	STU	C19-N3	-3.15	1.34	1.39
3	A	903	STU	C12-C11	-2.90	1.40	1.46
3	A	902	STU	C5-C6	-2.69	1.40	1.46
3	A	903	STU	C7-C8	-2.67	1.40	1.49
3	A	901	STU	C8-N1	-2.66	1.33	1.35
3	A	902	STU	C25-N3	2.65	1.47	1.45
3	A	901	STU	C7-C8	-2.51	1.41	1.49
3	A	901	STU	C24-C25	2.46	1.56	1.51
3	A	903	STU	C5-C6	-2.45	1.41	1.46
3	A	902	STU	C8-N1	-2.39	1.33	1.35
3	A	901	STU	C26-C21	2.36	1.54	1.51
3	A	902	STU	C9-N1	2.35	1.48	1.45
3	A	902	STU	C7-C8	-2.23	1.42	1.49
3	A	901	STU	C5-C6	-2.12	1.41	1.46
3	A	902	STU	C22-C23	2.04	1.55	1.53
3	A	901	STU	C19-N3	-2.02	1.36	1.39

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	STU	C7-C8-N1	5.72	111.84	106.38
3	A	902	STU	C9-N1-C8	-5.70	108.67	113.86
3	A	903	STU	O5-C8-N1	-5.61	119.11	125.35
3	A	903	STU	C7-C8-N1	4.93	111.08	106.38
3	A	902	STU	O5-C8-N1	-4.90	119.90	125.35
3	A	902	STU	C9-C10-C7	-4.84	108.11	109.88
3	A	901	STU	C28-N4-C23	4.43	119.69	114.39
3	A	903	STU	C9-C10-C7	-4.32	108.30	109.88
3	A	901	STU	C7-C8-N1	3.62	109.83	106.38
3	A	903	STU	C6-C7-C10	-3.58	118.27	120.76
3	A	903	STU	C16-C17-C12	-3.34	117.69	121.59
3	A	901	STU	C16-C17-C12	-3.30	117.73	121.59
3	A	902	STU	C16-C17-C12	-3.21	117.84	121.59
3	A	903	STU	C11-C10-C7	2.89	123.15	121.25
3	A	902	STU	C9-C10-C11	2.81	130.87	128.73
3	A	903	STU	C9-N1-C8	-2.63	111.46	113.86
3	A	903	STU	C16-C17-N2	2.62	134.15	130.22
3	A	903	STU	C6-C7-C8	2.61	134.21	130.53
3	A	901	STU	C9-C10-C7	-2.56	108.94	109.88
3	A	902	STU	C16-C17-N2	2.53	134.02	130.22
3	A	902	STU	C18-C19-N3	2.31	131.38	129.42
3	A	903	STU	C15-C14-C13	-2.18	117.55	120.24
3	A	901	STU	C20-N3-C25	-2.15	121.96	127.45
3	A	902	STU	C6-C7-C10	-2.14	119.27	120.76
3	A	903	STU	C18-N2-C17	2.14	109.95	107.37
3	A	903	STU	C27-O6-C22	-2.14	110.84	114.46
3	A	902	STU	C27-O6-C22	2.13	118.06	114.46
3	A	902	STU	C6-C7-C8	2.07	133.46	130.53
3	A	901	STU	C16-C17-N2	2.05	133.29	130.22
3	A	902	STU	C19-C18-N2	-2.04	129.65	131.38
3	A	901	STU	C6-C7-C8	2.03	133.40	130.53

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	903	STU	C24-C23-N4-C28


There are no ring outliers.

3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	902	STU	3	0
3	A	903	STU	2	0
3	A	901	STU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand STU A 902




Bond lengths



Bond angles




Torsions




Rings


Ligand STU A 903




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/377 (94%)	1.89	139 (39%)  	21, 37, 52, 63	13 (3%)
2	B	5/6 (83%)	1.50	1 (20%)  	31, 32, 38, 51	0
All	All	361/383 (94%)	1.88	140 (38%)  	21, 37, 52, 63	13 (3%)

All (140) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	545	LEU	6.1
1	A	546	HIS	5.9
1	A	544	VAL	5.8
1	A	478	GLY	5.6
1	A	549	VAL	5.2
1	A	515	LEU	5.1
1	A	734	TYR	5.0
1	A	537	LEU	4.9
1	A	571[A]	ASN	4.9
1	A	567	ILE	4.9
1	A	473	LEU	4.9
1	A	576	PHE	4.9
1	A	586	LEU	4.8
1	A	511	ASN	4.7
1	A	475	VAL	4.7
1	A	489	GLN	4.6
1	A	512	LEU	4.5
1	A	580	LEU	4.5
1	A	531	LEU	4.5
1	A	477	SER	4.4
1	A	470	VAL	4.3
1	A	471	ALA	4.3
1	A	520	PHE	4.2
1	A	562	VAL	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	587	VAL	4.1
1	A	582	ALA	4.1
1	A	577	SER	4.1
1	A	575	VAL	4.1
1	A	589	VAL	4.1
1	A	506	ILE	4.1
1	A	732	LEU	4.0
1	A	480	PHE	4.0
1	A	543	VAL	4.0
1	A	588	ALA	4.0
1	A	456	HIS	3.9
1	A	563	LEU	3.9
1	A	474	LEU	3.9
1	A	530	LEU	3.9
1	A	821	HIS	3.9
1	A	448	ILE	3.8
1	A	523	ILE	3.8
1	A	716	SER	3.8
1	A	504	PHE	3.7
1	A	548	ALA	3.7
1	A	490	GLU	3.6
1	A	579	ARG	3.5
1	A	568	GLY	3.5
1	A	564	GLY	3.5
1	A	569	ARG	3.4
1	A	593	ARG	3.4
1	A	459	LEU	3.4
1	A	513	TYR	3.4
1	A	536	PRO	3.4
1	A	578	GLY	3.3
1	A	597	PRO	3.3
1	A	585	THR	3.3
1	A	501	PRO	3.3
1	A	454	PRO	3.3
1	A	595	THR	3.2
1	A	495	VAL	3.2
1	A	533	THR	3.2
1	A	560	ASP	3.2
1	A	584	ASN	3.1
1	A	535	GLN	3.1
1	A	561	LEU	3.1
1	A	565	GLU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	717	GLY	3.1
1	A	600	LEU	3.1
1	A	592	CYS	3.1
1	A	468	ALA	3.0
1	A	479	ASP	2.9
1	A	534	GLN	2.9
1	A	547	ARG	2.9
1	A	465	ILE	2.9
1	A	528	ASP	2.9
1	A	570	GLY	2.8
1	A	460	TRP	2.8
1	A	654	ARG	2.8
1	A	583	ASP	2.8
1	A	596	LEU	2.8
1	A	505	ILE	2.8
1	A	653	ALA	2.8
1	A	472	GLU	2.7
1	A	554	TRP	2.7
1	A	815	GLN	2.7
1	A	632	ILE	2.7
1	A	733[A]	ASN	2.7
1	A	625	VAL	2.7
1	A	532	SER	2.7
1	A	503	HIS	2.7
1	A	527	ILE	2.7
1	A	446	SER	2.6
1	A	455	LEU	2.6
1	A	556	LEU	2.6
1	A	492	VAL	2.6
1	A	629	LYS	2.6
1	A	493	LEU	2.6
1	A	451[A]	VAL	2.6
2	B	1	ILE	2.6
1	A	602	ALA	2.6
1	A	573	GLY	2.5
1	A	818[A]	ARG	2.5
1	A	521	PRO	2.5
1	A	652	GLY	2.5
1	A	566	GLN	2.5
1	A	481	LEU	2.4
1	A	581	ARG	2.4
1	A	614	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	491	TYR	2.4
1	A	609	ARG	2.4
1	A	447	MET	2.4
1	A	476	HIS	2.4
1	A	529	HIS	2.4
1	A	631	PRO	2.3
1	A	467	ARG	2.3
1	A	449	PRO	2.3
1	A	714	ALA	2.3
1	A	635	VAL	2.3
1	A	791	ARG	2.3
1	A	710	ASP	2.2
1	A	767	GLN	2.2
1	A	610	ILE	2.2
1	A	623[A]	ILE	2.2
1	A	650	THR	2.2
1	A	572	PHE	2.2
1	A	598	PRO	2.2
1	A	482	VAL	2.1
1	A	594	GLU	2.1
1	A	525	LEU	2.1
1	A	599	ASP	2.1
1	A	604	PHE	2.1
1	A	620	VAL	2.1
1	A	691	VAL	2.1
1	A	658	LYS	2.1
1	A	656	ARG	2.1
1	A	461	TYR	2.1
1	A	634	ILE	2.1
1	A	713	TYR	2.1
1	A	638	LEU	2.0
1	A	788	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	STU	A	903	35/35	0.91	0.14	30,35,43,47	0
3	STU	A	902	35/35	0.94	0.12	24,30,34,38	0
3	STU	A	901	35/35	0.97	0.10	18,22,27,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around STU A 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around STU A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around STU A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.