



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:52 AM UTC

PDB ID : 5CB1 / pdb_00005cb1
Title : Apo enzyme of human Polymerase lambda
Authors : Liu, M.S.; Tsai, M.D.
Deposited on : 2015-06-30
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	 80% 16% . .
1	B	326	 59% 15% . 25%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	0	0
			2567	1612	471	472	12			
1	B	245	Total	C	N	O	S	0	0	0
			1927	1205	356	356	10			

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA polymerase lambda

Chain A: 



• Molecule 1: DNA polymerase lambda

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	205.28Å 205.28Å 111.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.97 – 3.30 24.97 – 3.30	Depositor EDS
% Data completeness (in resolution range)	94.3 (24.97-3.30) 93.3 (24.97-3.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.82 (at 3.30Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.230 , 0.264 0.232 , 0.263	Depositor DCC
R_{free} test set	1808 reflections (9.88%)	wwPDB-VP
Wilson B-factor (Å ²)	110.0	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4494	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	1/2620 (0.0%)	0.83	10/3536 (0.3%)
1	B	0.28	0/1967	0.78	4/2657 (0.2%)
All	All	0.48	1/4587 (0.0%)	0.81	14/6193 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	462	VAL	CA-CB	-8.03	1.46	1.55

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	462	VAL	N-CA-C	-9.91	93.67	107.75
1	A	326	ASP	CB-CA-C	-9.69	105.40	116.54
1	A	463	SER	N-CA-C	-9.45	97.96	110.24
1	A	464	GLN	N-CA-C	-8.88	94.76	109.24
1	A	321	HIS	N-CA-CB	-6.70	100.78	110.36
1	A	461	LEU	N-CA-C	6.62	119.35	110.35
1	B	332	VAL	CA-C-N	-6.56	112.93	119.56
1	B	332	VAL	C-N-CA	-6.56	112.93	119.56
1	A	485	ARG	NE-CZ-NH1	-5.95	115.56	121.50
1	B	434	HIS	CA-C-N	5.67	126.93	119.84
1	B	434	HIS	C-N-CA	5.67	126.93	119.84
1	A	434	HIS	CA-C-N	5.50	125.59	119.32
1	A	434	HIS	C-N-CA	5.50	125.59	119.32
1	A	326	ASP	N-CA-C	5.34	116.42	108.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2567	0	2568	72	2
1	B	1927	0	1910	28	0
All	All	4494	0	4478	99	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:LEU:HD22	1:A:325:LEU:N	1.55	1.17
1:A:325:LEU:HD23	1:A:325:LEU:O	1.43	1.17
1:A:321:HIS:HA	1:A:323:ARG:HG2	1.26	1.16
1:A:326:ASP:OD2	1:A:353:TYR:OH	1.65	1.12
1:A:322:LEU:O	1:A:323:ARG:HG3	1.66	0.95
1:A:325:LEU:HD22	1:A:325:LEU:H	1.29	0.93
1:A:321:HIS:CA	1:A:323:ARG:HG2	2.01	0.91
1:A:325:LEU:N	1:A:325:LEU:CD2	2.30	0.88
1:A:391:GLU:OE1	1:A:487:ARG:NH2	2.09	0.84
1:A:321:HIS:HA	1:A:323:ARG:CG	2.08	0.80
1:A:322:LEU:O	1:A:323:ARG:CG	2.30	0.79
1:A:325:LEU:HD12	1:A:354:GLN:HE22	1.47	0.78
1:A:325:LEU:O	1:A:325:LEU:CD2	2.30	0.77
1:A:321:HIS:HA	1:A:322:LEU:C	2.10	0.76
1:A:342:TRP:HE1	1:A:488:ARG:HH21	1.38	0.72
1:A:321:HIS:ND1	1:A:321:HIS:N	2.34	0.72
1:A:342:TRP:NE1	1:A:488:ARG:HH21	1.89	0.70
1:A:462:VAL:O	1:A:463:SER:C	2.32	0.69
1:A:336:GLU:O	1:A:340:ASN:ND2	2.26	0.69
1:A:322:LEU:C	1:A:323:ARG:CG	2.67	0.68
1:B:511:HIS:HA	1:B:514:ARG:HD3	1.76	0.68
1:A:464:GLN:N	1:A:464:GLN:OE1	2.29	0.66
1:B:444:PHE:HE2	1:B:464:GLN:HE22	1.42	0.66
1:A:351:MET:O	1:A:355:GLN:HG3	1.96	0.65
1:A:342:TRP:CZ2	1:A:488:ARG:NE	2.65	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:ASP:OD1	1:B:452:ARG:NH1	2.30	0.64
1:A:342:TRP:CZ2	1:A:488:ARG:CZ	2.81	0.64
1:A:441:ARG:NH2	1:B:464:GLN:O	2.26	0.64
1:A:462:VAL:HG23	1:A:474:LEU:O	1.97	0.63
1:B:431:LEU:HD23	1:B:503:LEU:HD12	1.80	0.63
1:A:462:VAL:CG2	1:A:474:LEU:O	2.47	0.62
1:A:342:TRP:CE2	1:A:488:ARG:NE	2.68	0.62
1:A:391:GLU:CD	1:A:487:ARG:HH21	2.09	0.59
1:A:565:LEU:HD12	1:A:566:PRO:HD2	1.85	0.58
1:A:320:GLY:C	1:A:323:ARG:HD2	2.29	0.58
1:A:325:LEU:CD2	1:A:325:LEU:C	2.74	0.57
1:A:325:LEU:HD23	1:A:325:LEU:C	2.14	0.56
1:B:336:GLU:O	1:B:340:ASN:ND2	2.35	0.55
1:A:322:LEU:C	1:A:323:ARG:HG2	2.33	0.54
1:A:462:VAL:C	1:A:463:SER:O	2.46	0.54
1:A:337:LEU:HD11	1:A:360:LEU:HD11	1.90	0.53
1:A:395:ILE:HG12	1:A:479:LEU:HD11	1.91	0.52
1:B:449:ASP:O	1:B:453:GLN:HG2	2.09	0.52
1:B:525:MET:HE3	1:B:563:LEU:HD23	1.89	0.52
1:B:568:ARG:O	1:B:573:ARG:NH1	2.44	0.51
1:B:387:MET:HE1	1:B:392:ALA:HB2	1.93	0.51
1:A:462:VAL:O	1:A:464:GLN:OE1	2.30	0.50
1:A:462:VAL:O	1:A:463:SER:O	2.30	0.50
1:B:501:CYS:SG	1:B:531:ALA:HA	2.51	0.50
1:A:509:SER:OG	1:A:510:ALA:N	2.44	0.50
1:B:420:ARG:HD3	1:B:574:ASP:HA	1.94	0.49
1:B:387:MET:HB3	1:B:487:ARG:CZ	2.42	0.49
1:A:296:TYR:HB2	1:A:314:ILE:HG13	1.95	0.48
1:B:334:VAL:HG13	1:B:335:LEU:HD12	1.95	0.48
1:B:379:HIS:HD2	1:B:486:HIS:CG	2.32	0.48
1:B:523:LYS:NZ	1:B:563:LEU:O	2.36	0.48
1:A:486:HIS:O	1:A:487:ARG:HD3	2.15	0.47
1:B:460:ASP:OD1	1:B:473:TYR:OH	2.28	0.47
1:B:391:GLU:O	1:B:395:ILE:N	2.41	0.47
1:A:350:GLN:O	1:A:354:GLN:HB2	2.15	0.46
1:B:508:GLY:HA3	1:B:509:SER:HA	1.76	0.46
1:A:460:ASP:OD1	1:A:461:LEU:N	2.43	0.46
1:B:434:HIS:HA	1:B:435:PRO:HD2	1.79	0.46
1:A:326:ASP:HB3	1:A:327:HIS:H	1.34	0.46
1:B:388:PRO:HA	1:B:424:THR:HG22	1.99	0.45
1:B:557:LYS:HG2	1:B:567:TYR:CD1	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:403:ALA:HB1	1:A:410:LEU:HD23	1.99	0.45
1:A:255:HIS:CD2	1:A:255:HIS:H	2.36	0.44
1:A:456:PHE:HD2	1:A:457:LEU:HD12	1.83	0.44
1:A:325:LEU:HD12	1:A:354:GLN:NE2	2.25	0.44
1:A:461:LEU:C	1:A:462:VAL:CG2	2.91	0.44
1:A:321:HIS:HA	1:A:322:LEU:O	2.18	0.44
1:A:354:GLN:HB2	1:A:354:GLN:HE21	1.59	0.44
1:B:477:CYS:SG	1:B:478:ARG:N	2.91	0.44
1:A:488:ARG:NH1	1:A:490:ASP:OD2	2.50	0.44
1:A:451:LEU:HB3	1:A:457:LEU:HD13	2.00	0.43
1:A:461:LEU:N	1:A:461:LEU:HD12	2.33	0.43
1:A:324:LYS:C	1:A:325:LEU:HD22	2.35	0.43
1:A:461:LEU:O	1:A:462:VAL:HG22	2.19	0.43
1:A:472:LYS:HD3	1:A:474:LEU:HD21	2.01	0.43
1:A:555:THR:OG1	1:A:558:ASP:OD1	2.29	0.42
1:A:436:ASP:OD1	1:A:437:GLY:N	2.52	0.42
1:A:508:GLY:HA3	1:A:509:SER:HA	1.84	0.42
1:A:394:GLU:HB3	1:A:479:LEU:HD21	2.01	0.42
1:B:467:ASN:ND2	1:B:529:GLU:OE1	2.52	0.42
1:B:387:MET:HG3	1:B:425:CYS:O	2.19	0.42
1:A:400:GLN:HE21	1:A:404:GLN:NE2	2.18	0.41
1:A:434:HIS:CG	1:A:439:SER:HB2	2.55	0.41
1:A:452:ARG:HH21	1:A:460:ASP:HB2	1.84	0.41
1:A:260:LEU:HD23	1:A:260:LEU:HA	1.95	0.41
1:A:458:THR:O	1:A:476:VAL:HG22	2.21	0.41
1:B:458:THR:OG1	1:B:459:ASP:N	2.53	0.41
1:A:253:ASN:N	1:A:253:ASN:OD1	2.54	0.41
1:A:419:ARG:NH1	1:A:507:THR:OG1	2.49	0.41
1:A:486:HIS:C	1:A:487:ARG:HD3	2.46	0.40
1:A:461:LEU:N	1:A:461:LEU:CD1	2.84	0.40
1:A:323:ARG:C	1:A:325:LEU:HD22	2.47	0.40
1:B:419:ARG:HG2	1:B:507:THR:HG22	2.04	0.40
1:B:416:GLY:H	1:B:507:THR:HA	1.86	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:485:ARG:CZ	1:A:485:ARG:NH1[6_568]	1.33	0.87
1:A:485:ARG:CZ	1:A:485:ARG:CZ[6_568]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/326 (99%)	316 (98%)	8 (2%)	0	100	100
1	B	243/326 (74%)	232 (96%)	10 (4%)	1 (0%)	30	60
All	All	567/652 (87%)	548 (97%)	18 (3%)	1 (0%)	43	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	435	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/274 (100%)	264 (96%)	10 (4%)	31	58
1	B	206/274 (75%)	202 (98%)	4 (2%)	50	68
All	All	480/548 (88%)	466 (97%)	14 (3%)	37	62

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	321	HIS
1	A	323	ARG
1	A	325	LEU
1	A	354	GLN
1	A	458	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	462	VAL
1	A	476	VAL
1	A	488	ARG
1	A	529	GLU
1	A	536	VAL
1	B	332	VAL
1	B	461	LEU
1	B	474	LEU
1	B	476	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	354	GLN
1	A	379	HIS
1	A	400	GLN
1	A	434	HIS
1	A	471	GLN
1	B	379	HIS
1	B	400	GLN
1	B	464	GLN
1	B	469	GLN
1	B	511	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/326 (100%)	-0.48	1 (0%) 90 82	25, 58, 100, 159	0
1	B	245/326 (75%)	-0.19	0 100 100	25, 85, 128, 168	0
All	All	571/652 (87%)	-0.36	1 (0%) 91 86	25, 67, 125, 168	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	322	LEU	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.