



# wwPDB X-ray Structure Validation Summary Report

Mar 8, 2026 – 02:04 PM UTC

PDB ID : 4CC9 / pdb\_00004cc9  
Title : Crystal structure of human SAMHD1 (amino acid residues 582-626) bound to Vpx isolated from sooty mangabey and human DCAF1 (amino acid residues 1058-1396)  
Authors : Schwefel, D.; Groom, H.C.T.; Boucherit, V.C.; Christodoulou, E.; Walker, P.A.; Stoye, J.P.; Bishop, K.N.; Taylor, I.A.  
Deposited on : 2013-10-19  
Resolution : 2.47 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	 3% 76% 9% 15%
2	B	119	 3% 74% 7% 18%
3	C	71	 20% 7% 73%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 3459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN VPRBP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	2448	1547	424	460	17	54	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1057	MET	-	expression tag	UNP Q9Y4B6
A	1397	GLU	-	expression tag	UNP Q9Y4B6
A	1398	LEU	-	expression tag	UNP Q9Y4B6
A	1399	ALA	-	expression tag	UNP Q9Y4B6
A	1400	LEU	-	expression tag	UNP Q9Y4B6
A	1401	VAL	-	expression tag	UNP Q9Y4B6
A	1402	PRO	-	expression tag	UNP Q9Y4B6
A	1403	ARG	-	expression tag	UNP Q9Y4B6
A	1404	GLY	-	expression tag	UNP Q9Y4B6
A	1405	SER	-	expression tag	UNP Q9Y4B6
A	1406	SER	-	expression tag	UNP Q9Y4B6
A	1407	ALA	-	expression tag	UNP Q9Y4B6
A	1408	HIS	-	expression tag	UNP Q9Y4B6
A	1409	HIS	-	expression tag	UNP Q9Y4B6
A	1410	HIS	-	expression tag	UNP Q9Y4B6
A	1411	HIS	-	expression tag	UNP Q9Y4B6
A	1412	HIS	-	expression tag	UNP Q9Y4B6
A	1413	HIS	-	expression tag	UNP Q9Y4B6
A	1414	HIS	-	expression tag	UNP Q9Y4B6
A	1415	HIS	-	expression tag	UNP Q9Y4B6
A	1416	HIS	-	expression tag	UNP Q9Y4B6
A	1417	HIS	-	expression tag	UNP Q9Y4B6

- Molecule 2 is a protein called PROTEIN VPX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	98	801	511	144	138	8	20	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	GLY	-	expression tag	UNP P19508
B	-5	PRO	-	expression tag	UNP P19508
B	-4	GLY	-	expression tag	UNP P19508
B	-3	TYR	-	expression tag	UNP P19508
B	-2	GLN	-	expression tag	UNP P19508
B	-1	ASP	-	expression tag	UNP P19508
B	0	PRO	-	expression tag	UNP P19508

- Molecule 3 is a protein called DEOXYNUCLEOSIDE TRIPHOSPHATE TRIPHOSPHO-HYDROLASE SAMHD1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	19	158	96	32	30	4	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	556	MET	-	expression tag	UNP Q9Y3Z3
C	557	ALA	-	expression tag	UNP Q9Y3Z3
C	558	SER	-	expression tag	UNP Q9Y3Z3
C	559	TRP	-	expression tag	UNP Q9Y3Z3
C	560	SER	-	expression tag	UNP Q9Y3Z3
C	561	HIS	-	expression tag	UNP Q9Y3Z3
C	562	PRO	-	expression tag	UNP Q9Y3Z3
C	563	GLN	-	expression tag	UNP Q9Y3Z3
C	564	PHE	-	expression tag	UNP Q9Y3Z3
C	565	GLU	-	expression tag	UNP Q9Y3Z3
C	566	LYS	-	expression tag	UNP Q9Y3Z3
C	567	GLY	-	expression tag	UNP Q9Y3Z3
C	568	ALA	-	expression tag	UNP Q9Y3Z3
C	569	LEU	-	expression tag	UNP Q9Y3Z3
C	570	GLU	-	expression tag	UNP Q9Y3Z3
C	571	VAL	-	expression tag	UNP Q9Y3Z3
C	572	LEU	-	expression tag	UNP Q9Y3Z3
C	573	PHE	-	expression tag	UNP Q9Y3Z3
C	574	GLN	-	expression tag	UNP Q9Y3Z3

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	575	GLY	-	expression tag	UNP Q9Y3Z3
C	576	PRO	-	expression tag	UNP Q9Y3Z3
C	577	GLY	-	expression tag	UNP Q9Y3Z3
C	578	TYR	-	expression tag	UNP Q9Y3Z3
C	579	GLN	-	expression tag	UNP Q9Y3Z3
C	580	ASP	-	expression tag	UNP Q9Y3Z3
C	581	PRO	-	expression tag	UNP Q9Y3Z3

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 13 8 5	0	0
4	A	1	Total C O 13 8 5	0	0

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Zn 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	21	Total O 21 21	0	0
6	B	3	Total O 3 3	0	0
6	C	1	Total O 1 1	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.25Å 82.88Å 115.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.23 – 2.47 29.23 – 2.47	Depositor EDS
% Data completeness (in resolution range)	97.8 (29.23-2.47) 97.6 (29.23-2.47)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.91 (at 2.48Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.176 , 0.216 0.189 , 0.224	Depositor DCC
$R_{free}$ test set	1303 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.8	Xtrriage
Anisotropy	0.208	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 59.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3459	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2505	0.83	1/3394 (0.0%)
2	B	0.53	0/828	0.83	2/1123 (0.2%)
3	C	0.53	0/159	0.79	0/211
All	All	0.53	0/3492	0.83	3/4728 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1383	VAL	N-CA-CB	-6.00	104.86	112.60
2	B	100	ARG	CA-C-N	-5.31	114.45	119.76
2	B	100	ARG	C-N-CA	-5.31	114.45	119.76

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2448	0	2354	13	0
2	B	801	0	765	6	0
3	C	158	0	161	1	0
4	A	26	0	36	0	0
5	B	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	21	0	0	0	0
6	B	3	0	0	0	0
6	C	1	0	0	0	0
All	All	3459	0	3316	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1329:PRO:HG3	2:B:85:LYS:HD3	1.69	0.73
2:B:16:GLU:HG3	2:B:20:GLU:HG3	1.73	0.68
1:A:1150:LEU:HD22	1:A:1166:MET:HE2	1.82	0.61
1:A:1245:ASP:HB2	1:A:1252:ILE:HD11	1.83	0.60
1:A:1158:GLN:NE2	1:A:1177:THR:OG1	2.35	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/361 (84%)	294 (97%)	8 (3%)	1 (0%)	36	53
2	B	94/119 (79%)	91 (97%)	3 (3%)	0	100	100
3	C	17/71 (24%)	17 (100%)	0	0	100	100
All	All	414/551 (75%)	402 (97%)	11 (3%)	1 (0%)	43	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1167	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/316 (86%)	258 (95%)	13 (5%)	23	43
2	B	85/99 (86%)	81 (95%)	4 (5%)	23	44
3	C	18/63 (29%)	15 (83%)	3 (17%)	2	3
All	All	374/478 (78%)	354 (95%)	20 (5%)	20	39

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	29	VAL
3	C	617	ARG
3	C	624	ASP
3	C	618	VAL
1	A	1213	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1303	HIS
1	A	1353	ASN
1	A	1371	ASN
1	A	1174	HIS
1	A	1140	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PG4	A	2393	-	12,12,12	0.54	0	11,11,11	0.33	0
4	PG4	A	2394	-	12,12,12	0.57	0	11,11,11	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG4	A	2393	-	-	6/10/10/10	-
4	PG4	A	2394	-	-	5/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2393	PG4	O1-C1-C2-O2
4	A	2394	PG4	O3-C5-C6-O4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	A	2393	PG4	O4-C7-C8-O5
4	A	2394	PG4	C5-C6-O4-C7
4	A	2394	PG4	C8-C7-O4-C6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/361 (85%)	-0.50	1 (0%) 90 88	13, 29, 52, 62	13 (4%)
2	B	98/119 (82%)	-0.48	4 (4%) 41 37	14, 26, 68, 84	5 (5%)
3	C	19/71 (26%)	0.22	1 (5%) 32 28	22, 40, 65, 80	1 (5%)
All	All	424/551 (76%)	-0.46	6 (1%) 73 71	13, 29, 54, 84	19 (4%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	90	LEU	4.3
1	A	1314	GLN	4.0
2	B	88	ARG	3.3
3	C	624	ASP	2.4
2	B	100	ARG	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PG4	A	2394	13/13	0.76	0.21	49,68,71,75	0
4	PG4	A	2393	13/13	0.84	0.14	47,54,66,68	0
5	ZN	B	1112	1/1	0.99	0.03	43,43,43,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.