



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:55 PM UTC

PDB ID : 5CCG / pdb_00005ccg
Title : Structure of the Ca²⁺-bound synaptotagmin-1 SNARE complex (long unit cell form)
Authors : Zhou, Q.; Zhao, M.; Lyubimov, A.Y.; Uervirojnangkoorn, M.; Zeldin, O.B.; Weis, W.I.; Brunger, A.T.
Deposited on : 2015-07-02
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	63	 3% 75% 25%
1	G	63	 3% 83% 17%
2	B	67	 73% 25% .
2	H	67	 .% 70% 27% .
3	C	77	 .% 74% 22% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	77	 3% 75% 21% .
4	D	65	 75% 23% .
4	J	65	 77% 20% .
5	E	281	 4% 78% 21% ..
5	F	281	 77% 20% ..
5	K	281	 2% 78% 21% ..

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vesicle-associated membrane protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	63	Total	C	N	O	S	0	0	0
			502	305	95	101	1			
1	G	63	Total	C	N	O	S	0	0	0
			492	299	91	101	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	GLY	-	expression tag	UNP P63045
G	27	GLY	-	expression tag	UNP P63045

- Molecule 2 is a protein called Syntaxin-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	66	Total	C	N	O	S	0	0	0
			515	315	87	108	5			
2	H	65	Total	C	N	O	S	0	0	0
			510	314	84	107	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	190	MET	-	initiating methionine	UNP P32851
H	190	MET	-	initiating methionine	UNP P32851

- Molecule 3 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	74	Total	C	N	O	S	0	0	0
			579	343	102	129	5			
3	I	74	Total	C	N	O	S	0	0	0
			549	325	98	121	5			

- Molecule 4 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	64	Total	C	N	O	S	0	0	0
			498	291	96	106	5			
4	J	63	Total	C	N	O	S	0	0	0
			496	292	94	105	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	140	MET	-	initiating methionine	UNP P60881
J	140	MET	-	initiating methionine	UNP P60881

- Molecule 5 is a protein called Synaptotagmin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	279	Total	C	N	O	S	0	0	1
			2163	1391	363	402	7			
5	F	275	Total	C	N	O	S	0	0	1
			2074	1323	350	394	7			
5	K	279	Total	C	N	O	S	0	0	1
			2160	1390	360	403	7			

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total	0	0
			2		
6	C	3	Total	0	0
			3		
6	D	1	Total	0	0
			1		
6	E	4	Total	0	0
			4		
6	F	4	Total	0	0
			4		
6	G	1	Total	0	0
			1		
6	K	4	Total	0	0
			4		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O 1 1	0	0
7	B	1	Total O 1 1	0	0
7	D	1	Total O 1 1	0	0
7	E	4	Total O 4 4	0	0
7	F	4	Total O 4 4	0	0
7	I	1	Total O 1 1	0	0
7	J	1	Total O 1 1	0	0
7	K	5	Total O 5 5	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vesicle-associated membrane protein 2

Chain A:  75% 25%



- Molecule 1: Vesicle-associated membrane protein 2

Chain G:  3% 83% 17%



- Molecule 2: Syntaxin-1A

Chain B:  73% 25%




- Molecule 2: Syntaxin-1A

Chain H:  % 70% 27%




- Molecule 3: Synaptosomal-associated protein 25


Chain C:  % 74% 22%




- Molecule 3: Synaptosomal-associated protein 25




- Molecule 4: Synaptosomal-associated protein 25




- Molecule 4: Synaptosomal-associated protein 25




- Molecule 5: Synaptotagmin-1



- Molecule 5: Synaptotagmin-1



- Molecule 5: Synaptotagmin-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.60Å 171.09Å 291.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 3.50 19.97 – 3.50	Depositor EDS
% Data completeness (in resolution range)	87.6 (19.97-3.50) 87.0 (19.97-3.50)	Depositor EDS
R_{merge}	0.40	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 3.52Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.322 , 0.352 0.324 , 0.353	Depositor DCC
R_{free} test set	1999 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtrriage
Anisotropy	0.385	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.30$, $\langle L^2 \rangle = 0.13$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.72	EDS
Total number of atoms	10575	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.26	0/505	0.62	0/675
1	G	0.25	0/495	0.68	0/667
2	B	0.27	0/519	0.69	0/697
2	H	0.27	0/515	0.69	0/693
3	C	0.24	0/579	0.68	0/774
3	I	0.24	0/549	0.70	0/730
4	D	0.35	0/498	0.75	1/662 (0.2%)
4	J	0.26	0/496	0.70	0/661
5	E	0.40	1/2211 (0.0%)	0.83	6/2997 (0.2%)
5	F	0.33	0/2118	0.79	4/2872 (0.1%)
5	K	0.30	0/2207	0.77	7/2990 (0.2%)
All	All	0.32	1/10692 (0.0%)	0.76	18/14418 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	178	ASP	C-O	-5.30	1.20	1.24

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	K	186	LEU	N-CA-C	7.71	123.23	112.75
5	F	172	ASP	N-CA-C	7.51	120.82	110.24
5	E	180	TYR	CA-C-N	-7.11	113.97	122.93
5	E	180	TYR	C-N-CA	-7.11	113.97	122.93
5	F	186	LEU	C-N-CD	-6.92	96.61	125.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	502	0	498	17	0
1	G	492	0	472	10	0
2	B	515	0	476	19	0
2	H	510	0	479	25	0
3	C	579	0	538	17	0
3	I	549	0	506	17	0
4	D	498	0	476	17	0
4	J	496	0	480	12	0
5	E	2163	0	2110	49	0
5	F	2074	0	1928	43	1
5	K	2160	0	2102	43	0
6	A	2	0	0	0	0
6	C	3	0	0	0	0
6	D	1	0	0	0	0
6	E	4	0	0	0	0
6	F	4	0	0	0	0
6	G	1	0	0	0	0
6	K	4	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	D	1	0	0	0	0
7	E	4	0	0	1	0
7	F	4	0	0	0	0
7	I	1	0	0	0	0
7	J	1	0	0	0	0
7	K	5	0	0	0	0
All	All	10575	0	10065	216	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:186:LEU:CB	5:F:187:PRO:HD3	1.71	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:188:ASP:O	5:F:192:LYS:NZ	1.88	1.07
3:C:40:LYS:NZ	5:E:295:GLU:OE2	1.94	0.99
5:E:185:LEU:CD1	5:E:212:PHE:CD2	2.48	0.95
2:B:234:GLU:OE2	3:C:59:ARG:NH2	2.03	0.92

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:233:ARG:N	5:F:346:GLU:OE2[4_445]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	61/63 (97%)	61 (100%)	0	0	100	100
1	G	61/63 (97%)	61 (100%)	0	0	100	100
2	B	64/67 (96%)	63 (98%)	1 (2%)	0	100	100
2	H	63/67 (94%)	63 (100%)	0	0	100	100
3	C	72/77 (94%)	72 (100%)	0	0	100	100
3	I	72/77 (94%)	72 (100%)	0	0	100	100
4	D	62/65 (95%)	62 (100%)	0	0	100	100
4	J	61/65 (94%)	60 (98%)	1 (2%)	0	100	100
5	E	277/281 (99%)	262 (95%)	15 (5%)	0	100	100
5	F	269/281 (96%)	259 (96%)	10 (4%)	0	100	100
5	K	275/281 (98%)	259 (94%)	16 (6%)	0	100	100
All	All	1337/1387 (96%)	1294 (97%)	43 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	53/54 (98%)	53 (100%)	0	100	100
1	G	51/54 (94%)	51 (100%)	0	100	100
2	B	55/61 (90%)	55 (100%)	0	100	100
2	H	56/61 (92%)	56 (100%)	0	100	100
3	C	62/69 (90%)	62 (100%)	0	100	100
3	I	57/69 (83%)	57 (100%)	0	100	100
4	D	53/56 (95%)	53 (100%)	0	100	100
4	J	54/56 (96%)	54 (100%)	0	100	100
5	E	229/251 (91%)	227 (99%)	2 (1%)	70	76
5	F	212/251 (84%)	212 (100%)	0	100	100
5	K	229/251 (91%)	228 (100%)	1 (0%)	84	81
All	All	1111/1233 (90%)	1108 (100%)	3 (0%)	86	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	376	VAL
5	E	378	VAL
5	K	376	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	396	ASN
5	F	146	GLN
5	K	396	ASN
5	K	333	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	156	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 19 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	63/63 (100%)	0.15	0 100 100	23, 38, 60, 72	0
1	G	63/63 (100%)	0.25	2 (3%) 50 29	44, 57, 68, 72	0
2	B	66/67 (98%)	0.07	0 100 100	25, 39, 54, 58	0
2	H	65/67 (97%)	0.34	1 (1%) 72 47	37, 50, 67, 77	0
3	C	74/77 (96%)	0.14	1 (1%) 73 48	25, 43, 76, 93	0
3	I	74/77 (96%)	0.11	2 (2%) 56 33	35, 50, 65, 78	0
4	D	64/65 (98%)	0.07	0 100 100	23, 38, 56, 67	0
4	J	63/65 (96%)	0.19	0 100 100	36, 54, 72, 90	0
5	E	279/281 (99%)	0.41	10 (3%) 46 25	23, 51, 67, 78	0
5	F	275/281 (97%)	0.21	1 (0%) 88 72	26, 50, 74, 95	0
5	K	279/281 (99%)	0.32	7 (2%) 58 35	25, 49, 70, 86	0
All	All	1365/1387 (98%)	0.26	24 (1%) 67 42	23, 49, 70, 95	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	374	GLY	3.6
5	E	304	VAL	3.4
5	K	186	LEU	3.4
5	K	372	ALA	3.0
5	K	176	THR	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	CA	D	301	1/1	0.87	0.07	27,27,27,27	0
6	CA	F	503	1/1	0.89	0.09	59,59,59,59	0
6	CA	C	101	1/1	0.91	0.18	22,22,22,22	0
6	CA	F	501	1/1	0.92	0.10	23,23,23,23	0
6	CA	K	501	1/1	0.92	0.04	53,53,53,53	0
6	CA	E	503	1/1	0.94	0.05	22,22,22,22	0
6	CA	F	504	1/1	0.95	0.05	22,22,22,22	0
6	CA	C	102	1/1	0.95	0.04	22,22,22,22	0
6	CA	G	101	1/1	0.96	0.03	30,30,30,30	0
6	CA	K	502	1/1	0.96	0.06	52,52,52,52	0
6	CA	A	102	1/1	0.97	0.03	22,22,22,22	0
6	CA	F	502	1/1	0.98	0.13	24,24,24,24	0
6	CA	E	504	1/1	0.98	0.13	46,46,46,46	0
6	CA	A	101	1/1	0.98	0.11	24,24,24,24	0
6	CA	K	503	1/1	0.98	0.07	23,23,23,23	0
6	CA	E	501	1/1	0.99	0.10	37,37,37,37	0
6	CA	E	502	1/1	0.99	0.09	35,35,35,35	0
6	CA	C	103	1/1	0.99	0.04	22,22,22,22	0
6	CA	K	504	1/1	0.99	0.04	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.