



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:34 PM UTC

PDB ID : 7CC8 / pdb_00007cc8
Title : Crystal structure of White Spot Syndrome Virus Thymidylate Synthase - Apo form
Authors : Kumar, S.; Panchal, N.V.; Shaikh, N.; Vasudevan, D.
Deposited on : 2020-06-16
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	6% 87% 12% ..
1	B	297	5% 84% 10% 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate Synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	0	0
			2371	1514	414	426	17			
1	B	280	Total	C	N	O	S	0	0	0
			2258	1444	391	408	15			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	LEU	-	expression tag	UNP Q77J90
A	291	GLU	-	expression tag	UNP Q77J90
A	292	HIS	-	expression tag	UNP Q77J90
A	293	HIS	-	expression tag	UNP Q77J90
A	294	HIS	-	expression tag	UNP Q77J90
A	295	HIS	-	expression tag	UNP Q77J90
A	296	HIS	-	expression tag	UNP Q77J90
A	297	HIS	-	expression tag	UNP Q77J90
B	290	LEU	-	expression tag	UNP Q77J90
B	291	GLU	-	expression tag	UNP Q77J90
B	292	HIS	-	expression tag	UNP Q77J90
B	293	HIS	-	expression tag	UNP Q77J90
B	294	HIS	-	expression tag	UNP Q77J90
B	295	HIS	-	expression tag	UNP Q77J90
B	296	HIS	-	expression tag	UNP Q77J90
B	297	HIS	-	expression tag	UNP Q77J90

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			10	6	4		
3	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	91	Total	O	0	0
			91	91		
4	B	73	Total	O	0	0
			73	73		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thymidylate Synthase



• Molecule 1: Thymidylate Synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	91.80Å 91.80Å 233.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.41 – 2.30 43.41 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.9 (43.41-2.30) 98.8 (43.41-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8	Depositor
R, R_{free}	0.185 , 0.209 0.207 , 0.229	Depositor DCC
R_{free} test set	2277 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 18.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4850	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	15/2430 (0.6%)	1.39	7/3283 (0.2%)
1	B	1.34	8/2314 (0.3%)	1.29	4/3128 (0.1%)
All	All	1.38	23/4744 (0.5%)	1.34	11/6411 (0.2%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	278	PRO	C-O	-7.20	1.15	1.23
1	A	220	PRO	C-O	-6.91	1.16	1.23
1	A	231	HIS	CG-ND1	-6.88	1.30	1.38
1	A	171	HIS	CG-ND1	-6.76	1.30	1.38
1	B	171	HIS	CG-ND1	-6.22	1.31	1.38
1	B	216	VAL	C-O	-5.94	1.17	1.24
1	B	171	HIS	CD2-NE2	-5.88	1.31	1.37
1	A	138	ASN	C-O	-5.84	1.17	1.24
1	A	231	HIS	CD2-NE2	-5.79	1.31	1.37
1	A	65	TRP	NE1-CE2	-5.66	1.31	1.37
1	B	178	VAL	C-O	-5.52	1.18	1.24
1	A	109	VAL	C-O	-5.39	1.17	1.24
1	A	92	PHE	C-O	5.36	1.30	1.24
1	A	149	ASP	C-O	5.35	1.30	1.23
1	B	131	LYS	C-O	5.34	1.30	1.24
1	A	293	HIS	CE1-NE2	5.29	1.37	1.32
1	A	210	HIS	CG-ND1	-5.26	1.32	1.38
1	B	161	ASP	C-O	-5.22	1.17	1.24
1	B	236	HIS	CG-ND1	-5.21	1.32	1.38
1	B	15	LEU	C-O	-5.21	1.18	1.24
1	A	171	HIS	CD2-NE2	-5.20	1.32	1.37
1	A	194	VAL	C-O	-5.14	1.17	1.24
1	A	283	LEU	C-O	5.11	1.29	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	70	ASN	OD1-CG-ND2	-6.03	116.57	122.60
1	B	34	GLY	O-C-N	-5.63	116.14	121.77
1	A	241	LYS	CA-C-O	-5.62	114.59	120.55
1	B	15	LEU	CA-C-O	-5.34	114.89	120.55
1	A	27	THR	CA-C-N	5.31	125.72	120.31
1	A	27	THR	C-N-CA	5.31	125.72	120.31
1	A	55	PHE	CA-C-N	5.18	127.17	120.44
1	A	55	PHE	C-N-CA	5.18	127.17	120.44
1	B	248	PRO	CB-CA-C	-5.13	104.66	111.23
1	B	78	LYS	CA-C-O	-5.06	114.39	120.10
1	A	198	VAL	O-C-N	-5.01	117.22	120.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2352	12	0
1	B	2258	0	2241	13	0
2	A	25	0	0	0	0
2	B	5	0	0	0	0
3	A	7	0	9	1	0
3	B	20	0	28	2	0
4	A	91	0	0	0	0
4	B	73	0	0	1	0
All	All	4850	0	4630	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ARG:HG2	1:A:101:ARG:HH11	1.57	0.70
1:B:171:HIS:HD2	1:B:205:TYR:OH	1.76	0.69
1:B:211:LEU:O	1:B:214:SER:HB2	1.99	0.63
1:B:138:ASN:HB3	3:B:302:PGE:H32	1.81	0.62
1:A:211:LEU:O	1:A:214:SER:HB2	2.00	0.61
1:A:165:MET:SD	1:A:169:PRO:HD3	2.44	0.56
1:A:101:ARG:HG2	1:A:101:ARG:NH1	2.16	0.56
1:A:68:ARG:HE	1:A:70:ASN:ND2	2.06	0.54
1:B:37:MET:CE	1:B:225:LEU:HD23	2.39	0.53
1:B:37:MET:HE3	1:B:225:LEU:HD23	1.89	0.53
1:A:157:TRP:CZ2	1:B:153:ILE:HB	2.45	0.52
1:A:139:ILE:HG13	3:A:306:PGE:H5	1.95	0.49
1:B:149:ASP:OD2	3:B:302:PGE:H52	2.13	0.48
1:A:33:PHE:HA	1:A:228:GLY:O	2.12	0.48
1:B:165:MET:SD	1:B:169:PRO:HD3	2.54	0.48
1:B:231:HIS:HE1	4:B:453:HOH:O	1.97	0.46
1:B:107:GLY:HA2	1:B:121:TYR:CE2	2.52	0.45
1:A:157:TRP:NE1	1:B:153:ILE:HG21	2.32	0.44
1:B:117:PHE:CE2	1:B:158:ASN:HB2	2.54	0.42
1:B:58:GLY:HA2	1:B:81:ILE:HD12	2.00	0.42
1:A:225:LEU:CD2	1:A:227:LEU:HG	2.49	0.42
1:A:101:ARG:HG3	1:A:105:ASP:HB3	2.01	0.41
1:A:101:ARG:HD2	1:A:105:ASP:CG	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/297 (98%)	284 (97%)	7 (2%)	1 (0%)	36	46
1	B	278/297 (94%)	269 (97%)	8 (3%)	1 (0%)	30	38
All	All	570/594 (96%)	553 (97%)	15 (3%)	2 (0%)	30	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	110	TYR
1	A	110	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/258 (99%)	249 (98%)	6 (2%)	43	62
1	B	243/258 (94%)	240 (99%)	3 (1%)	63	79
All	All	498/516 (96%)	489 (98%)	9 (2%)	51	70

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	21	LYS
1	A	27	THR
1	A	80	LYS
1	A	239	VAL
1	A	286	LYS
1	B	80	LYS
1	B	125	SER
1	B	238	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	B	5	HIS
1	B	171	HIS
1	B	279	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PGE	B	302	-	9,9,9	0.40	0	8,8,8	0.37	0
2	SO4	A	305	-	4,4,4	0.27	0	6,6,6	0.23	0
2	SO4	B	301	-	4,4,4	0.69	0	6,6,6	0.31	0
2	SO4	A	301	-	4,4,4	0.53	0	6,6,6	0.17	0
2	SO4	A	302	-	4,4,4	0.31	0	6,6,6	0.14	0
3	PGE	B	303	-	9,9,9	0.31	0	8,8,8	0.16	0
3	PGE	A	306	-	6,6,9	0.43	0	5,5,8	0.44	0
2	SO4	A	304	-	4,4,4	0.31	0	6,6,6	0.14	0
2	SO4	A	303	-	4,4,4	0.31	0	6,6,6	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	B	303	-	-	4/7/7/7	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	B	302	-	-	5/7/7/7	-
3	PGE	A	306	-	-	1/4/4/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	PGE	O2-C3-C4-O3
3	B	302	PGE	O1-C1-C2-O2
3	B	302	PGE	O3-C5-C6-O4
3	B	303	PGE	O1-C1-C2-O2
3	B	303	PGE	O3-C5-C6-O4
3	B	303	PGE	O2-C3-C4-O3
3	B	302	PGE	C6-C5-O3-C4
3	B	303	PGE	C6-C5-O3-C4
3	A	306	PGE	O2-C3-C4-O3
3	B	302	PGE	C3-C4-O3-C5

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	PGE	2	0
3	A	306	PGE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/297 (98%)	0.01	17 (5%) 29 31	23, 29, 56, 82	0
1	B	280/297 (94%)	0.12	14 (5%) 34 35	22, 31, 49, 93	0
All	All	574/594 (96%)	0.06	31 (5%) 31 33	22, 30, 55, 93	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	285	MET	4.3
1	B	24	ARG	4.2
1	B	1	MET	4.2
1	A	24	ARG	3.9
1	A	292	HIS	3.8
1	A	293	HIS	3.8
1	B	27	THR	3.7
1	A	286	LYS	3.7
1	B	25	THR	3.6
1	A	289	VAL	3.6
1	A	288	ALA	3.5
1	B	280	SER	3.4
1	A	294	HIS	3.3
1	A	287	MET	3.3
1	B	26	GLY	3.2
1	A	27	THR	2.8
1	A	290	LEU	2.7
1	A	114	TRP	2.7
1	A	283	LEU	2.6
1	B	92	PHE	2.6
1	B	23	ASP	2.4
1	B	238	GLU	2.4
1	A	26	GLY	2.3
1	B	80	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	28	GLY	2.2
1	A	291	GLU	2.2
1	B	22	ASP	2.1
1	B	234	ASN	2.1
1	B	124	CYS	2.0
1	A	282	ASN	2.0
1	A	21	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	303	5/5	0.84	0.16	103,105,121,125	0
2	SO4	A	302	5/5	0.85	0.10	81,83,94,103	0
2	SO4	A	304	5/5	0.85	0.18	90,102,111,115	0
3	PGE	B	302	10/10	0.85	0.24	45,60,70,77	0
3	PGE	A	306	7/10	0.86	0.21	40,49,58,64	0
2	SO4	A	305	5/5	0.89	0.20	80,108,120,127	0
3	PGE	B	303	10/10	0.89	0.17	64,69,72,74	0
2	SO4	A	301	5/5	0.96	0.10	54,56,63,71	0
2	SO4	B	301	5/5	0.97	0.14	45,51,56,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.