



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:04 AM UTC

PDB ID : 1CDG / pdb_00001cdg
Title : NUCLEOTIDE SEQUENCE AND X-RAY STRUCTURE OF CYCLODEX-
TRIN GLYCOSYLTRANSFERASE FROM BACILLUS CIRCULANS
STRAIN 251 IN A MALTOSE-DEPENDENT CRYSTAL FORM
Authors : Lawson, C.L.; Van Montfort, R.; Strokopytov, B.V.; Kalk, K.H.; Rozeboom,
H.J.; Dijkstra, B.W.
Deposited on : 1993-08-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	
2	B	2	
2	C	2	
2	D	2	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5806 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLODEXTRIN GLYCOSYL-TRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	686	Total	C	N	O	S	0	0	0
			5264	3321	900	1027	16			

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	471	Total	O	0	0
			471	471		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYCLODEXTRIN GLYCOSYL-TRANSFERASE




- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	120.36Å 110.94Å 66.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.00 81.57 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 80.2 (81.57-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.05 (at 2.00Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.164 , (Not available) (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 114.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5806	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	5/5394 (0.1%)	1.58	59/7352 (0.8%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	28	GLY	C-N	-5.89	1.28	1.33
1	A	378	ILE	C-N	-5.50	1.27	1.34
1	A	553	ILE	C-N	5.38	1.40	1.33
1	A	304	LYS	N-CA	-5.08	1.40	1.46
1	A	546	ASP	N-CA	-5.04	1.40	1.46

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	ASN	CB-CA-C	-7.19	102.62	111.82
1	A	579	PHE	CA-CB-CG	-7.08	106.72	113.80
1	A	29	ASN	CA-C-N	6.78	126.41	119.56
1	A	29	ASN	C-N-CA	6.78	126.41	119.56
1	A	233	HIS	CA-CB-CG	-6.70	107.10	113.80
1	A	668	HIS	N-CA-C	-6.66	102.22	110.41
1	A	327	HIS	CA-CB-CG	-6.57	107.23	113.80
1	A	652	PHE	CA-CB-CG	-6.34	107.46	113.80
1	A	11	ASN	CA-CB-CG	6.30	118.90	112.60
1	A	543	SER	N-CA-CB	6.25	121.17	111.24
1	A	639	ASP	CA-CB-CG	-6.24	106.36	112.60
1	A	65	TYR	N-CA-C	6.17	117.67	111.07
1	A	546	ASP	CA-CB-CG	-6.02	106.58	112.60
1	A	255	PHE	CA-C-N	-6.01	117.12	122.47
1	A	255	PHE	C-N-CA	-6.01	117.12	122.47
1	A	92	VAL	N-CA-C	5.93	116.84	108.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	511	PRO	CA-C-N	-5.91	114.23	122.86
1	A	511	PRO	C-N-CA	-5.91	114.23	122.86
1	A	336	ASN	N-CA-CB	-5.88	101.62	110.91
1	A	617	ASP	CA-C-O	5.84	123.08	119.29
1	A	29	ASN	O-C-N	5.83	125.31	121.23
1	A	412	ARG	N-CA-CB	5.80	119.17	110.65
1	A	333	HIS	CA-CB-CG	-5.79	108.01	113.80
1	A	554	PRO	CB-CA-C	-5.79	103.65	111.12
1	A	5	SER	N-CA-CB	-5.75	101.59	110.04
1	A	266	SER	CA-C-N	5.72	125.25	119.19
1	A	266	SER	C-N-CA	5.72	125.25	119.19
1	A	609	SER	N-CA-C	5.69	119.93	113.15
1	A	275	GLU	N-CA-C	5.64	121.58	114.31
1	A	336	ASN	CA-C-N	-5.57	113.11	122.92
1	A	336	ASN	C-N-CA	-5.57	113.11	122.92
1	A	182	ASP	N-CA-C	-5.57	105.69	112.88
1	A	525	SER	N-CA-CB	-5.57	101.73	110.69
1	A	308	GLU	N-CA-CB	5.56	118.14	110.07
1	A	412	ARG	NE-CZ-NH1	5.54	127.04	121.50
1	A	569	ALA	CA-C-N	-5.53	113.18	122.25
1	A	569	ALA	C-N-CA	-5.53	113.18	122.25
1	A	269	ASN	CA-CB-CG	5.49	118.09	112.60
1	A	184	SER	N-CA-C	5.49	116.94	111.07
1	A	402	PRO	N-CA-CB	5.47	108.95	103.48
1	A	298	ASP	CA-CB-CG	-5.44	107.16	112.60
1	A	277	GLY	N-CA-C	-5.38	107.92	114.92
1	A	663	GLU	CA-C-N	-5.31	115.57	121.83
1	A	663	GLU	C-N-CA	-5.31	115.57	121.83
1	A	479	ASN	CA-CB-CG	-5.31	107.29	112.60
1	A	148	GLN	CA-C-N	5.28	126.44	119.84
1	A	148	GLN	C-N-CA	5.28	126.44	119.84
1	A	191	TYR	N-CA-C	5.26	119.68	113.16
1	A	375	ARG	NE-CZ-NH1	5.22	126.72	121.50
1	A	539	ALA	N-CA-C	5.17	117.66	111.71
1	A	281	LEU	N-CA-C	-5.08	102.33	109.96
1	A	205	SER	N-CA-C	5.06	117.45	111.33
1	A	431	VAL	CA-C-N	-5.06	116.52	123.14
1	A	431	VAL	C-N-CA	-5.06	116.52	123.14
1	A	586	GLN	N-CA-C	5.04	117.82	109.85
1	A	640	VAL	N-CA-C	5.03	115.72	108.48
1	A	440	ALA	CA-C-N	5.03	125.04	120.21
1	A	440	ALA	C-N-CA	5.03	125.04	120.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	346	LEU	CB-CA-C	-5.02	103.00	110.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5264	0	5025	77	0
2	B	23	0	21	1	0
2	C	23	0	21	0	0
2	D	23	0	21	4	0
3	A	2	0	0	0	0
4	A	471	0	0	8	0
All	All	5806	0	5088	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ASN:HD21	1:A:91:GLY:H	1.10	0.91
1:A:560:ASN:HD21	1:A:578:ASN:HA	1.37	0.87
1:A:620:LYS:HE3	4:A:937:HOH:O	1.84	0.77
1:A:88:ASN:HD21	1:A:91:GLY:N	1.83	0.75
1:A:336:ASN:OD1	1:A:336:ASN:N	2.25	0.69
1:A:562:ASN:HB3	1:A:575:VAL:HG13	1.76	0.67
1:A:617:ASP:OD2	1:A:620:LYS:HE2	1.95	0.66
1:A:47:ARG:HD2	4:A:908:HOH:O	1.95	0.65
1:A:653:LEU:HD23	1:A:653:LEU:N	2.13	0.64
1:A:293:PHE:O	1:A:341:LYS:HD2	1.98	0.64
1:A:142:SER:HB2	1:A:143:PRO:HD2	1.81	0.63
2:D:1:GLC:H62	2:D:2:GLC:C5	2.29	0.63
1:A:142:SER:HB2	1:A:143:PRO:CD	2.30	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:1:GLC:H62	2:D:2:GLC:O5	2.01	0.60
1:A:374:ASN:N	1:A:374:ASN:OD1	2.35	0.59
1:A:395:ALA:HB3	1:A:396:PRO:HD3	1.83	0.59
1:A:664:GLY:N	1:A:685:GLN:O	2.32	0.59
1:A:445:SER:HB3	1:A:479:ASN:ND2	2.18	0.59
1:A:670:PHE:CE2	1:A:680:ILE:HD11	2.39	0.57
1:A:669:THR:HG22	1:A:670:PHE:N	2.20	0.57
1:A:674:SER:HB2	4:A:1126:HOH:O	2.05	0.55
1:A:26:SER:O	1:A:56:GLY:HA3	2.07	0.55
1:A:227:ARG:HG2	1:A:255:PHE:CE2	2.44	0.53
1:A:107:LYS:NZ	4:A:776:HOH:O	2.41	0.53
1:A:227:ARG:HH11	1:A:257:GLU:HB2	1.74	0.52
1:A:562:ASN:HB3	1:A:575:VAL:CG1	2.40	0.51
1:A:87:ILE:HD12	1:A:143:PRO:HG2	1.92	0.51
1:A:649:GLU:OE1	1:A:667:ASN:ND2	2.28	0.51
1:A:444:ILE:HD12	1:A:482:LEU:HB2	1.93	0.50
1:A:327:HIS:ND1	1:A:327:HIS:N	2.57	0.50
1:A:479:ASN:ND2	4:A:856:HOH:O	2.44	0.50
1:A:48:LEU:HD12	1:A:95:THR:HG23	1.93	0.50
1:A:148:GLN:N	1:A:149:PRO:HD3	2.26	0.50
1:A:2:PRO:HD2	1:A:5:SER:HB2	1.95	0.49
1:A:651:LYS:NZ	1:A:665:GLY:O	2.46	0.49
1:A:20:ILE:HG23	1:A:25:PHE:HB2	1.96	0.48
1:A:594:ASN:HB2	1:A:683:ASN:OD1	2.14	0.48
1:A:444:ILE:HD12	1:A:482:LEU:CB	2.44	0.48
1:A:603:ASN:HB3	1:A:624:PRO:HB3	1.96	0.48
1:A:542:THR:HG21	4:A:892:HOH:O	2.14	0.47
1:A:397:LEU:HA	1:A:400:CYS:SG	2.54	0.47
1:A:669:THR:CG2	1:A:670:PHE:N	2.78	0.47
1:A:304:LYS:O	1:A:308:GLU:HG2	2.15	0.47
2:D:1:GLC:H62	2:D:2:GLC:H5	1.95	0.47
1:A:227:ARG:HD2	1:A:227:ARG:C	2.40	0.47
1:A:330:GLU:HB3	1:A:369:GLY:HA2	1.96	0.46
1:A:594:ASN:ND2	1:A:682:VAL:O	2.48	0.46
1:A:627:ASN:HD21	2:D:2:GLC:H61	1.80	0.46
1:A:663:GLU:CD	1:A:668:HIS:HE2	2.21	0.46
1:A:29:ASN:ND2	1:A:32:ASN:ND2	2.64	0.46
1:A:564:LYS:HB3	1:A:564:LYS:HE3	1.82	0.46
1:A:630:VAL:HG12	1:A:631:TYR:CE1	2.51	0.45
1:A:680:ILE:HD13	1:A:680:ILE:HG21	1.74	0.45
1:A:331:ARG:NH2	1:A:360:TYR:O	2.35	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:GLN:HG2	4:A:883:HOH:O	2.16	0.45
1:A:445:SER:CB	1:A:479:ASN:ND2	2.80	0.44
1:A:664:GLY:O	1:A:685:GLN:HB2	2.17	0.44
1:A:371:ASP:OD1	1:A:372:PRO:HA	2.18	0.44
1:A:119:GLN:HE21	1:A:119:GLN:HB2	1.54	0.44
1:A:149:PRO:HA	1:A:154:ASN:ND2	2.33	0.44
1:A:413:TRP:CE2	2:B:1:GLC:H2	2.52	0.44
1:A:652:PHE:C	1:A:653:LEU:HD23	2.43	0.44
1:A:87:ILE:CD1	1:A:143:PRO:HG2	2.47	0.43
1:A:395:ALA:N	1:A:396:PRO:CD	2.80	0.43
1:A:258:TRP:CD1	1:A:258:TRP:C	2.96	0.43
1:A:378:ILE:HA	1:A:379:PRO:HD3	1.89	0.43
1:A:148:GLN:C	1:A:150:SER:H	2.27	0.43
1:A:194:LEU:O	1:A:195:TYR:HB2	2.19	0.43
1:A:670:PHE:CZ	1:A:680:ILE:HD11	2.54	0.42
1:A:75:TRP:C	1:A:75:TRP:CD1	2.97	0.41
1:A:148:GLN:C	1:A:150:SER:N	2.78	0.41
1:A:21:PHE:CE1	1:A:327:HIS:HB3	2.55	0.41
1:A:418:VAL:HA	1:A:434:VAL:O	2.21	0.41
1:A:172:GLN:O	1:A:173:ASN:HB2	2.21	0.41
1:A:331:ARG:HG2	4:A:809:HOH:O	2.20	0.41
1:A:378:ILE:HG13	1:A:379:PRO:HD2	2.03	0.41
1:A:445:SER:HB3	1:A:479:ASN:CG	2.45	0.41
1:A:88:ASN:ND2	1:A:88:ASN:C	2.78	0.40
1:A:164:LEU:HA	1:A:164:LEU:HD23	1.88	0.40
1:A:290:ARG:NH2	1:A:330:GLU:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	684/686 (100%)	655 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1	2	12,12,12	0.38	0	17,17,17	0.93	1 (5%)
2	GLC	B	2	2	11,11,12	0.41	0	15,15,17	0.92	1 (6%)
2	GLC	C	1	2	12,12,12	0.28	0	17,17,17	1.40	2 (11%)
2	GLC	C	2	2	11,11,12	0.46	0	15,15,17	0.88	0
2	GLC	D	1	2	12,12,12	0.33	0	17,17,17	1.86	2 (11%)
2	GLC	D	2	2	11,11,12	0.27	0	15,15,17	2.08	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	1	2	-	2/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	GLC	C1-O5-C5	6.38	120.74	112.19
2	D	1	GLC	O4-C4-C3	-5.05	98.46	110.38
2	D	2	GLC	O5-C1-C2	3.74	119.71	110.79
2	C	1	GLC	O5-C1-C2	3.19	115.91	110.30
2	D	1	GLC	O5-C1-C2	3.01	115.60	110.30
2	B	2	GLC	O2-C2-C1	2.42	114.77	109.22
2	B	1	GLC	O5-C5-C4	-2.29	105.57	109.70
2	C	1	GLC	C1-C2-C3	2.17	114.79	110.36

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	GLC	O5-C5-C6-O6
2	D	2	GLC	O5-C5-C6-O6
2	C	1	GLC	C4-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	GLC	1	0
2	D	1	GLC	3	0
2	D	2	GLC	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	686/686 (100%)	-0.75	0 100 100	12, 23, 40, 76	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	2	11/12	0.60	0.15	84,86,88,90	0
2	GLC	D	1	12/12	0.79	0.11	77,79,81,82	0
2	GLC	C	1	12/12	0.79	0.12	68,74,77,79	0
2	GLC	C	2	11/12	0.83	0.10	60,62,64,65	0
2	GLC	B	2	11/12	0.91	0.07	35,37,38,40	0
2	GLC	B	1	12/12	0.96	0.05	25,32,37,38	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain B:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	692	1/1	0.99	0.04	21,21,21,21	0
3	CA	A	691	1/1	1.00	0.04	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.