



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:51 PM UTC

PDB ID : 1CDW / pdb_00001cdw
Title : HUMAN TBP CORE DOMAIN COMPLEXED WITH DNA
Authors : Nikolov, D.B.; Chen, H.; Halay, E.D.; Hoffmann, A.; Roeder, R.G.; Burley, S.K.
Deposited on : 1996-04-11
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	B	16	100%
2	C	16	81% 19%
3	A	179	78% 16% 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*TP*GP*CP*TP*AP*TP*AP*AP*AP*AP*GP*GP*CP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	16	Total	C	N	O	P	0	0	0
			327	157	62	93	15			

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*AP*GP*CP*CP*TP*TP*TP*TP*AP*TP*AP*GP*CP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	16	Total	C	N	O	P	0	0	0
			323	156	57	95	15			

- Molecule 3 is a protein called PROTEIN (TATA BINDING PROTEIN (TBP)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	179	Total	C	N	O	S	0	0	0
			1421	922	251	241	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	225	ASN	GLN	conflict	UNP P20226

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	41	Total	O	0	0
			41	41		
4	C	50	Total	O	0	0
			50	50		
4	A	189	Total	O	0	0
			189	189		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(*CP*TP*GP*CP*TP*AP*TP*AP*AP*AP*AP*GP*GP*CP*TP*G)-3')

Chain B:  100%

There are no outlier residues recorded for this chain.


- Molecule 2: DNA (5'-D(*CP*AP*GP*CP*CP*TP*TP*TP*TP*AP*TP*AP*GP*CP*AP*G)-3')

Chain C:  81% 19%



- Molecule 3: PROTEIN (TATA BINDING PROTEIN (TBP))

Chain A:  78% 16% 6%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.80Å 78.00Å 97.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.90	Depositor
% Data completeness (in resolution range)	88.4 (6.00-1.90)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.189 , 0.258	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2351	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.57	0/367	1.09	0/565
2	C	0.59	0/361	1.09	0/555
3	A	0.59	0/1447	0.98	6/1947 (0.3%)
All	All	0.59	0/2175	1.02	6/3067 (0.2%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	277	SER	N-CA-C	6.00	119.50	109.95
3	A	283	LEU	N-CA-C	-5.66	105.41	112.93
3	A	156	GLY	N-CA-C	-5.61	107.52	115.43
3	A	182	ARG	N-CA-C	5.58	120.21	112.45
3	A	247	LEU	N-CA-C	5.47	118.73	109.76
3	A	311	ALA	N-CA-C	5.42	118.14	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	327	0	182	0	0
2	C	323	0	183	2	0
3	A	1421	0	1512	32	0
4	A	189	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	41	0	0	0	0
4	C	50	0	0	1	0
All	All	2351	0	1877	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:296:ILE:HD11	3:A:308:LEU:HB3	1.59	0.84
3:A:184:ARG:H	3:A:184:ARG:HD2	1.48	0.77
3:A:231:ARG:HH21	3:A:235:ARG:NH1	1.88	0.72
3:A:279:TYR:HB2	3:A:287:LEU:HD22	1.72	0.71
3:A:231:ARG:O	3:A:235:ARG:HD2	1.94	0.68
3:A:201:ARG:HG3	4:A:1096:HOH:O	1.95	0.67
3:A:296:ILE:CD1	3:A:308:LEU:HB3	2.26	0.63
3:A:235:ARG:HG3	4:A:1249:HOH:O	1.99	0.62
3:A:231:ARG:HH21	3:A:235:ARG:HH12	1.48	0.60
3:A:253:ASN:HD21	3:A:310:GLY:H	1.51	0.59
3:A:235:ARG:HD3	4:A:1267:HOH:O	2.02	0.58
3:A:296:ILE:HB	4:A:1232:HOH:O	2.06	0.55
3:A:231:ARG:HG2	3:A:246:PHE:CZ	2.46	0.51
3:A:326:PRO:HA	3:A:329:LYS:HZ3	1.76	0.51
2:C:105:DC:H2'	2:C:106:DT:H71	1.93	0.51
3:A:267:GLU:O	3:A:271:LEU:HG	2.11	0.51
3:A:223:GLU:OE1	3:A:314:ARG:NH1	2.46	0.48
3:A:265:ARG:NH2	3:A:268:GLY:HA3	2.29	0.48
3:A:223:GLU:OE2	3:A:314:ARG:HD2	2.13	0.48
3:A:299:LEU:N	3:A:299:LEU:HD12	2.29	0.47
3:A:294:PRO:HG2	3:A:296:ILE:CG2	2.46	0.46
3:A:235:ARG:HG3	4:A:1231:HOH:O	2.16	0.46
3:A:231:ARG:HE	3:A:235:ARG:HH11	1.64	0.45
3:A:296:ILE:HG13	3:A:297:VAL:N	2.29	0.45
3:A:231:ARG:HE	3:A:235:ARG:NH1	2.15	0.45
3:A:263:PRO:HA	3:A:302:VAL:O	2.17	0.45
3:A:176:LEU:HG	3:A:196:VAL:HG23	1.99	0.44
3:A:261:LYS:HE3	3:A:261:LYS:HB2	1.76	0.43
3:A:267:GLU:H	3:A:267:GLU:CD	2.27	0.43
3:A:326:PRO:HA	3:A:329:LYS:NZ	2.34	0.42
3:A:274:GLN:HG2	4:A:1109:HOH:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:253:ASN:HD21	3:A:310:GLY:N	2.15	0.41
2:C:112:DA:H8	4:C:1118:HOH:O	2.03	0.41
3:A:296:ILE:HG21	3:A:320:ALA:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	177/179 (99%)	174 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	154/154 (100%)	138 (90%)	16 (10%)	7	2

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	174	LEU
3	A	176	LEU
3	A	184	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	196	VAL
3	A	201	ARG
3	A	225	ASN
3	A	231	ARG
3	A	235	ARG
3	A	240	LEU
3	A	267	GLU
3	A	274	GLN
3	A	282	GLU
3	A	283	LEU
3	A	287	LEU
3	A	295	ARG
3	A	296	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	189	ASN
3	A	225	ASN
3	A	253	ASN
3	A	273	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.