



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:18 PM UTC

PDB ID : 2CDO / pdb\_00002cdo  
Title : structure of agarase carbohydrate binding module in complex with neoagaro-hexaose  
Authors : Henshaw, J.; Horne, A.; Van Bueren, A.L.; Money, V.A.; Bolam, D.N.; Czjzek, M.; Weiner, R.M.; Hutcheson, S.W.; Davies, G.J.; Boraston, A.B.; Gilbert, H.J.  
Deposited on : 2006-01-25  
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	 2% 74% 11% 14%
1	B	160	 % 72% 13% 14%
1	C	160	 5% 71% 13% 15%
1	D	160	 2% 71% 14% 14%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	E	6	 17% 83%
2	F	6	 83% 17%
2	G	6	 83% 17%
2	H	6	 17% 83%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GAL	E	1	X	-	-	-
2	GAL	F	1	X	-	-	-
2	GAL	G	1	X	-	-	-
2	GAL	H	1	X	-	-	-
5	EDO	C	1146	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-AGARASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	138	1046	651	177	215	3	0	4	0
1	B	137	1052	656	179	214	3	0	6	0
1	C	136	1039	648	177	211	3	0	5	0
1	D	137	1026	638	175	210	3	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	MSE	ILE	engineered mutation	UNP Q6DN99
A	77	MSE	LEU	engineered mutation	UNP Q6DN99
B	5	MSE	ILE	engineered mutation	UNP Q6DN99
B	77	MSE	LEU	engineered mutation	UNP Q6DN99
C	5	MSE	ILE	engineered mutation	UNP Q6DN99
C	77	MSE	LEU	engineered mutation	UNP Q6DN99
D	5	MSE	ILE	engineered mutation	UNP Q6DN99
D	77	MSE	LEU	engineered mutation	UNP Q6DN99

- Molecule 2 is an oligosaccharide called 3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	E	6	64	36	28	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	F	6	Total	C	O	0	0	0
			64	36	28			
2	G	6	Total	C	O	0	0	0
			64	36	28			
2	H	6	Total	C	O	0	0	0
			64	36	28			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		
3	B	2	Total	Ca	0	0
			2	2		
3	C	2	Total	Ca	0	0
			2	2		
3	D	2	Total	Ca	0	0
			2	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Cl	0	0
			3	3		
4	B	2	Total	Cl	0	0
			2	2		
4	C	3	Total	Cl	0	0
			3	3		
4	D	2	Total	Cl	0	0
			2	2		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0



- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	154	Total O 154 154	0	0
6	B	159	Total O 159 159	0	0
6	C	139	Total O 139 139	0	0
6	D	147	Total O 147 147	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: BETA-AGARASE 1



- Molecule 1: BETA-AGARASE 1




- Molecule 1: BETA-AGARASE 1



- Molecule 1: BETA-AGARASE 1





- Molecule 2: 3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose

Chain E:  17% 83%


 GAL1  
AAL2  
GAL3  
AAL4  
GAL5  
AAL6


- Molecule 2: 3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose

Chain F:  83% 17%

 GAL1  
AAL2  
GAL3  
AAL4  
GAL5  
AAL6


- Molecule 2: 3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose

Chain G:  83% 17%

 GAL1  
AAL2  
GAL3  
AAL4  
GAL5  
AAL6

- Molecule 2: 3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-3,6-anhydro-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose

Chain H:  17% 83%

 GAL1  
AAL2  
GAL3  
AAL4  
GAL5  
AAL6

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.04Å 54.98Å 196.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	98.53 – 1.64 98.41 – 1.65	Depositor EDS
% Data completeness (in resolution range)	69.4 (98.53-1.64) 69.4 (98.41-1.65)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 1.64Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.176 , 0.241 0.176 , 0.241	Depositor DCC
$R_{free}$ test set	2565 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.3	Xtrriage
Anisotropy	0.170	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5064	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AAL, CA, CL, GAL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.04	1/1077 (0.1%)	0.88	0/1468
1	B	1.07	0/1092	0.86	0/1488
1	C	1.02	0/1076	0.92	0/1466
1	D	1.09	1/1051 (0.1%)	1.04	2/1433 (0.1%)
All	All	1.06	2/4296 (0.0%)	0.93	2/5855 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	25	GLN	N-CA	7.78	1.52	1.45
1	A	53	VAL	CA-CB	5.39	1.60	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	24	ALA	N-CA-C	-17.45	90.04	112.90
1	D	138	ASN	N-CA-C	6.47	129.12	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	23	GLN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1046	0	967	15	0
1	B	1052	0	978	19	0
1	C	1039	0	964	23	0
1	D	1026	0	938	17	0
2	E	64	0	51	0	0
2	F	64	0	51	1	0
2	G	64	0	51	1	0
2	H	64	0	51	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
4	C	3	0	0	0	0
4	D	2	0	0	0	0
5	B	12	0	18	5	0
5	C	12	0	17	8	0
5	D	4	0	6	0	0
6	A	154	0	0	1	0
6	B	159	0	0	1	0
6	C	139	0	0	5	0
6	D	147	0	0	6	0
All	All	5064	0	4092	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:VAL:HG21	1:A:51[B]:ILE:HD11	1.51	0.92
1:C:108:VAL:HG23	5:C:1146:EDO:H12	1.53	0.88
1:D:55:GLN:HG2	6:D:2059:HOH:O	1.76	0.86
1:C:109:TYR:H	5:C:1146:EDO:H21	1.42	0.85
1:B:58:ASN:HB2	5:B:1142:EDO:H21	1.62	0.80
1:C:87:LYS:HD2	5:C:1146:EDO:H11	1.70	0.73
1:C:108:VAL:CG2	5:C:1146:EDO:H12	2.19	0.73
1:B:58:ASN:HB2	5:B:1142:EDO:C2	2.18	0.72
5:B:1145:EDO:H22	6:D:2051:HOH:O	1.89	0.72
1:D:137:SER:O	1:D:138:ASN:HB2	1.90	0.70
1:A:64:GLN:HE22	1:A:102:PRO:HG3	1.56	0.70
1:D:111:SER:O	1:D:115:HIS:HE1	1.76	0.68
1:A:111:SER:O	1:A:115:HIS:HE1	1.77	0.67
1:C:111:SER:O	1:C:115:HIS:HE1	1.78	0.67
1:C:5[A]:MSE:HE3	1:C:136:LEU:HB2	1.79	0.64
1:C:108:VAL:HG23	5:C:1146:EDO:C1	2.28	0.64
1:C:133[A]:LYS:HD3	6:C:2120:HOH:O	1.98	0.63
1:A:115:HIS:HD2	6:A:2077:HOH:O	1.82	0.62
1:C:75:GLU:OE1	1:C:120:HIS:HD2	1.82	0.62
1:C:109:TYR:H	5:C:1146:EDO:C2	2.12	0.62
1:B:111:SER:O	1:B:115:HIS:HE1	1.83	0.61
1:C:115:HIS:HD2	6:C:2058:HOH:O	1.84	0.60
1:B:75:GLU:OE1	1:B:120:HIS:HD2	1.85	0.59
1:D:75:GLU:OE1	1:D:120:HIS:HD2	1.87	0.58
1:D:11:ASN:HB3	6:D:2009:HOH:O	2.04	0.57
1:A:12:PHE:CD1	1:A:15:VAL:HG13	2.41	0.56
1:B:59:TYR:CE1	1:B:138:ASN:HB2	2.41	0.55
1:D:115:HIS:HD2	6:D:2078:HOH:O	1.90	0.54
1:D:64:GLN:HE22	1:D:102:PRO:HG3	1.72	0.54
1:B:123:GLY:H	1:B:128:GLN:HE22	1.56	0.54
1:B:123:GLY:H	1:B:128:GLN:NE2	2.08	0.51
1:A:75:GLU:OE1	1:A:120:HIS:HD2	1.94	0.50
1:C:73:SER:H	1:C:128:GLN:HE22	1.59	0.50
1:B:44:GLY:HA2	1:B:120:HIS:HE1	1.77	0.50
1:D:123:GLY:H	1:D:128:GLN:NE2	2.10	0.50
1:C:58[B]:ASN:ND2	6:C:2047:HOH:O	2.36	0.48
1:D:73:SER:H	1:D:128:GLN:HE22	1.60	0.48
1:C:74:ILE:HG12	1:C:129:TRP:CE2	2.50	0.47
1:C:123:GLY:H	1:C:128:GLN:NE2	2.12	0.47
1:A:5[A]:MSE:CE	1:A:51[A]:ILE:CG2	2.92	0.47
1:B:58:ASN:HB2	5:B:1142:EDO:H22	1.96	0.47
1:D:73:SER:H	1:D:128:GLN:NE2	2.12	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:123:GLY:H	1:D:128:GLN:HE22	1.63	0.46
1:A:5[A]:MSE:CE	1:A:51[A]:ILE:HG21	2.46	0.46
1:B:125:ASN:HB3	1:B:127:TRP:O	2.15	0.46
1:D:5[B]:MSE:HE3	1:D:53:VAL:HG22	1.98	0.46
1:B:73:SER:H	1:B:128:GLN:HE22	1.64	0.46
1:C:73:SER:H	1:C:128:GLN:NE2	2.14	0.45
1:B:5[A]:MSE:HE3	1:B:53:VAL:HG22	1.99	0.45
1:A:80:GLU:O	1:A:81:ASN:HB2	2.16	0.45
1:C:6:ALA:HB1	6:C:2120:HOH:O	2.16	0.45
1:D:44:GLY:HA2	1:D:120:HIS:HE1	1.82	0.45
1:A:5[A]:MSE:HE3	1:A:51[A]:ILE:CG2	2.46	0.44
1:A:74:ILE:HG12	1:A:129:TRP:CE2	2.52	0.44
1:B:16:GLY:HA3	1:B:46:TYR:CE2	2.53	0.43
1:B:74:ILE:HG12	1:B:129:TRP:CE2	2.53	0.43
1:A:5[A]:MSE:HE1	1:A:53:VAL:HG22	2.00	0.43
1:C:44:GLY:HA2	1:C:120:HIS:HE1	1.83	0.43
1:C:123:GLY:H	1:C:128:GLN:HE22	1.66	0.42
1:B:73:SER:H	1:B:128:GLN:NE2	2.17	0.42
1:A:128:GLN:HB3	1:A:129:TRP:CD1	2.54	0.42
1:D:128:GLN:HB3	1:D:129:TRP:CD1	2.53	0.42
5:B:1145:EDO:C2	6:D:2051:HOH:O	2.60	0.42
1:A:50:THR:HG21	1:B:95:GLN:NE2	2.35	0.42
1:C:109:TYR:N	5:C:1146:EDO:H21	2.23	0.42
1:B:115:HIS:HD2	6:B:2079:HOH:O	2.03	0.41
1:C:97:TRP:CG	2:G:6:AAL:H5	2.55	0.41
1:B:5[B]:MSE:HE1	1:B:59:TYR:CE1	2.55	0.41
1:D:120:HIS:HE1	6:D:2120:HOH:O	2.04	0.41
1:A:58:ASN:HB2	1:C:13:ASN:HD22	1.85	0.41
1:D:44:GLY:HA2	1:D:120:HIS:CE1	2.55	0.41
1:C:133[B]:LYS:HB3	1:C:133[B]:LYS:HE2	1.89	0.40
5:C:1146:EDO:H22	6:C:2139:HOH:O	2.21	0.40
1:D:97:TRP:CG	2:F:6:AAL:H5	2.56	0.40
1:B:59:TYR:CD1	1:B:138:ASN:HB2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/160 (88%)	135 (96%)	5 (4%)	0	100	100
1	B	141/160 (88%)	139 (99%)	2 (1%)	0	100	100
1	C	139/160 (87%)	134 (96%)	5 (4%)	0	100	100
1	D	136/160 (85%)	133 (98%)	3 (2%)	0	100	100
All	All	556/640 (87%)	541 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/120 (92%)	110 (100%)	0	100	100
1	B	111/120 (92%)	110 (99%)	1 (1%)	70	54
1	C	109/120 (91%)	105 (96%)	4 (4%)	30	5
1	D	106/120 (88%)	103 (97%)	3 (3%)	38	12
All	All	436/480 (91%)	428 (98%)	8 (2%)	53	26

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	20	SER
1	C	4	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	133[A]	LYS
1	C	133[B]	LYS
1	C	137	SER
1	D	2	THR
1	D	133	LYS
1	D	138	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	GLN
1	A	25	GLN
1	A	55	GLN
1	A	64	GLN
1	A	115	HIS
1	A	116	GLN
1	A	120	HIS
1	B	13	ASN
1	B	25	GLN
1	B	81	ASN
1	B	95	GLN
1	B	115	HIS
1	B	116	GLN
1	B	120	HIS
1	B	128	GLN
1	C	13	ASN
1	C	25	GLN
1	C	35	ASN
1	C	81	ASN
1	C	115	HIS
1	C	120	HIS
1	C	128	GLN
1	D	23	GLN
1	D	25	GLN
1	D	64	GLN
1	D	95	GLN
1	D	115	HIS
1	D	116	GLN
1	D	120	HIS
1	D	128	GLN
1	D	138	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GAL	E	1	2	12,12,12	0.91	0	17,17,17	1.53	2 (11%)
2	AAL	E	2	2	11,11,12	0.64	0	13,16,18	1.19	2 (15%)
2	GAL	E	3	2	11,11,12	0.43	0	15,15,17	1.21	0
2	AAL	E	4	2	11,11,12	0.86	0	13,16,18	2.05	3 (23%)
2	GAL	E	5	2	11,11,12	0.50	0	15,15,17	1.24	2 (13%)
2	AAL	E	6	2	11,11,12	0.72	0	13,16,18	1.65	3 (23%)
2	GAL	F	1	2	12,12,12	0.90	0	17,17,17	2.45	7 (41%)
2	AAL	F	2	2	11,11,12	0.55	0	13,16,18	1.65	4 (30%)
2	GAL	F	3	2	11,11,12	0.61	0	15,15,17	1.23	1 (6%)
2	AAL	F	4	2	11,11,12	0.67	0	13,16,18	1.67	3 (23%)
2	GAL	F	5	2	11,11,12	0.69	0	15,15,17	1.35	2 (13%)
2	AAL	F	6	2	11,11,12	0.99	0	13,16,18	1.65	3 (23%)
2	GAL	G	1	2	12,12,12	0.80	0	17,17,17	2.98	7 (41%)
2	AAL	G	2	2	11,11,12	0.78	0	13,16,18	1.24	1 (7%)
2	GAL	G	3	2	11,11,12	0.56	0	15,15,17	1.05	1 (6%)
2	AAL	G	4	2	11,11,12	0.60	0	13,16,18	1.57	2 (15%)
2	GAL	G	5	2	11,11,12	0.50	0	15,15,17	1.29	2 (13%)
2	AAL	G	6	2	11,11,12	0.76	0	13,16,18	1.53	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GAL	H	1	2	12,12,12	0.76	0	17,17,17	0.80	0
2	AAL	H	2	2	11,11,12	0.71	0	13,16,18	1.71	5 (38%)
2	GAL	H	3	2	11,11,12	0.61	0	15,15,17	1.11	1 (6%)
2	AAL	H	4	2	11,11,12	0.81	0	13,16,18	1.31	1 (7%)
2	GAL	H	5	2	11,11,12	0.79	0	15,15,17	1.35	1 (6%)
2	AAL	H	6	2	11,11,12	0.85	0	13,16,18	1.42	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	E	1	2	1/1/5/5	0/2/22/22	0/1/1/1
2	AAL	E	2	2	-	-	0/3/2/2
2	GAL	E	3	2	-	0/2/19/22	0/1/1/1
2	AAL	E	4	2	-	-	0/3/2/2
2	GAL	E	5	2	-	0/2/19/22	0/1/1/1
2	AAL	E	6	2	-	-	0/3/2/2
2	GAL	F	1	2	1/1/5/5	0/2/22/22	0/1/1/1
2	AAL	F	2	2	-	-	0/3/2/2
2	GAL	F	3	2	-	0/2/19/22	0/1/1/1
2	AAL	F	4	2	-	-	0/3/2/2
2	GAL	F	5	2	-	0/2/19/22	0/1/1/1
2	AAL	F	6	2	-	-	0/3/2/2
2	GAL	G	1	2	1/1/5/5	0/2/22/22	0/1/1/1
2	AAL	G	2	2	-	-	0/3/2/2
2	GAL	G	3	2	-	2/2/19/22	0/1/1/1
2	AAL	G	4	2	-	-	0/3/2/2
2	GAL	G	5	2	-	0/2/19/22	0/1/1/1
2	AAL	G	6	2	-	-	0/3/2/2
2	GAL	H	1	2	1/1/5/5	2/2/22/22	0/1/1/1
2	AAL	H	2	2	-	-	0/3/2/2
2	GAL	H	3	2	-	0/2/19/22	0/1/1/1
2	AAL	H	4	2	-	-	0/3/2/2
2	GAL	H	5	2	-	0/2/19/22	0/1/1/1
2	AAL	H	6	2	-	-	0/3/2/2

There are no bond length outliers.

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	GAL	O4-C4-C3	-8.67	89.94	110.38
2	E	4	AAL	O5-C5-C6	-4.86	106.23	113.33
2	F	1	GAL	O3-C3-C2	4.84	121.79	110.38
2	F	1	GAL	C3-C4-C5	4.54	118.47	110.23
2	G	6	AAL	C1-O5-C5	4.36	118.03	112.19
2	H	5	GAL	O3-C3-C2	-4.32	101.25	110.05
2	E	6	AAL	C1-O5-C5	4.28	117.93	112.19
2	F	1	GAL	C4-C3-C2	4.14	118.10	110.83
2	G	1	GAL	C3-C4-C5	4.08	117.64	110.23
2	E	1	GAL	C1-C2-C3	3.85	118.20	110.36
2	G	1	GAL	C4-C3-C2	3.80	117.51	110.83
2	G	4	AAL	O5-C5-C6	-3.72	107.89	113.33
2	G	1	GAL	O3-C3-C4	3.69	119.08	110.38
2	G	5	GAL	O3-C3-C2	-3.67	102.56	110.05
2	E	4	AAL	C1-O5-C5	3.66	117.10	112.19
2	F	1	GAL	O5-C5-C4	3.66	116.29	109.70
2	G	1	GAL	C1-C2-C3	3.56	117.62	110.36
2	E	1	GAL	C4-C3-C2	3.54	117.04	110.83
2	F	4	AAL	C1-O5-C5	3.53	116.92	112.19
2	F	6	AAL	C2-C3-C4	-3.49	105.22	113.85
2	F	6	AAL	O5-C5-C6	-3.47	108.26	113.33
2	H	2	AAL	C1-O5-C5	3.24	116.53	112.19
2	F	1	GAL	C1-O5-C5	3.08	119.60	113.65
2	F	4	AAL	O5-C5-C6	-3.06	108.85	113.33
2	G	4	AAL	C1-O5-C5	2.95	116.14	112.19
2	F	5	GAL	O3-C3-C2	-2.94	104.05	110.05
2	F	2	AAL	O5-C5-C6	-2.90	109.09	113.33
2	H	6	AAL	C1-O5-C5	2.82	115.96	112.19
2	F	2	AAL	C1-O5-C5	2.76	115.88	112.19
2	H	6	AAL	O5-C5-C6	-2.58	109.57	113.33
2	H	4	AAL	C1-O5-C5	2.55	115.60	112.19
2	G	3	GAL	C1-O5-C5	2.55	115.60	112.19
2	G	2	AAL	C1-O5-C5	2.50	115.54	112.19
2	E	2	AAL	C2-C3-C4	-2.49	107.69	113.85
2	E	5	GAL	C1-C2-C3	2.49	113.27	109.64
2	E	2	AAL	O5-C5-C6	-2.38	109.85	113.33
2	E	4	AAL	C2-C3-C4	-2.38	107.97	113.85
2	F	4	AAL	O4-C4-C5	-2.36	104.29	111.08
2	F	1	GAL	O4-C4-C3	-2.35	104.83	110.38
2	E	6	AAL	C2-C3-C4	-2.35	108.04	113.85
2	E	5	GAL	O6-C6-C5	-2.32	103.42	111.33
2	F	2	AAL	C1-C2-C3	2.29	112.43	109.29
2	E	6	AAL	O5-C5-C6	-2.28	110.00	113.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	3	GAL	O2-C2-C3	2.26	114.84	110.15
2	H	6	AAL	C2-C3-C4	-2.26	108.27	113.85
2	G	1	GAL	O1-C1-C2	-2.25	102.44	108.98
2	F	2	AAL	C2-C3-C4	-2.25	108.28	113.85
2	H	2	AAL	O4-C4-C3	-2.24	104.91	111.19
2	H	2	AAL	C1-C2-C3	2.22	112.34	109.29
2	F	6	AAL	C1-O5-C5	2.22	115.17	112.19
2	H	2	AAL	C3-C4-C5	-2.20	97.26	101.99
2	H	2	AAL	O5-C5-C6	-2.09	110.28	113.33
2	F	1	GAL	O2-C2-C3	2.07	115.26	110.38
2	H	3	GAL	O5-C5-C6	2.05	111.65	107.66
2	G	5	GAL	O6-C6-C5	-2.03	104.41	111.33
2	F	5	GAL	O3-C3-C4	2.02	115.14	110.38
2	G	1	GAL	O3-C3-C2	2.02	115.13	110.38

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	E	1	GAL	C3
2	F	1	GAL	C3
2	G	1	GAL	C3
2	H	1	GAL	C3

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	H	1	GAL	C4-C5-C6-O6
2	H	1	GAL	O5-C5-C6-O6
2	G	3	GAL	O5-C5-C6-O6
2	G	3	GAL	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	6	AAL	1	0
2	F	6	AAL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 18 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	C	1141	-	3,3,3	0.61	0	2,2,2	0.22	0
5	EDO	B	1145	-	3,3,3	0.92	0	2,2,2	0.48	0
5	EDO	C	1146	-	3,3,3	0.72	0	2,2,2	0.61	0
5	EDO	C	1145	-	3,3,3	0.32	0	2,2,2	1.11	0
5	EDO	B	1141	-	3,3,3	0.39	0	2,2,2	0.50	0
5	EDO	D	1143	-	3,3,3	0.42	0	2,2,2	0.26	0
5	EDO	B	1142	-	3,3,3	0.33	0	2,2,2	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	1141	-	-	0/1/1/1	-
5	EDO	B	1145	-	-	1/1/1/1	-
5	EDO	C	1146	-	-	1/1/1/1	-
5	EDO	C	1145	-	-	0/1/1/1	-
5	EDO	B	1141	-	-	1/1/1/1	-
5	EDO	D	1143	-	-	1/1/1/1	-
5	EDO	B	1142	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1145	EDO	O1-C1-C2-O2
5	D	1143	EDO	O1-C1-C2-O2
5	B	1141	EDO	O1-C1-C2-O2
5	C	1146	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1145	EDO	2	0
5	C	1146	EDO	8	0
5	B	1142	EDO	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	136/160 (85%)	-0.11	3 (2%) 62 68	5, 13, 27, 36	3 (2%)
1	B	135/160 (84%)	-0.37	2 (1%) 72 77	5, 10, 22, 40	5 (3%)
1	C	134/160 (83%)	0.22	8 (5%) 27 31	6, 13, 33, 40	4 (2%)
1	D	135/160 (84%)	-0.13	4 (2%) 52 57	5, 13, 25, 31	0
All	All	540/640 (84%)	-0.09	17 (3%) 51 56	5, 13, 27, 40	12 (2%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	22	GLY	5.1
1	D	2	THR	4.0
1	C	3	ALA	3.9
1	C	19	PHE	3.8
1	C	18	THR	3.2
1	C	20	SER	3.2
1	A	81	ASN	3.1
1	B	2	THR	2.9
1	A	2	THR	2.9
1	D	34	GLY	2.5
1	B	20	SER	2.4
1	C	24	ALA	2.3
1	A	1	SER	2.3
1	C	138	ASN	2.2
1	D	138	ASN	2.1
1	C	29	VAL	2.0
1	D	82	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GAL	E	1	12/12	0.59	0.18	43,49,52,52	0
2	GAL	F	1	12/12	0.70	0.16	41,45,48,49	0
2	GAL	H	1	12/12	0.78	0.15	40,47,49,49	0
2	GAL	G	1	12/12	0.86	0.10	19,24,27,28	0
2	AAL	E	2	10/11	0.88	0.10	22,30,33,35	0
2	AAL	G	2	10/11	0.91	0.08	18,20,22,23	0
2	AAL	F	2	10/11	0.91	0.09	18,28,33,34	0
2	AAL	H	2	10/11	0.92	0.09	22,29,32,35	0
2	GAL	G	3	11/12	0.93	0.08	14,16,25,31	0
2	AAL	G	6	10/11	0.94	0.06	10,14,16,17	0
2	GAL	H	3	11/12	0.94	0.07	11,17,24,28	0
2	GAL	E	3	11/12	0.95	0.06	13,18,23,26	0
2	GAL	G	5	11/12	0.95	0.06	12,14,16,19	0
2	AAL	E	6	10/11	0.95	0.06	10,11,13,14	0
2	AAL	G	4	10/11	0.96	0.04	8,10,12,14	0
2	GAL	F	3	11/12	0.96	0.06	12,14,20,23	0
2	AAL	F	6	10/11	0.97	0.05	8,10,12,13	0
2	GAL	E	5	11/12	0.97	0.05	8,11,14,16	0
2	AAL	H	4	10/11	0.97	0.04	8,10,12,13	0
2	GAL	F	5	11/12	0.98	0.04	9,10,12,16	0
2	AAL	E	4	10/11	0.98	0.04	10,10,12,13	0
2	AAL	F	4	10/11	0.98	0.04	9,10,12,13	0
2	GAL	H	5	11/12	0.98	0.03	6,8,9,15	0
2	AAL	H	6	10/11	0.98	0.03	7,8,9,11	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	B	1145	4/4	0.86	0.15	18,19,21,26	0
5	EDO	D	1143	4/4	0.90	0.12	29,31,31,32	0
5	EDO	C	1146	4/4	0.91	0.18	9,15,20,22	0
5	EDO	B	1142	4/4	0.92	0.16	21,22,26,27	0
5	EDO	C	1145	4/4	0.94	0.10	11,18,20,21	0
3	CA	C	1140	1/1	0.94	0.04	17,17,17,17	0
4	CL	C	1143	1/1	0.94	0.18	35,35,35,35	0
4	CL	C	1142	1/1	0.96	0.07	27,27,27,27	0
4	CL	D	1142	1/1	0.97	0.07	23,23,23,23	0
3	CA	A	1139	1/1	0.97	0.05	14,14,14,14	0
4	CL	C	1144	1/1	0.97	0.05	24,24,24,24	0
3	CA	C	1139	1/1	0.98	0.04	14,14,14,14	0
4	CL	A	1141	1/1	0.98	0.05	24,24,24,24	0
5	EDO	C	1141	4/4	0.98	0.05	10,13,14,15	0
4	CL	A	1142	1/1	0.98	0.13	19,19,19,19	0
4	CL	D	1141	1/1	0.98	0.07	25,25,25,25	0
4	CL	B	1143	1/1	0.98	0.06	22,22,22,22	0
5	EDO	B	1141	4/4	0.99	0.03	16,16,16,20	0
4	CL	B	1144	1/1	0.99	0.04	23,23,23,23	0
3	CA	D	1139	1/1	0.99	0.02	13,13,13,13	0
3	CA	B	1140	1/1	0.99	0.04	10,10,10,10	0
3	CA	A	1140	1/1	0.99	0.02	15,15,15,15	0
4	CL	A	1143	1/1	0.99	0.09	21,21,21,21	0
3	CA	B	1139	1/1	0.99	0.01	10,10,10,10	0
3	CA	D	1140	1/1	1.00	0.01	11,11,11,11	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.