



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 09:01 PM UTC

PDB ID : 7CE9 / pdb\_00007ce9  
Title : PQQ-soaked Apo-methanol dehydrogenase (MDH) from *Methylococcus capsulatus* (Bath)  
Authors : Chuankhayan, P.; Chan, S.I.; Nareddy, P.K.R.; Tsai, I.K.; Tsai, Y.F.; Chen, K.H.-C.; Yu, S.S.-F.; Chen, C.J.  
Deposited on : 2020-06-22  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	573	92% 7% .
1	B	573	95% . .
1	C	573	93% 6%
1	D	573	93% 6% .

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	573	 93% 7%
1	H	573	 93% 6% .
1	M	573	 92% 8%
1	N	573	 94% 6% .
2	E	72	 93% 6% .
2	F	72	 11% 82% 15% ..
2	I	72	 3% 90% 7% ..
2	J	72	 90% 7% ..
2	K	72	 % 93% 6% .
2	L	72	 85% 11% ...
2	O	72	 8% 86% 11% ..
2	P	72	 4% 86% 11% ..

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 43200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methanol dehydrogenase protein, large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	573	4491	2871	765	832	23	0	0	0
1	B	573	4490	2871	765	831	23	0	0	0
1	C	573	4491	2871	765	832	23	0	0	0
1	D	573	4490	2871	765	831	23	0	0	0
1	G	573	4491	2871	765	832	23	0	0	0
1	H	573	4490	2871	765	831	23	0	0	0
1	M	573	4491	2871	765	832	23	0	0	0
1	N	573	4490	2871	765	831	23	0	0	0

- Molecule 2 is a protein called Methanol dehydrogenase [cytochrome c] subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	71	568	356	100	109	3	0	0	0
2	F	71	568	356	100	109	3	0	0	0
2	I	71	568	356	100	109	3	0	0	0
2	J	71	568	356	100	109	3	0	0	0
2	K	71	568	356	100	109	3	0	0	0
2	L	71	568	356	100	109	3	0	0	0

*Continued on next page...*



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			24	14	2	8		
4	B	1	Total	C	N	O	0	0
			24	14	2	8		
4	C	1	Total	C	N	O	0	0
			24	14	2	8		
4	D	1	Total	C	N	O	0	0
			24	14	2	8		
4	G	1	Total	C	N	O	0	0
			24	14	2	8		
4	H	1	Total	C	N	O	0	0
			24	14	2	8		
4	M	1	Total	C	N	O	0	0
			24	14	2	8		
4	N	1	Total	C	N	O	0	0
			24	14	2	8		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	254	Total	O	0	0
			254	254		
5	B	309	Total	O	0	0
			309	309		
5	C	286	Total	O	0	0
			286	286		
5	D	288	Total	O	0	0
			288	288		
5	E	34	Total	O	0	0
			34	34		
5	F	43	Total	O	0	0
			43	43		
5	G	250	Total	O	0	0
			250	250		
5	H	284	Total	O	0	0
			284	284		
5	I	25	Total	O	0	0
			25	25		
5	J	38	Total	O	0	0
			38	38		
5	K	48	Total	O	0	0
			48	48		
5	L	59	Total	O	0	0
			59	59		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	M	308	Total 308	O 308	0	0
5	N	238	Total 238	O 238	0	0
5	O	42	Total 42	O 42	0	0
5	P	26	Total 26	O 26	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methanol dehydrogenase protein, large subunit




- Molecule 1: Methanol dehydrogenase protein, large subunit



- Molecule 1: Methanol dehydrogenase protein, large subunit



- Molecule 1: Methanol dehydrogenase protein, large subunit







- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain G: 93% 7%





- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain H: 93% 6%



- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain M: 92% 8%



- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain N: 94% 6%




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2


Chain E: 93% 6%




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2




- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2



- Molecule 2: Methanol dehydrogenase [cytochrome c] subunit 2





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.56Å 211.76Å 223.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	153.00 – 2.20 153.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (153.00-2.20) 99.9 (153.00-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.10 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.161 , 0.209 0.171 , 0.215	Depositor DCC
$R_{free}$ test set	15016 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtrriage
Anisotropy	0.021	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 23.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	43200	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PQQ, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.15	8/4622 (0.2%)	1.07	9/6281 (0.1%)
1	B	1.17	4/4621 (0.1%)	1.10	9/6281 (0.1%)
1	C	1.18	8/4622 (0.2%)	1.10	14/6281 (0.2%)
1	D	1.19	5/4621 (0.1%)	1.08	9/6281 (0.1%)
1	G	1.14	4/4622 (0.1%)	1.06	6/6281 (0.1%)
1	H	1.17	2/4621 (0.0%)	1.10	8/6281 (0.1%)
1	M	1.14	8/4622 (0.2%)	1.08	11/6281 (0.2%)
1	N	1.12	3/4621 (0.1%)	1.06	9/6281 (0.1%)
2	E	1.19	1/583 (0.2%)	1.13	1/785 (0.1%)
2	F	1.25	1/583 (0.2%)	1.21	4/785 (0.5%)
2	I	1.10	1/583 (0.2%)	1.09	1/785 (0.1%)
2	J	1.22	0/583	1.23	1/785 (0.1%)
2	K	1.14	0/583	1.15	1/785 (0.1%)
2	L	1.17	1/583 (0.2%)	1.19	5/785 (0.6%)
2	O	1.26	0/583	1.11	2/785 (0.3%)
2	P	1.11	0/583	1.15	0/785
All	All	1.16	46/41636 (0.1%)	1.09	90/56528 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	G	0	1
1	H	0	3
1	M	0	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	#Chirality outliers	#Planarity outliers
2	O	0	1
All	All	0	10

The worst 5 of 46 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	96	HIS	C-N	10.77	1.46	1.32
1	C	97	THR	C-N	-7.60	1.22	1.33
1	B	522	THR	C-O	-7.44	1.14	1.24
1	A	98	PRO	CA-C	7.25	1.62	1.52
1	M	147	HIS	CA-C	7.04	1.62	1.52

The worst 5 of 90 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	147	HIS	N-CA-C	14.88	130.73	112.87
1	M	147	HIS	N-CA-C	14.27	130.13	113.01
1	B	147	HIS	N-CA-C	11.80	127.17	113.01
1	H	147	HIS	N-CA-C	11.43	126.73	113.01
1	C	147	HIS	N-CA-C	11.41	126.70	113.01

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	131	CYS	Peptide
1	B	97	THR	Mainchain
1	C	97	THR	Mainchain
1	D	131	CYS	Peptide
1	G	97	THR	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4491	0	4320	15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4490	0	4320	9	0
1	C	4491	0	4320	13	0
1	D	4490	0	4320	14	0
1	G	4491	0	4320	20	0
1	H	4490	0	4320	20	0
1	M	4491	0	4320	15	0
1	N	4490	0	4320	17	0
2	E	568	0	545	1	0
2	F	568	0	545	3	0
2	I	568	0	545	2	0
2	J	568	0	545	2	0
2	K	568	0	545	0	0
2	L	568	0	545	5	0
2	O	568	0	545	3	0
2	P	568	0	545	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	M	1	0	0	0	0
3	N	1	0	0	0	0
4	A	24	0	3	1	0
4	B	24	0	3	2	0
4	C	24	0	3	2	0
4	D	24	0	3	2	0
4	G	24	0	3	2	0
4	H	24	0	3	2	0
4	M	24	0	3	2	0
4	N	24	0	3	1	0
5	A	254	0	0	1	0
5	B	309	0	0	1	0
5	C	286	0	0	0	0
5	D	288	0	0	3	0
5	E	34	0	0	0	0
5	F	43	0	0	0	0
5	G	250	0	0	0	0
5	H	284	0	0	2	0
5	I	25	0	0	0	0
5	J	38	0	0	0	0
5	K	48	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	59	0	0	2	0
5	M	308	0	0	2	0
5	N	238	0	0	1	0
5	O	42	0	0	1	0
5	P	26	0	0	0	0
All	All	43200	0	38944	134	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:90:ASP:HB2	5:H:1055:HOH:O	1.76	0.84
2:O:90:VAL:HG12	2:O:93:ILE:HD12	1.61	0.82
1:N:145:GLY:HA3	1:N:148:GLY:O	1.83	0.76
1:A:114:MET:HE1	1:B:591:GLN:HE21	1.53	0.72
1:C:591:GLN:HE21	1:D:114:MET:HE1	1.55	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	571/573 (100%)	539 (94%)	31 (5%)	1 (0%)	43 51
1	B	571/573 (100%)	538 (94%)	32 (6%)	1 (0%)	43 51
1	C	571/573 (100%)	540 (95%)	30 (5%)	1 (0%)	43 51
1	D	571/573 (100%)	535 (94%)	34 (6%)	2 (0%)	30 34
1	G	571/573 (100%)	542 (95%)	27 (5%)	2 (0%)	30 34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	571/573 (100%)	541 (95%)	27 (5%)	3 (0%)	24	27
1	M	571/573 (100%)	539 (94%)	31 (5%)	1 (0%)	43	51
1	N	571/573 (100%)	539 (94%)	30 (5%)	2 (0%)	30	34
2	E	69/72 (96%)	69 (100%)	0	0	100	100
2	F	69/72 (96%)	68 (99%)	1 (1%)	0	100	100
2	I	69/72 (96%)	68 (99%)	1 (1%)	0	100	100
2	J	69/72 (96%)	69 (100%)	0	0	100	100
2	K	69/72 (96%)	69 (100%)	0	0	100	100
2	L	69/72 (96%)	69 (100%)	0	0	100	100
2	O	69/72 (96%)	68 (99%)	1 (1%)	0	100	100
2	P	69/72 (96%)	68 (99%)	1 (1%)	0	100	100
All	All	5120/5160 (99%)	4861 (95%)	246 (5%)	13 (0%)	36	42

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	145	GLY
1	N	145	GLY
1	H	557	GLY
1	H	145	GLY
1	A	557	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/464 (100%)	458 (99%)	6 (1%)	61	76
1	B	464/464 (100%)	458 (99%)	6 (1%)	61	76
1	C	464/464 (100%)	458 (99%)	6 (1%)	61	76
1	D	464/464 (100%)	455 (98%)	9 (2%)	50	66
1	G	464/464 (100%)	457 (98%)	7 (2%)	57	73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	464/464 (100%)	456 (98%)	8 (2%)	53	69
1	M	464/464 (100%)	454 (98%)	10 (2%)	45	61
1	N	464/464 (100%)	455 (98%)	9 (2%)	50	66
2	E	60/61 (98%)	59 (98%)	1 (2%)	53	69
2	F	60/61 (98%)	54 (90%)	6 (10%)	7	7
2	I	60/61 (98%)	57 (95%)	3 (5%)	22	28
2	J	60/61 (98%)	57 (95%)	3 (5%)	22	28
2	K	60/61 (98%)	57 (95%)	3 (5%)	22	28
2	L	60/61 (98%)	57 (95%)	3 (5%)	22	28
2	O	60/61 (98%)	57 (95%)	3 (5%)	22	28
2	P	60/61 (98%)	54 (90%)	6 (10%)	7	7
All	All	4192/4200 (100%)	4103 (98%)	89 (2%)	47	63

5 of 89 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	K	85	LYS
1	N	146	GLU
2	L	43	ASP
1	M	161	ILE
1	N	337	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 54 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	117	GLN
2	J	74	GLN
1	N	591	GLN
1	H	156	GLN
1	H	545	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PQQ	G	702	3	26,26,26	3.61	10 (38%)	34,40,40	2.15	12 (35%)
4	PQQ	B	702	3	26,26,26	3.42	10 (38%)	34,40,40	1.96	10 (29%)
4	PQQ	H	702	3	26,26,26	3.59	7 (26%)	34,40,40	1.97	12 (35%)
4	PQQ	M	702	3	26,26,26	3.66	7 (26%)	34,40,40	2.37	16 (47%)
4	PQQ	N	702	3	26,26,26	3.64	9 (34%)	34,40,40	2.32	13 (38%)
4	PQQ	D	702	3	26,26,26	3.35	8 (30%)	34,40,40	2.32	16 (47%)
4	PQQ	C	702	3	26,26,26	3.40	8 (30%)	34,40,40	2.00	13 (38%)
4	PQQ	A	702	3	26,26,26	3.64	10 (38%)	34,40,40	2.25	16 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PQQ	G	702	3	-	2/12/28/28	0/3/3/3
4	PQQ	B	702	3	-	4/12/28/28	0/3/3/3
4	PQQ	H	702	3	-	3/12/28/28	0/3/3/3
4	PQQ	M	702	3	-	2/12/28/28	0/3/3/3

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PQQ	N	702	3	-	3/12/28/28	0/3/3/3
4	PQQ	D	702	3	-	2/12/28/28	0/3/3/3
4	PQQ	C	702	3	-	3/12/28/28	0/3/3/3
4	PQQ	A	702	3	-	5/12/28/28	0/3/3/3

The worst 5 of 69 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	N	702	PQQ	C5-C4	-12.75	1.36	1.54
4	A	702	PQQ	C5-C4	-12.32	1.37	1.54
4	M	702	PQQ	C5-C4	-12.24	1.37	1.54
4	B	702	PQQ	C5-C4	-11.97	1.37	1.54
4	H	702	PQQ	C5-C4	-11.67	1.38	1.54

The worst 5 of 108 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	N	702	PQQ	O9B-C9X-O9A	-5.31	111.94	123.35
4	M	702	PQQ	C3-C2-C2X	-4.93	121.17	132.53
4	D	702	PQQ	O9B-C9X-O9A	-4.78	113.08	123.35
4	A	702	PQQ	C3-C2-C2X	-4.71	121.69	132.53
4	B	702	PQQ	C3-C2-C2X	-4.62	121.88	132.53

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	702	PQQ	C3-C2-C2X-O2A
4	B	702	PQQ	N1-C2-C2X-O2A
4	B	702	PQQ	N1-C2-C2X-O2B
4	B	702	PQQ	C3-C2-C2X-O2A
4	B	702	PQQ	C3-C2-C2X-O2B

There are no ring outliers.

8 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	702	PQQ	2	0
4	B	702	PQQ	2	0
4	H	702	PQQ	2	0


Continued on next page...


*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	M	702	PQQ	2	0
4	N	702	PQQ	1	0
4	D	702	PQQ	2	0
4	C	702	PQQ	2	0
4	A	702	PQQ	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	573/573 (100%)	-0.58	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	14, 22, 35, 67	0
1	B	573/573 (100%)	-0.66	2 (0%) <span style="border: 1px solid blue; padding: 2px;">90</span> <span style="border: 1px solid blue; padding: 2px;">88</span>	13, 20, 34, 54	0
1	C	573/573 (100%)	-0.69	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	13, 19, 31, 52	0
1	D	573/573 (100%)	-0.69	1 (0%) <span style="border: 1px solid blue; padding: 2px;">91</span> <span style="border: 1px solid blue; padding: 2px;">90</span>	13, 19, 33, 52	0
1	G	573/573 (100%)	-0.55	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	13, 23, 38, 62	0
1	H	573/573 (100%)	-0.66	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	14, 19, 34, 55	0
1	M	573/573 (100%)	-0.66	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	14, 20, 33, 53	0
1	N	573/573 (100%)	-0.49	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	14, 25, 40, 62	0
2	E	71/72 (98%)	-0.14	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	21, 32, 49, 66	0
2	F	71/72 (98%)	0.18	8 (11%) <span style="border: 1px solid red; padding: 2px;">10</span> <span style="border: 1px solid red; padding: 2px;">8</span>	16, 26, 78, 113	0
2	I	71/72 (98%)	0.20	2 (2%) <span style="border: 1px solid gray; padding: 2px;">55</span> <span style="border: 1px solid gray; padding: 2px;">52</span>	25, 37, 55, 74	0
2	J	71/72 (98%)	-0.24	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	18, 25, 48, 63	0
2	K	71/72 (98%)	-0.29	1 (1%) <span style="border: 1px solid blue; padding: 2px;">73</span> <span style="border: 1px solid blue; padding: 2px;">71</span>	17, 26, 45, 58	0
2	L	71/72 (98%)	-0.30	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	18, 28, 49, 68	0
2	O	71/72 (98%)	0.15	6 (8%) <span style="border: 1px solid red; padding: 2px;">16</span> <span style="border: 1px solid red; padding: 2px;">14</span>	20, 28, 80, 130	0
2	P	71/72 (98%)	0.29	3 (4%) <span style="border: 1px solid red; padding: 2px;">40</span> <span style="border: 1px solid red; padding: 2px;">37</span>	30, 39, 58, 72	0
All	All	5152/5160 (99%)	-0.56	23 (0%) <span style="border: 1px solid blue; padding: 2px;">88</span> <span style="border: 1px solid blue; padding: 2px;">87</span>	13, 21, 41, 130	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	O	90	VAL	6.2
2	O	93	ILE	5.9
2	F	93	ILE	5.4
2	O	91	GLU	4.4
2	F	82	LYS	4.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	M	701	1/1	0.94	0.15	58,58,58,58	0
3	CA	N	701	1/1	0.94	0.13	77,77,77,77	0
3	CA	B	701	1/1	0.95	0.21	61,61,61,61	0
3	CA	G	701	1/1	0.95	0.15	67,67,67,67	0
4	PQQ	A	702	24/24	0.95	0.06	18,23,26,31	0
4	PQQ	H	702	24/24	0.95	0.06	20,24,25,27	0
4	PQQ	N	702	24/24	0.95	0.06	20,24,27,28	0
3	CA	H	701	1/1	0.96	0.23	60,60,60,60	0
4	PQQ	C	702	24/24	0.96	0.05	16,22,27,27	0
4	PQQ	D	702	24/24	0.96	0.05	16,19,21,24	0
4	PQQ	G	702	24/24	0.96	0.05	17,23,25,27	0
3	CA	D	701	1/1	0.96	0.21	59,59,59,59	0
4	PQQ	M	702	24/24	0.96	0.06	16,21,23,28	0
3	CA	A	701	1/1	0.96	0.16	57,57,57,57	0
4	PQQ	B	702	24/24	0.97	0.05	16,19,20,23	0
3	CA	C	701	1/1	0.98	0.19	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around PQQ A 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PQQ H 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PQQ N 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PQQ C 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PQQ D 702:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around PQQ G 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.