



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:57 AM UTC

PDB ID : 3CFC / pdb\_00003cfc  
Title : High-resolution structure of blue fluorescent antibody EP2-19G2  
Authors : Debler, E.W.; Wilson, I.A.  
Deposited on : 2008-03-03  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	213	
2	L	219	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BLUE FLUORESCENT ANTIBODY EP2-19G2-IGG2B HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	213	1662	1041	278	334	9	0	11	0

- Molecule 2 is a protein called BLUE FLUORESCENT ANTIBODY EP2-19G2-KAPPA LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	219	1703	1061	286	348	8	0	2	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	H	1	6	3	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	187	Total 187	O 187	0	0
4	L	145	Total 145	O 145	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BLUE FLUORESCENT ANTIBODY EP2-19G2-IGG2B HEAVY CHAIN



- Molecule 2: BLUE FLUORESCENT ANTIBODY EP2-19G2-KAPPA LIGHT CHAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.89Å 87.89Å 114.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.76 – 1.70 28.76 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (28.76-1.70) 99.9 (28.76-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.3.0011	Depositor
R, $R_{free}$	0.191 , 0.231 0.261 , 0.294	Depositor DCC
$R_{free}$ test set	2518 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtrriage
Anisotropy	0.048	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 36.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3703	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	0.92	1/1701 (0.1%)	0.94	0/2319
2	L	0.79	1/1740 (0.1%)	0.86	0/2363
All	All	0.86	2/3441 (0.1%)	0.90	0/4682

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	82	MET	SD-CE	-6.91	1.62	1.79
2	L	96	PRO	N-CA	-5.00	1.41	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1662	0	1632	10	0
2	L	1703	0	1639	4	1
3	H	6	0	8	0	0
4	H	187	0	0	2	1
4	L	145	0	0	0	0
All	All	3703	0	3279	14	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:12:VAL:HG11	1:H:82(C):LEU:HD13	1.92	0.51
1:H:140:LEU:HB3	1:H:223:LEU:HD22	1.93	0.50
1:H:40:ILE:HG23	1:H:41:PRO:HD2	1.94	0.50
1:H:2:VAL:HA	1:H:25[A]:SER:O	2.10	0.49
1:H:71:ARG:NH1	4:H:451:HOH:O	2.25	0.49
4:H:448:HOH:O	2:L:96:PRO:HG3	2.13	0.48
2:L:39:LYS:HE2	2:L:81:GLU:O	2.13	0.48
1:H:134:THR:CG2	1:H:135:GLY:N	2.79	0.45
1:H:82:MET:HE1	1:H:109:VAL:HG21	1.99	0.44
2:L:163:TRP:CE2	2:L:175:MET:HE3	2.55	0.41
1:H:134:THR:HG22	1:H:135:GLY:N	2.35	0.41
2:L:185:GLU:HA	2:L:188:ARG:HD2	2.03	0.41
1:H:71:ARG:NE	1:H:73:ASN:OD1	2.51	0.40
1:H:41:PRO:C	1:H:43:LYS:H	2.29	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:1:ASP:O	4:H:413:HOH:O[8_665]	2.17	0.03

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	222/213 (104%)	217 (98%)	5 (2%)	0	100	100
2	L	219/219 (100%)	217 (99%)	2 (1%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	441/432 (102%)	434 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	195/184 (106%)	193 (99%)	2 (1%)	68	58
2	L	196/194 (101%)	194 (99%)	2 (1%)	68	58
All	All	391/378 (103%)	387 (99%)	4 (1%)	65	58

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	223	LEU
1	H	226	GLU
2	L	11	ASN
2	L	143	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	H	301	-	5,5,5	0.36	0	5,5,5	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	H	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	213/213 (100%)	2.16	106 (49%) <b>0</b> <b>0</b>	19, 40, 55, 64	11 (5%)
2	L	219/219 (100%)	1.80	77 (35%) <b>1</b> <b>1</b>	21, 42, 49, 59	2 (0%)
All	All	432/432 (100%)	1.98	183 (42%) <b>0</b> <b>0</b>	19, 41, 52, 64	13 (3%)

All (183) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	227	PRO	6.1
2	L	9	PHE	5.9
1	H	42	GLU	5.6
1	H	41	PRO	5.6
1	H	134	THR	5.4
2	L	40	PRO	5.0
2	L	7	ALA	5.0
1	H	223	LEU	4.8
1	H	43	LYS	4.7
1	H	40	ILE	4.7
1	H	166	LEU	4.6
2	L	210	ASN	4.6
2	L	209	PHE	4.5
2	L	41	GLY	4.5
2	L	15	LEU	4.5
1	H	103	TRP	4.3
2	L	69	THR	4.1
2	L	188	ARG	4.0
1	H	133	THR	3.8
1	H	45	LEU	3.8
1	H	222	LYS	3.7
1	H	109	VAL	3.7
1	H	169	SER	3.7
2	L	153	SER	3.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	L	150	ILE	3.6
1	H	168	SER	3.6
1	H	130	ASP	3.6
1	H	36	TRP	3.5
1	H	37	VAL	3.5
1	H	80	LEU	3.5
2	L	194	CYS	3.5
1	H	105	GLN	3.5
1	H	63	VAL	3.5
1	H	44	ARG	3.5
2	L	68	GLY	3.5
2	L	74	ARG	3.5
1	H	163	SER	3.4
1	H	49	ALA	3.3
2	L	181	LEU	3.3
1	H	171	VAL	3.3
2	L	8	ALA	3.3
1	H	82(C)	LEU	3.3
2	L	202	THR	3.2
1	H	69	ILE	3.2
1	H	78	LEU	3.2
1	H	177	LEU	3.2
1	H	48	VAL	3.2
2	L	190	ASN	3.1
1	H	102	TYR	3.1
2	L	214	CYS	3.1
1	H	226	GLU	3.1
1	H	68	THR	3.1
1	H	167	SER	3.1
2	L	122	SER	3.1
1	H	179	GLN	3.1
1	H	16	GLY	3.1
2	L	156	GLN	3.1
1	H	1	GLU	3.1
2	L	43	SER	3.1
1	H	206	THR	3.0
1	H	67	PHE	3.0
2	L	213	GLU	3.0
1	H	18	LEU	3.0
1	H	88	ALA	3.0
2	L	152	GLY	3.0
2	L	163	TRP	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	H	82(A)[A]	SER	3.0
2	L	10	SER	3.0
2	L	56	SER	3.0
2	L	157	ASN	3.0
1	H	205	VAL	2.9
1	H	219	VAL	2.9
1	H	90	TYR	2.9
2	L	26	THR	2.9
2	L	212	ASN	2.9
1	H	164	GLY	2.9
1	H	51	ILE	2.9
1	H	82	MET	2.9
1	H	17[A]	SER	2.9
1	H	12	VAL	2.8
1	H	135	GLY	2.8
1	H	20	LEU	2.8
1	H	89	LEU	2.8
1	H	25[A]	SER	2.8
1	H	4	LEU	2.7
1	H	5	VAL	2.7
2	L	21	ILE	2.7
2	L	189	HIS	2.7
2	L	191	SER	2.7
1	H	47	TRP	2.7
2	L	115	VAL	2.7
1	H	156	THR	2.6
1	H	74	VAL	2.6
1	H	79	TYR	2.6
2	L	145	ASN	2.6
1	H	138	VAL	2.6
1	H	93	ALA	2.6
2	L	126	THR	2.5
1	H	136	SER	2.5
2	L	130	ALA	2.5
2	L	42	GLN	2.5
2	L	148	TRP	2.5
1	H	24	ALA	2.5
1	H	111	VAL	2.5
2	L	197	THR	2.5
1	H	29	PHE	2.5
2	L	46	LEU	2.5
1	H	106	GLY	2.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	L	58	VAL	2.5
2	L	206	VAL	2.5
2	L	185	GLU	2.5
1	H	59	TYR	2.4
1	H	23	THR	2.4
1	H	107	THR	2.4
2	L	11	ASN	2.4
1	H	87	THR	2.4
1	H	144	VAL	2.4
2	L	144	ILE	2.4
2	L	128	GLY	2.4
1	H	9	GLY	2.3
2	L	27(A)	SER	2.3
2	L	187	GLU	2.3
1	H	153	THR	2.3
2	L	146	VAL	2.3
1	H	15	GLY	2.3
2	L	81	GLU	2.3
1	H	19	LYS	2.3
1	H	33	ILE	2.3
1	H	53[A]	SER	2.3
1	H	195	SER	2.3
1	H	221	LYS	2.3
1	H	218	THR	2.3
2	L	193	THR	2.3
2	L	105	GLU	2.3
2	L	27(E)	SER	2.3
1	H	92[A]	CYS	2.2
1	H	128	CYS	2.2
1	H	70	SER	2.2
1	H	112[A]	SER	2.2
2	L	208	SER	2.2
2	L	192	TYR	2.2
1	H	39	GLN	2.2
2	L	204	PRO	2.2
2	L	53	ASN	2.2
1	H	13	LYS	2.2
2	L	18	SER	2.2
1	H	11	LEU	2.2
2	L	129	GLY	2.2
2	L	71	PHE	2.2
1	H	110	THR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	L	44	PRO	2.2
1	H	84[A]	SER	2.2
1	H	208	CYS	2.2
2	L	45	GLN	2.2
1	H	174	PHE	2.2
1	H	21[A]	SER	2.2
2	L	70	ASP	2.1
2	L	143	ASP	2.1
2	L	211	ARG	2.1
2	L	104	LEU	2.1
1	H	7	SER	2.1
2	L	182	THR	2.1
2	L	184	ASP	2.1
2	L	39	LYS	2.1
2	L	54	LEU	2.1
1	H	82(B)	SER	2.1
1	H	85	GLU	2.1
1	H	2	VAL	2.1
1	H	152	VAL	2.1
2	L	13	VAL	2.1
1	H	22	CYS	2.1
1	H	3	LYS	2.1
1	H	64	LYS	2.1
1	H	91	TYR	2.1
2	L	205	ILE	2.1
1	H	54	GLY	2.0
1	H	81	GLN	2.0
2	L	1	ASP	2.0
2	L	203	SER	2.0
2	L	36	TYR	2.0
1	H	113[A]	SER	2.0
2	L	127	SER	2.0
1	H	193	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	H	301	6/6	0.81	0.17	33,40,43,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.