



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:11 PM UTC

PDB ID : 3CHT / pdb_00003cht
Title : Crystal Structure of Di-iron AurF with partially bound Ligand
Authors : Zhang, H.; Brunzelle, J.S.; Nair, S.K.
Deposited on : 2008-03-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	 7% 75% 12% • 10%
1	B	336	 13% 74% 14% • 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	4NB	A	502	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called p-Aminobenzoate N-Oxygenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	301	2404	1525	431	438	10	0	0	0
1	B	300	2388	1517	427	434	10	0	0	0

- Molecule 2 is MU-OXO-DIIRON (CCD ID: FEO) (formula: Fe₂O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	O		
2	A	1	3	2	1	0	0
2	B	1	3	2	1	0	0

- Molecule 3 is 4-NITROBENZOIC ACID (CCD ID: 4NB) (formula: C₇H₅NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	12	7	1	4	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	139	139	139	0	0
4	B	91	91	91	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: p-Aminobenzoate N-Oxygenase

Chain A: 



- Molecule 1: p-Aminobenzoate N-Oxygenase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.14Å 77.19Å 140.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.00 25.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.6 (25.00-2.00) 96.5 (25.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.256 0.204 , 0.257	Depositor DCC
R_{free} test set	2135 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtrriage
Anisotropy	0.465	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5040	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 4NB, FEO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2454	0.89	4/3338 (0.1%)
1	B	0.76	4/2438 (0.2%)	0.89	2/3318 (0.1%)
All	All	0.73	4/4892 (0.1%)	0.89	6/6656 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	247	GLU	C-N	13.83	1.52	1.34
1	B	247	GLU	C-O	10.13	1.35	1.24
1	B	246	ASN	CG-OD1	9.23	1.41	1.23
1	B	246	ASN	CG-ND2	7.58	1.49	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	247	GLU	O-C-N	6.66	129.18	122.12
1	A	52	ALA	N-CA-C	5.26	117.42	111.11
1	A	178	MET	CA-C-N	5.08	125.11	119.32
1	A	178	MET	C-N-CA	5.08	125.11	119.32
1	A	162	GLN	N-CA-C	5.00	120.87	109.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	246	ASN	OD1-CG-ND2	5.00	127.60	122.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	GLN	Peptide
1	B	178	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2404	0	2403	33	0
1	B	2388	0	2383	28	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	12	0	4	4	0
4	A	139	0	0	8	0
4	B	91	0	0	2	0
All	All	5040	0	4790	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ALA:HB1	4:A:637:HOH:O	1.16	1.25
3:A:502:4NB:H2	4:A:599:HOH:O	1.84	0.77
1:A:323:ASP:HB2	4:A:630:HOH:O	1.84	0.76
1:A:78:GLU:H	1:A:81:GLN:HE21	1.33	0.74
1:B:78:GLU:H	1:B:81:GLN:HE21	1.35	0.74
1:A:89:MET:HE3	1:A:272:LEU:HD11	1.72	0.70
1:A:168:TYR:CE1	1:A:172:ARG:HD2	2.27	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:267:GLN:HE21	1:B:267:GLN:HA	1.57	0.68
1:A:168:TYR:CE1	1:A:172:ARG:CD	2.78	0.65
1:A:225:ARG:HD2	4:A:567:HOH:O	1.95	0.65
1:A:225:ARG:NH2	4:A:639:HOH:O	2.00	0.63
1:A:68:PHE:HE2	1:A:272:LEU:HD12	1.65	0.60
1:B:48:LEU:HD13	1:B:145:LEU:HD22	1.84	0.59
1:B:78:GLU:H	1:B:81:GLN:NE2	2.01	0.57
1:A:203:LEU:HD22	3:A:502:4NB:H6	1.89	0.55
1:A:68:PHE:CE2	1:A:272:LEU:HD12	2.41	0.54
1:A:148:ASP:OD2	4:A:636:HOH:O	2.18	0.53
1:A:221:THR:O	1:A:225:ARG:HG3	2.09	0.53
1:A:168:TYR:CE1	1:A:172:ARG:HD3	2.44	0.52
1:B:32:VAL:HG11	1:B:127:LYS:HE2	1.92	0.52
1:A:168:TYR:CZ	1:A:172:ARG:HD2	2.45	0.51
1:A:37:ARG:NH1	4:A:561:HOH:O	2.43	0.51
1:B:181:GLN:NE2	1:B:184:ARG:HH11	2.09	0.51
1:A:78:GLU:H	1:A:81:GLN:NE2	2.06	0.50
1:A:51:TRP:CE2	1:A:215:MET:HG3	2.46	0.50
1:B:267:GLN:NE2	1:B:268:ASP:H	2.10	0.49
1:B:77:ALA:O	1:B:82:ARG:NH2	2.46	0.48
1:A:312:GLU:HG3	1:A:317:ASP:OD1	2.13	0.48
1:B:267:GLN:HE21	1:B:268:ASP:H	1.61	0.48
1:B:267:GLN:HE21	1:B:267:GLN:CA	2.25	0.48
1:A:93:TYR:OH	3:A:502:4NB:O1	2.31	0.48
1:B:171:LEU:HG	1:B:191:TRP:CG	2.49	0.48
1:B:179:PRO:HA	4:B:553:HOH:O	2.14	0.47
1:A:216:HIS:HE1	4:A:521:HOH:O	1.97	0.47
1:A:269:LEU:H	1:A:269:LEU:CD2	2.28	0.47
1:A:51:TRP:CD1	1:A:215:MET:HE3	2.49	0.47
1:A:238:VAL:HG12	1:A:314:LEU:HD21	1.97	0.47
1:B:60:ASP:O	1:B:216:HIS:HD2	1.98	0.46
1:A:236:GLU:CD	1:A:239:ARG:HH12	2.23	0.46
1:B:236:GLU:CD	1:B:239:ARG:HH12	2.23	0.46
1:B:179:PRO:HD3	4:B:541:HOH:O	2.16	0.45
1:B:252:ALA:O	1:B:256:CYS:HB2	2.16	0.45
1:A:181:GLN:NE2	1:A:184:ARG:HH11	2.15	0.45
1:B:89:MET:HG3	1:B:275:GLU:HG3	1.98	0.44
1:A:171:LEU:HG	1:A:191:TRP:CG	2.52	0.44
1:B:299:ARG:C	1:B:301:VAL:H	2.26	0.43
1:A:272:LEU:HD13	1:A:272:LEU:HA	1.93	0.43
1:B:171:LEU:HG	1:B:191:TRP:CD1	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:132:ALA:O	1:B:136:GLU:HG2	2.19	0.43
1:B:149:ARG:HH12	1:B:152:GLU:CD	2.28	0.42
1:A:97:VAL:O	1:A:101:GLU:HG2	2.20	0.42
1:A:203:LEU:HB2	3:A:502:4NB:O1	2.20	0.42
1:A:213:GLN:NE2	1:A:215:MET:HE1	2.35	0.42
1:B:122:ASP:HA	1:B:123:PRO:HD3	1.94	0.42
1:B:212:ILE:O	1:B:213:GLN:C	2.63	0.42
1:A:160:PRO:HA	1:A:161:PRO:HD3	1.86	0.41
1:B:111:LEU:HD23	1:B:111:LEU:HA	1.91	0.41
1:B:247:GLU:N	1:B:247:GLU:CD	2.79	0.41
1:B:108:ALA:HB2	1:B:192:GLY:HA3	2.02	0.41
1:B:277:ASN:HD22	1:B:277:ASN:HA	1.62	0.41
1:A:60:ASP:O	1:A:216:HIS:HD2	2.03	0.41
1:B:250:ARG:O	1:B:254:VAL:HG23	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/336 (88%)	289 (97%)	7 (2%)	1 (0%)	36	35
1	B	296/336 (88%)	286 (97%)	7 (2%)	3 (1%)	12	8
All	All	593/672 (88%)	575 (97%)	14 (2%)	4 (1%)	18	14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	52	ALA
1	B	179	PRO
1	A	162	GLN
1	B	300	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/281 (90%)	234 (92%)	19 (8%)	12	9
1	B	250/281 (89%)	239 (96%)	11 (4%)	25	24
All	All	503/562 (90%)	473 (94%)	30 (6%)	17	14

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	44	LYS
1	A	56	THR
1	A	57	LEU
1	A	70	GLU
1	A	171	LEU
1	A	173	ARG
1	A	177	ASP
1	A	189	LEU
1	A	191	TRP
1	A	215	MET
1	A	269	LEU
1	A	270	SER
1	A	272	LEU
1	A	281	ILE
1	A	312	GLU
1	A	317	ASP
1	A	327	ARG
1	A	331	SER
1	B	28	LEU
1	B	37	ARG
1	B	121	ASP
1	B	171	LEU
1	B	189	LEU
1	B	215	MET
1	B	247	GLU
1	B	267	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	277	ASN
1	B	281	ILE
1	B	317	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	83	GLN
1	A	181	GLN
1	A	213	GLN
1	A	216	HIS
1	A	248	GLN
1	B	81	GLN
1	B	181	GLN
1	B	213	GLN
1	B	216	HIS
1	B	246	ASN
1	B	267	GLN
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	4NB	A	502	-	12,12,12	1.13	2 (16%)	14,16,16	1.07	1 (7%)
2	FEO	B	501	1	0,2,2	-	-	-		
2	FEO	A	501	1	0,2,2	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4NB	A	502	-	-	3/6/8/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	4NB	C4-N	-2.29	1.39	1.45
3	A	502	4NB	O1-C	-2.10	1.24	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	4NB	C5-C4-N	2.06	121.13	119.34

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	4NB	C3-C4-N-O2'
3	A	502	4NB	C5-C4-N-O2'
3	A	502	4NB	O1-C-C1-C2

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	4NB	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/336 (89%)	0.92	23 (7%) 20 18	27, 40, 56, 65	0
1	B	300/336 (89%)	1.23	44 (14%) 6 5	29, 39, 55, 64	0
All	All	601/672 (89%)	1.07	67 (11%) 10 9	27, 40, 55, 65	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	GLN	4.0
1	B	318	ASP	4.0
1	A	52	ALA	3.4
1	B	268	ASP	3.4
1	A	72	PRO	3.3
1	A	284	ALA	3.2
1	A	318	ASP	3.1
1	B	176	ALA	3.0
1	B	70	GLU	2.9
1	A	163	PRO	2.9
1	B	43	ASN	2.9
1	B	184	ARG	2.9
1	A	214	PRO	2.9
1	B	247	GLU	2.9
1	B	329	ASP	2.8
1	B	137	SER	2.8
1	A	180	GLU	2.8
1	B	52	ALA	2.8
1	A	316	LEU	2.8
1	B	134	VAL	2.8
1	B	132	ALA	2.7
1	B	179	PRO	2.7
1	B	42	VAL	2.7
1	A	256	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	54	TYR	2.6
1	B	182	TRP	2.5
1	B	301	VAL	2.5
1	A	207	ALA	2.5
1	B	243	ALA	2.5
1	B	252	ALA	2.5
1	B	320	VAL	2.5
1	B	140	THR	2.5
1	B	255	ARG	2.5
1	B	161	PRO	2.4
1	A	319	ALA	2.4
1	B	272	LEU	2.4
1	A	327	ARG	2.4
1	B	116	VAL	2.4
1	B	121	ASP	2.4
1	B	330	TRP	2.4
1	B	189	LEU	2.4
1	B	299	ARG	2.4
1	B	24	GLY	2.4
1	B	157	SER	2.4
1	A	330	TRP	2.4
1	A	161	PRO	2.3
1	A	158	GLU	2.3
1	B	180	GLU	2.3
1	A	267	GLN	2.3
1	B	160	PRO	2.3
1	B	166	VAL	2.3
1	B	174	VAL	2.3
1	A	225	ARG	2.3
1	A	63	LEU	2.3
1	A	73	LEU	2.3
1	B	75	LEU	2.2
1	A	49	ASP	2.2
1	B	245	MET	2.2
1	B	45	ALA	2.2
1	B	286	GLU	2.2
1	B	41	VAL	2.1
1	B	79	PRO	2.1
1	B	248	GLN	2.1
1	B	319	ALA	2.1
1	B	64	GLU	2.1
1	A	218	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	208	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	4NB	A	502	12/12	0.67	0.18	62,64,65,65	0
2	FEO	B	501	3/3	0.99	0.13	30,30,30,30	0
2	FEO	A	501	3/3	1.00	0.12	25,25,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.