



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 07:58 AM UTC

PDB ID : 3CHX / pdb\_00003chx  
Title : Crystal structure of Methylosinus trichosporium OB3b particulate methane monooxygenase (pMMO)  
Authors : Hakemian, A.S.  
Deposited on : 2008-03-10  
Resolution : 3.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1270 (4.10-3.70)
Clashscore	190562	1034 (4.08-3.72)
Ramachandran outliers	187476	1251 (4.10-3.70)
Sidechain outliers	187428	1243 (4.10-3.70)
RSRZ outliers	180081	1269 (4.10-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	
1	E	392	
1	I	392	
2	B	252	
2	F	252	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	J	252	
3	C	256	
3	G	256	
3	K	256	
4	D	20	
4	H	20	
4	L	20	
5	M	26	
5	N	26	
5	O	26	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 18945 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PmoB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	362	2832	1831	478	518	5	0	0	0
1	E	362	2832	1831	478	518	5	0	0	0
1	I	362	2832	1831	478	518	5	0	0	0

- Molecule 2 is a protein called PmoA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	238	1925	1303	301	310	11	0	0	0
2	F	238	1925	1303	301	310	11	0	0	0
2	J	238	1925	1303	301	310	11	0	0	0

- Molecule 3 is a protein called PmoC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	159	1324	886	211	223	4	0	0	0
3	G	159	1324	886	211	223	4	0	0	0
3	K	159	1324	886	211	223	4	0	0	0

- Molecule 4 is a protein called 20-residue peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	20	100	60	20	20	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	20	Total	C	N	O	0	0	0
			100	60	20	20			
4	L	20	Total	C	N	O	0	0	0
			100	60	20	20			

- Molecule 5 is a protein called 26-residue peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	M	26	Total	C	N	O	0	0	0
			131	78	26	27			
5	N	26	Total	C	N	O	0	0	0
			131	78	26	27			
5	O	26	Total	C	N	O	0	0	0
			131	78	26	27			


- Molecule 6 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Cu	0	0
			2	2		
6	C	1	Total	Cu	0	0
			1	1		
6	E	2	Total	Cu	0	0
			2	2		
6	G	1	Total	Cu	0	0
			1	1		
6	I	2	Total	Cu	0	0
			2	2		
6	K	1	Total	Cu	0	0
			1	1		


### 3 Residue-property plots i


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: PmoB




#### • Molecule 1: PmoB






• Molecule 1: PmoB




• Molecule 2: PmoA







• Molecule 2: PmoA




• Molecule 2: PmoA






• Molecule 3: PmoC




• Molecule 3: PmoC



• Molecule 3: PmoC





- Molecule 5: 26-residue peptide

Chain O: 19% 81%

A horizontal bar chart for Chain O showing 19% green and 81% yellow.



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.81Å 184.07Å 203.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.00 – 3.90 38.00 – 3.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.00-3.90) 99.8 (38.00-3.90)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.84 (at 3.87Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.342 , 0.377 0.331 , 0.365	Depositor DCC
$R_{free}$ test set	3992 reflections (10.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	121.3	Xtrriage
Anisotropy	0.991	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 115.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.37$ , $\langle L^2 \rangle = 0.20$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	18945	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	141.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.56	0/2902	1.25	36/3944 (0.9%)
1	E	0.56	0/2902	1.25	36/3944 (0.9%)
1	I	0.56	0/2902	1.25	36/3944 (0.9%)
2	B	0.71	2/2002 (0.1%)	1.35	35/2740 (1.3%)
2	F	0.71	2/2002 (0.1%)	1.35	35/2740 (1.3%)
2	J	0.71	2/2002 (0.1%)	1.35	35/2740 (1.3%)
3	C	0.57	0/1371	1.13	16/1876 (0.9%)
3	G	0.57	0/1371	1.13	16/1876 (0.9%)
3	K	0.57	0/1371	1.13	16/1876 (0.9%)
All	All	0.62	6/18825 (0.0%)	1.26	261/25680 (1.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	197	SER	CA-C	6.50	1.57	1.53
2	B	197	SER	CA-C	6.49	1.57	1.53
2	F	197	SER	CA-C	6.37	1.57	1.53
2	J	197	SER	N-CA	5.99	1.50	1.46
2	F	197	SER	N-CA	5.91	1.49	1.46

The worst 5 of 261 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	53	TRP	N-CA-C	-17.32	90.10	113.30
2	B	53	TRP	N-CA-C	-17.28	90.14	113.30
2	F	53	TRP	N-CA-C	-17.25	90.19	113.30
2	F	197	SER	N-CA-C	15.18	121.23	108.78
2	J	197	SER	N-CA-C	15.17	121.22	108.78

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2832	0	2837	741	1
1	E	2832	0	2837	740	0
1	I	2832	0	2837	746	1
2	B	1925	0	1884	626	0
2	F	1925	0	1884	636	0
2	J	1925	0	1884	642	0
3	C	1324	0	1292	349	0
3	G	1324	0	1292	346	0
3	K	1324	0	1292	343	0
4	D	100	0	22	18	0
4	H	100	0	22	18	0
4	L	100	0	22	19	0
5	M	131	0	28	30	0
5	N	131	0	28	30	0
5	O	131	0	28	30	0
6	A	2	0	0	0	0
6	C	1	0	0	0	0
6	E	2	0	0	0	0
6	G	1	0	0	0	0
6	I	2	0	0	0	0
6	K	1	0	0	0	0
All	All	18945	0	18189	4765	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 128.

The worst 5 of 4765 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:114:GLU:HG3	1:E:280:THR:HB	1.25	1.19
1:E:309:ARG:HD2	1:E:381:ALA:HA	1.22	1.16
1:I:359:LEU:HD21	1:I:376:VAL:HG13	1.17	1.15

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359:LEU:HD21	1:A:376:VAL:HG13	1.17	1.15
1:A:309:ARG:HD2	1:A:381:ALA:HA	1.22	1.14

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:LYS:NZ	1:I:230:LYS:NZ[3_555]	1.64	0.56

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/392 (90%)	206 (58%)	86 (24%)	62 (18%)	0	2
1	E	354/392 (90%)	206 (58%)	85 (24%)	63 (18%)	0	2
1	I	354/392 (90%)	206 (58%)	86 (24%)	62 (18%)	0	2
2	B	236/252 (94%)	87 (37%)	74 (31%)	75 (32%)	0	0
2	F	236/252 (94%)	87 (37%)	74 (31%)	75 (32%)	0	0
2	J	236/252 (94%)	87 (37%)	74 (31%)	75 (32%)	0	0
3	C	157/256 (61%)	72 (46%)	51 (32%)	34 (22%)	0	1
3	G	157/256 (61%)	72 (46%)	51 (32%)	34 (22%)	0	1
3	K	157/256 (61%)	72 (46%)	51 (32%)	34 (22%)	0	1
All	All	2241/2700 (83%)	1095 (49%)	632 (28%)	514 (23%)	0	1

5 of 514 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42	GLU
1	A	48	PHE

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	68	VAL
1	A	81	PHE
1	A	140	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/322 (92%)	254 (85%)	44 (15%)	3	16
1	E	298/322 (92%)	254 (85%)	44 (15%)	3	16
1	I	298/322 (92%)	254 (85%)	44 (15%)	3	16
2	B	197/207 (95%)	147 (75%)	50 (25%)	0	4
2	F	197/207 (95%)	147 (75%)	50 (25%)	0	4
2	J	197/207 (95%)	147 (75%)	50 (25%)	0	4
3	C	135/208 (65%)	114 (84%)	21 (16%)	2	15
3	G	135/208 (65%)	114 (84%)	21 (16%)	2	15
3	K	135/208 (65%)	114 (84%)	21 (16%)	2	15
All	All	1890/2211 (86%)	1545 (82%)	345 (18%)	2	11

5 of 345 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	42	GLU
2	J	106	TRP
1	I	116	PHE
1	I	295	VAL
2	J	188	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	398	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	G	32	ASN
3	K	91	ASN
2	J	112	ASN
2	F	14	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	362/392 (92%)	0.43	25 (6%) 23 20	145, 145, 145, 145	0
1	E	362/392 (92%)	0.34	12 (3%) 49 34	145, 145, 145, 145	0
1	I	362/392 (92%)	0.45	18 (4%) 34 26	145, 145, 145, 145	0
2	B	238/252 (94%)	0.58	18 (7%) 20 19	145, 145, 145, 145	0
2	F	238/252 (94%)	0.73	24 (10%) 12 15	145, 145, 145, 145	0
2	J	238/252 (94%)	0.77	25 (10%) 11 14	145, 145, 145, 145	0
3	C	159/256 (62%)	0.64	13 (8%) 17 18	145, 145, 145, 145	0
3	G	159/256 (62%)	0.69	14 (8%) 15 17	145, 145, 145, 145	0
3	K	159/256 (62%)	0.62	14 (8%) 15 17	145, 145, 145, 145	0
4	D	0/20	-	-	-	-
4	H	0/20	-	-	-	-
4	L	0/20	-	-	-	-
5	M	0/26	-	-	-	-
5	N	0/26	-	-	-	-
5	O	0/26	-	-	-	-
All	All	2277/2838 (80%)	0.55	163 (7%) 21 20	145, 145, 145, 145	0

The worst 5 of 163 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	139	ASP	9.0
2	B	199	PRO	7.6
1	I	82	SER	7.0
1	I	73	VAL	6.5
2	F	197	SER	6.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	CU	E	722	1/1	0.89	0.06	127,127,127,127	0
6	CU	I	732	1/1	0.94	0.05	127,127,127,127	0
6	CU	I	731	1/1	0.95	0.05	119,119,119,119	0
6	CU	E	721	1/1	0.95	0.05	119,119,119,119	0
6	CU	G	662	1/1	0.97	0.16	118,118,118,118	0
6	CU	K	663	1/1	0.97	0.04	118,118,118,118	0
6	CU	A	712	1/1	0.98	0.04	127,127,127,127	0
6	CU	C	661	1/1	0.98	0.03	118,118,118,118	0
6	CU	A	711	1/1	0.99	0.05	119,119,119,119	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.