



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:45 PM UTC

PDB ID : 6CH0 / pdb_00006ch0
Title : Structure of the Quorum Quenching lactonase from Alicyclobacillus acidoterrestris bound to a glycerol molecule
Authors : Bergonzi, C.; Schwab, M.; Naik, T.; Daude, D.; Chabriere, E.; Elias, M.
Deposited on : 2018-02-21
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	282	3% 89% 8% .
1	F	282	91% 6% .
1	I	282	3% 90% 6% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	275	Total	C	N	O	S	0	2	0
			2223	1417	380	413	13			
1	F	275	Total	C	N	O	S	0	5	0
			2254	1434	388	420	12			
1	I	275	Total	C	N	O	S	0	8	0
			2278	1447	388	430	13			

- Molecule 2 is COBALT (II) ION (CCD ID: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	2	Total	Co	0	0
			2	2		
2	F	2	Total	Co	0	0
			2	2		
2	I	2	Total	Co	0	0
			2	2		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			13	8	5		
4	F	1	Total	C	O	0	0
			13	8	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	F	1	Total	C	O	0	0
			13	8	5		
4	I	1	Total	C	O	0	0
			13	8	5		
4	I	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	I	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	49	Total	O	0	0
			49	49		
6	F	50	Total	O	0	0
			50	50		
6	I	28	Total	O	0	0
			28	28		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase

Chain C: 





- Molecule 1: Beta-lactamase

Chain F: 



- Molecule 1: Beta-lactamase

Chain I: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	110.84Å 113.50Å 79.10Å 90.00° 108.91° 90.00°	Depositor
Resolution (Å)	74.83 – 2.15 74.83 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.4 (74.83-2.15) 99.4 (74.83-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.185 , 0.229 (Not available) , 0.230	Depositor DCC
R_{free} test set	2502 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 25.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6979	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CO, GOL, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.52	0/2284	0.74	0/3096
1	F	0.50	0/2316	0.75	0/3139
1	I	0.48	0/2339	0.73	0/3169
All	All	0.50	0/6939	0.74	0/9404

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2223	0	2141	20	0
1	F	2254	0	2161	12	0
1	I	2278	0	2174	17	0
2	C	2	0	0	0	0
2	F	2	0	0	0	0
2	I	2	0	0	0	0
3	C	8	0	12	0	0
4	C	13	0	18	1	0
4	F	26	0	36	0	0
4	I	26	0	36	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	6	0	6	0	0
5	F	6	0	6	1	0
5	I	6	0	6	0	0
6	C	49	0	0	1	0
6	F	50	0	0	0	0
6	I	28	0	0	0	0
All	All	6979	0	6596	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:273[A]:ARG:CG	1:I:273[A]:ARG:HH11	1.79	0.96
1:C:73[B]:MET:HE2	1:C:73[B]:MET:HA	1.46	0.94
1:I:273[A]:ARG:HH11	1:I:273[A]:ARG:HG3	1.35	0.92
1:C:29:ALA:HB2	1:C:236:GLY:HA2	1.79	0.64
1:C:73[B]:MET:HA	1:C:73[B]:MET:CE	2.23	0.63
1:I:273[A]:ARG:HG3	1:I:273[A]:ARG:NH1	2.11	0.62
1:F:69:ASN:HD22	1:F:71:ASP:H	1.49	0.59
1:C:238:ILE:HA	6:C:408:HOH:O	2.03	0.59
1:I:107:ARG:HG3	1:I:108:PRO:HD2	1.84	0.58
1:I:273[A]:ARG:HH11	1:I:273[A]:ARG:HG2	1.66	0.57
1:I:9:GLN:HA	1:I:9:GLN:OE1	2.03	0.56
1:C:258:THR:HG21	4:C:305:PG4:H21	1.87	0.56
1:I:198:HIS:CD2	1:I:220:ASP:HB3	2.41	0.55
1:C:69:ASN:HD22	1:C:71:ASP:H	1.55	0.55
1:I:29:ALA:HB2	1:I:236:GLY:HA2	1.89	0.54
1:I:69:ASN:HD22	1:I:71:ASP:H	1.54	0.54
1:I:273[A]:ARG:CG	1:I:273[A]:ARG:NH1	2.51	0.52
1:I:9:GLN:OE1	1:I:9:GLN:CA	2.58	0.52
1:F:23:ASP:HB3	1:F:26:TRP:CD1	2.48	0.48
1:C:29:ALA:CB	1:C:236:GLY:HA2	2.45	0.46
1:F:21:ARG:HA	1:F:46:ILE:O	2.16	0.46
1:F:29:ALA:HB2	1:F:236:GLY:HA2	1.97	0.46
1:C:9:GLN:HA	1:C:9:GLN:OE1	2.17	0.45
1:F:108:PRO:CG	1:F:129:MET:HE3	2.47	0.45
1:I:93:GLU:HG2	1:I:96:TYR:CE2	2.51	0.44
1:C:73[B]:MET:HE1	1:C:84:GLN:HG2	1.99	0.44
1:I:190:LYS:NZ	4:I:303:PG4:H72	2.32	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:198:HIS:CD2	1:C:220:ASP:HB3	2.52	0.44
1:F:69:ASN:ND2	1:F:71:ASP:H	2.14	0.44
1:I:21:ARG:HA	1:I:46:ILE:O	2.17	0.44
1:I:11:LYS:HD3	1:I:281:TYR:CE1	2.53	0.43
1:I:72:SER:HB3	1:I:79:TRP:CE2	2.53	0.43
1:C:267:ASP:OD1	1:C:270:GLN:N	2.46	0.43
1:F:115:ILE:HD11	1:F:185:LEU:CD2	2.49	0.43
1:F:122:ASP:OD2	5:F:305:GOL:H2	2.18	0.43
1:C:73[B]:MET:HE2	1:C:73[B]:MET:CA	2.32	0.43
1:F:108:PRO:HG3	1:F:129:MET:HE3	2.01	0.42
1:F:72:SER:HB3	1:F:79:TRP:CE2	2.55	0.42
1:C:115:ILE:HD11	1:C:185:LEU:HD21	2.02	0.42
1:F:115:ILE:HD11	1:F:185:LEU:HD21	2.02	0.42
1:C:73[B]:MET:CE	1:C:84:GLN:HG2	2.49	0.41
1:C:22:MET:HE2	1:C:48:PHE:CE2	2.54	0.41
1:C:72:SER:HB3	1:C:79:TRP:CE2	2.56	0.41
1:I:29:ALA:CB	1:I:236:GLY:HA2	2.51	0.41
1:C:73[B]:MET:CE	1:C:73[B]:MET:CA	2.97	0.41
1:C:12:LEU:HD23	1:C:12:LEU:C	2.46	0.41
1:C:115:ILE:HD11	1:C:185:LEU:CD2	2.51	0.40
1:F:198:HIS:CD2	1:F:220:ASP:HB3	2.57	0.40
1:C:69:ASN:ND2	1:C:71:ASP:H	2.17	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	275/282 (98%)	266 (97%)	9 (3%)	0	100	100
1	F	278/282 (99%)	270 (97%)	8 (3%)	0	100	100
1	I	281/282 (100%)	269 (96%)	12 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	834/846 (99%)	805 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	236/239 (99%)	233 (99%)	3 (1%)	61	68
1	F	239/239 (100%)	238 (100%)	1 (0%)	84	89
1	I	242/239 (101%)	234 (97%)	8 (3%)	33	34
All	All	717/717 (100%)	705 (98%)	12 (2%)	55	60

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	132	ASN
1	C	155	GLU
1	C	168	LYS
1	F	36	ILE
1	I	9	GLN
1	I	45	PHE
1	I	76	GLN
1	I	93	GLU
1	I	107	ARG
1	I	273[A]	ARG
1	I	273[B]	ARG
1	I	283	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	25	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	31	HIS
1	C	38	ASN
1	C	69	ASN
1	C	98	HIS
1	F	17	ASN
1	F	31	HIS
1	F	38	ASN
1	F	69	ASN
1	F	132	ASN
1	I	25	ASN
1	I	31	HIS
1	I	38	ASN
1	I	69	ASN
1	I	103	GLN
1	I	132	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 6 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	I	305	2	5,5,5	0.35	0	5,5,5	0.30	0
4	PG4	I	303	-	12,12,12	0.55	0	11,11,11	0.29	0
5	GOL	F	305	2	5,5,5	0.34	0	5,5,5	0.40	0
4	PG4	F	304	-	12,12,12	0.50	0	11,11,11	0.33	0
5	GOL	C	306	2	5,5,5	0.50	0	5,5,5	0.58	0
4	PG4	I	304	-	12,12,12	0.50	0	11,11,11	0.22	0
3	EDO	C	303	-	3,3,3	0.53	0	2,2,2	0.08	0
4	PG4	F	303	-	12,12,12	0.55	0	11,11,11	0.11	0
4	PG4	C	305	-	12,12,12	0.50	0	11,11,11	0.28	0
3	EDO	C	304	-	3,3,3	0.45	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	I	305	2	-	2/4/4/4	-
4	PG4	I	303	-	-	6/10/10/10	-
5	GOL	F	305	2	-	2/4/4/4	-
4	PG4	F	304	-	-	5/10/10/10	-
5	GOL	C	306	2	-	0/4/4/4	-
4	PG4	I	304	-	-	4/10/10/10	-
3	EDO	C	303	-	-	1/1/1/1	-
4	PG4	F	303	-	-	1/10/10/10	-
4	PG4	C	305	-	-	5/10/10/10	-
3	EDO	C	304	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	305	GOL	O1-C1-C2-C3
5	I	305	GOL	C1-C2-C3-O3
4	I	304	PG4	O3-C5-C6-O4
4	C	305	PG4	O2-C3-C4-O3
4	I	304	PG4	O2-C3-C4-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	C	305	PG4	O1-C1-C2-O2
4	I	303	PG4	O4-C7-C8-O5
5	F	305	GOL	O1-C1-C2-O2
5	I	305	GOL	O2-C2-C3-O3
4	F	303	PG4	O2-C3-C4-O3
4	I	303	PG4	O1-C1-C2-O2
3	C	303	EDO	O1-C1-C2-O2
4	I	304	PG4	C3-C4-O3-C5
4	F	304	PG4	C4-C3-O2-C2
4	I	303	PG4	C4-C3-O2-C2
4	F	304	PG4	O4-C7-C8-O5
4	I	303	PG4	C3-C4-O3-C5
3	C	304	EDO	O1-C1-C2-O2
4	C	305	PG4	O3-C5-C6-O4
4	F	304	PG4	C5-C6-O4-C7
4	C	305	PG4	C8-C7-O4-C6
4	F	304	PG4	O1-C1-C2-O2
4	I	304	PG4	O1-C1-C2-O2
4	I	303	PG4	O2-C3-C4-O3
4	F	304	PG4	O3-C5-C6-O4
4	C	305	PG4	C1-C2-O2-C3
4	I	303	PG4	C6-C5-O3-C4

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	I	303	PG4	1	0
5	F	305	GOL	1	0
4	C	305	PG4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	275/282 (97%)	-0.03	3 (1%) 78 80	19, 38, 65, 84	2 (0%)
1	F	275/282 (97%)	0.02	1 (0%) 88 90	22, 40, 67, 89	5 (1%)
1	I	275/282 (97%)	0.33	8 (2%) 53 57	20, 47, 77, 106	8 (2%)
All	All	825/846 (97%)	0.11	12 (1%) 72 76	19, 42, 70, 106	15 (1%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	273[A]	ARG	3.1
1	F	237	ILE	3.0
1	C	236	GLY	2.9
1	I	110[A]	ASP	2.7
1	C	237	ILE	2.7
1	I	184	VAL	2.4
1	I	107	ARG	2.4
1	I	210	PRO	2.3
1	C	36	ILE	2.3
1	I	236	GLY	2.1
1	I	185	LEU	2.1
1	I	106	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PG4	I	303	13/13	0.82	0.21	56,59,64,75	13
3	EDO	C	303	4/4	0.84	0.17	38,40,47,52	0
4	PG4	I	304	13/13	0.85	0.16	45,57,67,69	6
3	EDO	C	304	4/4	0.86	0.18	52,55,56,59	0
4	PG4	F	303	13/13	0.87	0.14	49,63,70,70	0
4	PG4	F	304	13/13	0.87	0.15	42,52,76,76	6
4	PG4	C	305	13/13	0.88	0.14	59,66,74,77	0
5	GOL	C	306	6/6	0.92	0.13	35,50,55,56	0
5	GOL	F	305	6/6	0.93	0.10	54,61,62,71	0
5	GOL	I	305	6/6	0.95	0.10	47,61,63,67	0
2	CO	C	301	1/1	0.99	0.02	31,31,31,31	0
2	CO	I	301	1/1	0.99	0.02	37,37,37,37	0
2	CO	C	302	1/1	1.00	0.01	30,30,30,30	0
2	CO	I	302	1/1	1.00	0.02	36,36,36,36	0
2	CO	F	301	1/1	1.00	0.01	36,36,36,36	0
2	CO	F	302	1/1	1.00	0.01	35,35,35,35	0

6.5 Other polymers

There are no such residues in this entry.