



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:43 AM UTC

PDB ID : 6CH4 / pdb_00006ch4
Title : Aminoglycoside Phosphotransferase (2'')-Ia S376N mutant in complex with GMPPNP and Magnesium
Authors : Caldwell, S.J.; Berghuis, A.M.
Deposited on : 2018-02-21
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	3% 88% 9% ..
1	B	305	3% 88% 9% .
1	C	305	2% 87% 10% ..
1	D	305	4% 86% 6% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10345 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional AAC/APH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2448	1561	381	496	10			
1	B	297	Total	C	N	O	S	0	1	0
			2462	1570	383	499	10			
1	C	297	Total	C	N	O	S	0	1	0
			2451	1563	383	495	10			
1	D	282	Total	C	N	O	S	0	1	0
			2271	1451	353	457	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	376	ASN	SER	engineered mutation	UNP P0A0C1
B	376	ASN	SER	engineered mutation	UNP P0A0C1
C	376	ASN	SER	engineered mutation	UNP P0A0C1
D	376	ASN	SER	engineered mutation	UNP P0A0C1

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (CCD ID: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	B	1	Total	C	N	O	P	0	0
			28	10	6	10	2		
2	C	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	D	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		
3	C	2	Total	Mg	0	0
			2	2		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	154	Total	O	0	0
			154	154		
5	B	184	Total	O	0	0
			184	184		
5	C	149	Total	O	0	0
			149	149		
5	D	92	Total	O	0	0
			92	92		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Bifunctional AAC/APH




• Molecule 1: Bifunctional AAC/APH




• Molecule 1: Bifunctional AAC/APH



• Molecule 1: Bifunctional AAC/APH





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.43Å 99.91Å 93.20Å 90.00° 105.17° 90.00°	Depositor
Resolution (Å)	58.89 – 2.30 58.89 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (58.89-2.30) 100.0 (58.89-2.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.170 , 0.210 0.176 , 0.212	Depositor DCC
R_{free} test set	3596 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10345	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.43	12/2492 (0.5%)	1.19	6/3362 (0.2%)
1	B	1.40	8/2509 (0.3%)	1.16	7/3382 (0.2%)
1	C	1.35	7/2498 (0.3%)	1.19	5/3369 (0.1%)
1	D	1.23	4/2312 (0.2%)	1.12	2/3125 (0.1%)
All	All	1.36	31/9811 (0.3%)	1.17	20/13238 (0.2%)

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	236	GLY	CA-C	7.81	1.61	1.51
1	A	235	LYS	CA-C	7.41	1.62	1.53
1	B	395	GLY	C-O	-6.68	1.16	1.24
1	A	353	SER	CA-C	6.24	1.60	1.52
1	C	445	GLU	C-O	-6.16	1.16	1.24
1	A	236	GLY	C-O	6.16	1.32	1.23
1	A	253	THR	C-O	-6.12	1.16	1.23
1	A	397	SER	C-O	-6.09	1.16	1.23
1	C	261	GLU	C-O	-6.05	1.16	1.24
1	B	432	ILE	C-O	-5.92	1.17	1.23
1	A	404	CYS	CA-CB	5.82	1.62	1.53
1	C	448	TYR	C-O	-5.82	1.18	1.24
1	C	257	ILE	C-O	-5.79	1.18	1.24
1	B	257	ILE	N-CA	-5.78	1.41	1.46
1	A	257	ILE	C-O	-5.74	1.18	1.24
1	C	428	MET	C-O	-5.69	1.17	1.24
1	A	235	LYS	N-CA	5.62	1.52	1.46
1	B	208	ILE	C-O	-5.46	1.18	1.24
1	A	432	ILE	C-O	-5.35	1.18	1.23
1	C	235	LYS	CA-C	5.34	1.60	1.53
1	D	418	GLY	N-CA	5.33	1.51	1.45
1	C	404	CYS	CA-CB	5.18	1.61	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	236	GLY	N-CA	5.16	1.52	1.44
1	D	261	GLU	C-O	-5.14	1.17	1.24
1	D	445	GLU	C-O	-5.12	1.18	1.24
1	B	418	GLY	N-CA	5.10	1.50	1.45
1	A	185	ASN	CA-C	-5.07	1.46	1.52
1	D	185	ASN	CA-C	-5.04	1.46	1.52
1	A	261	GLU	C-O	-5.03	1.17	1.24
1	A	436	LYS	C-O	-5.01	1.17	1.24
1	B	251	LEU	C-O	-5.01	1.17	1.23

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	404	CYS	CA-CB-SG	-6.92	98.47	114.40
1	A	404	CYS	CA-CB-SG	-6.53	99.38	114.40
1	B	236	GLY	CA-C-O	6.35	128.38	121.90
1	C	198	ASP	N-CA-C	6.33	118.71	111.11
1	A	198	ASP	N-CA-C	6.17	118.52	111.11
1	A	198	ASP	CB-CA-C	-5.89	101.38	110.81
1	B	229	PHE	CA-C-O	-5.86	114.45	121.89
1	A	463	GLU	CA-CB-CG	5.63	125.37	114.10
1	C	198	ASP	CB-CA-C	-5.58	101.88	110.81
1	A	339	ILE	N-CA-C	5.56	118.33	112.83
1	B	478	LYS	N-CA-C	-5.55	105.31	111.36
1	B	236	GLY	N-CA-C	5.44	120.38	112.60
1	C	377	CYS	N-CA-C	5.36	117.55	111.11
1	B	417	ILE	N-CA-C	5.33	115.54	110.53
1	B	450	ILE	N-CA-C	-5.32	105.53	110.53
1	C	475	ARG	NH1-CZ-NH2	-5.27	112.45	119.30
1	B	198	ASP	CA-C-O	-5.22	114.20	120.10
1	A	478	LYS	N-CA-C	-5.11	105.79	111.36
1	D	185	ASN	O-C-N	5.08	127.31	122.07
1	D	377	CYS	N-CA-C	5.03	117.14	111.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2448	0	2348	10	1
1	B	2462	0	2364	9	0
1	C	2451	0	2359	14	0
1	D	2271	0	2140	12	0
2	A	32	0	13	0	0
2	B	28	0	12	0	0
2	C	32	0	13	0	0
2	D	32	0	13	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	154	0	0	1	0
5	B	184	0	0	2	0
5	C	149	0	0	2	1
5	D	92	0	0	2	0
All	All	10345	0	9262	39	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:359:ASN:OD1	5:D:1050:HOH:O	2.00	0.79
1:B:359:ASN:OD1	5:B:1050:HOH:O	2.07	0.71
1:D:189:MET:SD	1:D:227:THR:HG21	2.31	0.70
1:C:198:ASP:OD1	5:C:1122:HOH:O	2.11	0.68
1:B:189:MET:SD	1:B:227:THR:HG21	2.33	0.68
1:C:243:ILE:HD13	1:C:397:SER:HB3	1.77	0.67
1:C:191:TYR:HH	1:D:196:TYR:HH	1.24	0.67
1:A:189:MET:SD	1:A:227:THR:HG21	2.38	0.63
1:C:184:THR:HB	1:D:185:ASN:OD1	2.00	0.62
1:D:186:VAL:HG13	1:D:205:ILE:HG23	1.82	0.62
1:A:184:THR:HB	1:B:185:ASN:ND2	2.17	0.60
1:D:326:GLN:NE2	5:D:1013:HOH:O	2.32	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:186:VAL:HG13	1:C:205:ILE:HG23	1.85	0.59
1:B:186:VAL:HG13	1:B:205:ILE:HG23	1.89	0.55
1:C:189:MET:HG2	1:C:217:TYR:CE1	2.42	0.54
1:C:191:TYR:OH	1:D:196:TYR:OH	2.04	0.52
1:B:189:MET:HG2	1:B:217:TYR:CE1	2.45	0.51
1:C:186:VAL:CG1	1:C:205:ILE:HG23	2.41	0.51
1:A:189:MET:HG2	1:A:217:TYR:CE1	2.45	0.51
1:B:186:VAL:CG1	1:B:205:ILE:HG23	2.42	0.49
1:B:298:LEU:HA	5:B:1268:HOH:O	2.12	0.49
1:D:186:VAL:CG1	1:D:205:ILE:HG23	2.42	0.48
1:D:189:MET:HG2	1:D:217:TYR:CE1	2.49	0.47
1:C:328:VAL:HG11	1:C:358:LEU:HD22	1.96	0.47
1:C:189:MET:SD	1:C:227:THR:HG21	2.55	0.46
1:A:186:VAL:HG13	1:A:205:ILE:HG23	1.97	0.46
5:A:1062:HOH:O	1:D:198:ASP:CB	2.62	0.46
1:C:446:GLU:OE2	5:C:1065:HOH:O	2.21	0.46
1:A:182:ASN:N	1:B:181:ASP:OD2	2.49	0.46
1:A:186:VAL:CG1	1:A:205:ILE:HG23	2.46	0.45
1:A:243:ILE:HD13	1:A:397:SER:HB2	1.99	0.45
1:A:185:ASN:ND2	1:B:179:TYR:OH	2.50	0.45
1:C:307[B]:ARG:HH12	1:C:431:ASN:HB2	1.83	0.44
1:A:317:ILE:HD12	1:A:317:ILE:HA	1.79	0.44
1:C:229:PHE:O	1:C:230:SER:C	2.61	0.42
1:A:407:ILE:HD13	1:A:407:ILE:HG21	1.91	0.41
1:C:471:GLU:HG3	1:C:475:ARG:HD2	2.02	0.41
1:D:420:ASN:HD22	1:D:420:ASN:HA	1.55	0.41
1:D:307[B]:ARG:NH1	1:D:431:ASN:HB2	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:GLU:OE1	5:C:1080:HOH:O[2_747]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/305 (97%)	285 (96%)	11 (4%)	0	100	100
1	B	294/305 (96%)	283 (96%)	11 (4%)	0	100	100
1	C	296/305 (97%)	284 (96%)	12 (4%)	0	100	100
1	D	277/305 (91%)	267 (96%)	8 (3%)	2 (1%)	18	23
All	All	1163/1220 (95%)	1119 (96%)	42 (4%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	198	ASP
1	D	400	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/281 (96%)	263 (97%)	7 (3%)	40	59
1	B	273/281 (97%)	267 (98%)	6 (2%)	45	65
1	C	271/281 (96%)	265 (98%)	6 (2%)	45	65
1	D	243/281 (86%)	241 (99%)	2 (1%)	73	86
All	All	1057/1124 (94%)	1036 (98%)	21 (2%)	48	67

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	THR
1	A	212	TYR
1	A	221	ASN
1	A	267	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	284	PRO
1	A	385	ASN
1	A	478	LYS
1	B	212	TYR
1	B	221	ASN
1	B	230	SER
1	B	245	ASN
1	B	285	GLU
1	B	420	ASN
1	C	221	ASN
1	C	232	ASN
1	C	267	ASP
1	C	297	LEU
1	C	300	ARG
1	C	397	SER
1	D	221	ASN
1	D	245	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	378	ASN
1	A	386	ASN
1	A	467	ASN
1	B	185	ASN
1	B	359	ASN
1	B	378	ASN
1	B	467	ASN
1	C	327	ASN
1	C	431	ASN
1	D	295	GLN
1	D	296	ASN
1	D	326	GLN
1	D	378	ASN
1	D	420	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GNP	B	500	3	29,30,34	2.20	3 (10%)	42,47,54	0.93	3 (7%)
2	GNP	C	500	3	34,34,34	2.68	5 (14%)	47,54,54	2.07	8 (17%)
2	GNP	D	500	3	34,34,34	4.18	7 (20%)	47,54,54	1.57	8 (17%)
2	GNP	A	500	3	34,34,34	3.48	5 (14%)	47,54,54	1.72	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GNP	B	500	3	-	1/13/32/38	0/3/3/3
2	GNP	C	500	3	-	4/18/38/38	0/3/3/3
2	GNP	D	500	3	-	1/18/38/38	0/3/3/3
2	GNP	A	500	3	-	4/18/38/38	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	500	GNP	PB-O1B	17.85	1.73	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	GNP	PB-O1B	15.18	1.69	1.46
2	D	500	GNP	PG-O1G	12.37	1.64	1.46
2	A	500	GNP	PG-O1G	11.28	1.63	1.46
2	C	500	GNP	PB-O1B	9.82	1.61	1.46
2	C	500	GNP	PG-O1G	9.46	1.60	1.46
2	D	500	GNP	PB-O3A	8.49	1.69	1.59
2	B	500	GNP	PB-O1B	7.78	1.58	1.46
2	B	500	GNP	PB-O3A	7.27	1.68	1.59
2	D	500	GNP	PB-O2B	-4.24	1.45	1.56
2	B	500	GNP	PB-O2B	-4.17	1.45	1.56
2	C	500	GNP	PG-O2G	-4.09	1.45	1.56
2	C	500	GNP	PB-O2B	-3.96	1.46	1.56
2	A	500	GNP	PG-O2G	-3.86	1.46	1.56
2	D	500	GNP	PG-O2G	-3.84	1.46	1.56
2	A	500	GNP	PB-O2B	-3.71	1.47	1.56
2	C	500	GNP	PB-O3A	2.64	1.62	1.59
2	D	500	GNP	PA-O5'	2.43	1.68	1.59
2	D	500	GNP	PG-O3G	2.35	1.63	1.56
2	A	500	GNP	C5-C6	-2.34	1.35	1.44

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	500	GNP	O1B-PB-N3B	-7.49	100.75	111.77
2	C	500	GNP	O1G-PG-N3B	-6.84	101.70	111.77
2	A	500	GNP	O2B-PB-O1B	-5.83	97.36	109.87
2	D	500	GNP	O3A-PB-N3B	5.59	122.10	106.59
2	D	500	GNP	O1B-PB-N3B	-5.32	103.93	111.77
2	A	500	GNP	O1G-PG-N3B	-5.24	104.06	111.77
2	C	500	GNP	O2B-PB-O3A	5.11	121.71	104.64
2	C	500	GNP	O2A-PA-O3A	4.23	118.70	107.27
2	A	500	GNP	O2G-PG-O3G	4.06	118.50	107.59
2	C	500	GNP	O2B-PB-O1B	-3.99	101.31	109.87
2	A	500	GNP	O2B-PB-O3A	3.74	117.13	104.64
2	A	500	GNP	O3A-PA-O1A	-3.39	100.51	110.70
2	A	500	GNP	O1B-PB-N3B	-3.16	107.12	111.77
2	C	500	GNP	O2G-PG-O3G	2.96	115.53	107.59
2	B	500	GNP	O2'-C2'-C3'	-2.76	102.96	111.82
2	D	500	GNP	O5'-PA-O1A	2.73	119.74	108.94
2	C	500	GNP	O5'-PA-O1A	2.35	118.23	108.94
2	B	500	GNP	O2A-PA-O5'	2.30	117.98	107.57
2	A	500	GNP	O2A-PA-O3A	2.27	113.42	107.27

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	500	GNP	O2G-PG-O1G	-2.26	107.78	113.45
2	B	500	GNP	O3'-C3'-C2'	-2.22	104.70	111.82
2	D	500	GNP	O1G-PG-N3B	-2.14	108.62	111.77
2	D	500	GNP	O2A-PA-O3A	2.12	112.99	107.27
2	C	500	GNP	O2'-C2'-C3'	-2.12	105.04	111.82
2	D	500	GNP	O3'-C3'-C2'	-2.11	105.06	111.82
2	D	500	GNP	O2B-PB-O3A	-2.04	97.84	104.64

There are no chirality outliers.



All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	500	GNP	PG-N3B-PB-O1B
2	A	500	GNP	PA-O3A-PB-O2B
2	C	500	GNP	PA-O3A-PB-O2B
2	D	500	GNP	PB-N3B-PG-O1G
2	B	500	GNP	PB-O3A-PA-O1A
2	A	500	GNP	PB-O3A-PA-O2A
2	C	500	GNP	PB-O3A-PA-O1A
2	C	500	GNP	PB-N3B-PG-O1G
2	C	500	GNP	PG-N3B-PB-O3A
2	A	500	GNP	PB-O3A-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	298/305 (97%)	-0.12	9 (3%)	52	54	30, 48, 82, 101	0
1	B	297/305 (97%)	-0.19	9 (3%)	52	54	24, 47, 79, 112	1 (0%)
1	C	297/305 (97%)	-0.13	7 (2%)	59	62	23, 48, 89, 111	1 (0%)
1	D	282/305 (92%)	0.09	12 (4%)	40	41	30, 58, 88, 101	1 (0%)
All	All	1174/1220 (96%)	-0.09	37 (3%)	50	52	23, 50, 87, 112	3 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	183	ALA	4.6
1	D	209	GLY	4.0
1	A	212	TYR	3.8
1	C	183	ALA	3.7
1	D	472	ILE	3.5
1	B	235	LYS	3.4
1	D	463	GLU	3.1
1	C	212	TYR	3.1
1	B	230	SER	3.1
1	B	179	TYR	3.0
1	B	212	TYR	3.0
1	D	208	ILE	2.9
1	D	229	PHE	2.9
1	B	208	ILE	2.9
1	D	460	ILE	2.8
1	A	234	LYS	2.8
1	A	235	LYS	2.8
1	A	232	ASN	2.7
1	C	184	THR	2.7
1	B	307[A]	ARG	2.5
1	D	476	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	215	VAL	2.5
1	B	183	ALA	2.4
1	A	208	ILE	2.4
1	C	233	LYS	2.3
1	A	236	GLY	2.3
1	A	182	ASN	2.3
1	D	464	PHE	2.2
1	D	459	ASN	2.2
1	B	211	GLY	2.2
1	C	235	LYS	2.1
1	A	186	VAL	2.1
1	D	462	GLN	2.1
1	C	307[A]	ARG	2.1
1	B	210	SER	2.0
1	C	185	ASN	2.0
1	D	467	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	C	702	1/1	0.88	0.09	71,71,71,71	0
4	CL	C	802	1/1	0.89	0.12	70,70,70,70	0
4	CL	B	802	1/1	0.92	0.12	74,74,74,74	0
2	GNP	D	500	32/32	0.92	0.08	49,56,73,81	4
3	MG	B	700	1/1	0.93	0.09	62,62,62,62	0
2	GNP	C	500	32/32	0.94	0.09	47,56,67,67	4
2	GNP	B	500	28/32	0.94	0.09	46,56,70,75	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CL	A	802	1/1	0.95	0.07	79,79,79,79	0
2	GNP	A	500	32/32	0.96	0.07	40,47,60,65	4
3	MG	B	702	1/1	0.97	0.04	66,66,66,66	0
3	MG	C	700	1/1	0.98	0.04	59,59,59,59	0
3	MG	A	702	1/1	0.98	0.08	66,66,66,66	0
3	MG	D	700	1/1	0.98	0.06	56,56,56,56	0
3	MG	A	700	1/1	0.99	0.04	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GNP D 500:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GNP C 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GNP B 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.