



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 08:44 AM UTC

PDB ID : 6CHG / pdb_00006chg
Title : Crystal structure of the yeast COMPASS catalytic module
Authors : Hsu, P.L.; Li, H.; Zheng, N.
Deposited on : 2018-02-22
Resolution : 2.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3580 (3.00-2.96)
Clashscore	190562	3904 (3.00-2.96)
Ramachandran outliers	187476	3761 (3.00-2.96)
Sidechain outliers	187428	3764 (3.00-2.96)
RSRZ outliers	180081	3579 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	7% 80% 17% ..
2	B	405	12% 77% 16% • 6%
3	C	153	11% 78% 16% 7%
4	D	439	9% 79% 13% • 7%
5	E	61	20% 59% 8% • 31%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	61	
6	J	4	

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 21039 atoms, of which 10402 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KLLA0E24487p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	305	Total	C	H	N	O	S	0	0	0
			4730	1500	2361	395	455	19			

- Molecule 2 is a protein called KLLA0C10945p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	382	Total	C	H	N	O	S	0	0	0
			6139	1985	3011	519	609	15			

- Molecule 3 is a protein called Histone-lysine N-methyltransferase, H3 lysine-4 specific.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	143	Total	C	H	N	O	S	0	0	0
			2297	722	1163	201	203	8			

- Molecule 4 is a protein called KLLA0A08800p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	408	Total	C	H	N	O	S	0	0	0
			6400	2095	3127	539	622	17			


- Molecule 5 is a protein called KLLA0E03521p.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
5	E	42	Total	C	H	N	O	S	0	0	0
			697	216	355	63	62	1			
5	F	43	Total	C	H	N	O	S	0	0	0
			670	211	341	57	59	2			

- Molecule 6 is a protein called H3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	3	Total	C	H	N	O	S	0	0
			47	14	23	4	5	1		0

- Molecule 7 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: $C_{15}H_{22}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total	C	H	N	O	S	0
			48	15	21	6	5	1	0

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	1	Total	Zn	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	O	0	0
			1	1		
9	B	3	Total	O	0	0
			3	3		
9	C	1	Total	O	0	0
			1	1		
9	D	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: KLLA0E24487p




- Molecule 2: KLLA0C10945p





- Molecule 3: Histone-lysine N-methyltransferase, H3 lysine-4 specific








• Molecule 4: KLLA0A08800p




• Molecule 5: KLLA0E03521p



• Molecule 5: KLLA0E03521p



• Molecule 6: H3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	163.84Å 138.32Å 136.13Å 90.00° 112.41° 90.00°	Depositor
Resolution (Å)	44.61 – 2.98 44.61 – 2.98	Depositor EDS
% Data completeness (in resolution range)	95.6 (44.61-2.98) 95.5 (44.61-2.98)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 3.01Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.228 , 0.271 0.233 , 0.280	Depositor DCC
R_{free} test set	2789 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	75.4	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	21039	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/2414	0.47	2/3267 (0.1%)
2	B	0.15	0/3198	0.39	1/4323 (0.0%)
3	C	0.16	0/1154	0.44	1/1550 (0.1%)
4	D	0.14	0/3360	0.56	6/4577 (0.1%)
5	E	0.12	0/347	0.38	0/466
5	F	0.14	0/333	0.43	0/447
6	J	0.08	0/23	0.16	0/29
All	All	0.14	0/10829	0.47	10/14659 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	D	324	LEU	N-CA-C	17.95	137.35	114.56
4	D	324	LEU	CB-CA-C	-16.75	83.00	109.13
1	A	107	MET	N-CA-C	10.76	126.01	112.24
4	D	264	LEU	CB-CA-C	-8.22	98.18	110.67
4	D	238	ARG	N-CA-C	-8.12	93.15	108.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2369	2361	2380	33	0
2	B	3128	3011	3011	45	0
3	C	1134	1163	1163	20	0
4	D	3273	3127	3136	40	0
5	E	342	355	355	5	0
5	F	329	341	341	10	0
6	J	24	23	23	0	0
7	C	27	21	22	1	0
8	C	1	0	0	0	0
9	A	1	0	0	0	0
9	B	3	0	0	0	0
9	C	1	0	0	0	0
9	D	5	0	0	0	0
All	All	10637	10402	10431	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:987:PRO:O	3:C:999:LEU:N	2.11	0.83
4:D:238:ARG:O	4:D:239:THR:HG22	1.79	0.81
5:F:124:ASP:OD1	5:F:125:ALA:N	2.18	0.77
4:D:238:ARG:O	4:D:239:THR:CG2	2.33	0.76
1:A:88:SER:OG	1:A:90:ASP:OD1	2.04	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/312 (96%)	277 (92%)	24 (8%)	0	100	100
2	B	376/405 (93%)	358 (95%)	17 (4%)	1 (0%)	36	67
3	C	139/153 (91%)	127 (91%)	11 (8%)	1 (1%)	18	50
4	D	400/439 (91%)	368 (92%)	32 (8%)	0	100	100
5	E	40/61 (66%)	37 (92%)	2 (5%)	1 (2%)	4	21
5	F	39/61 (64%)	35 (90%)	4 (10%)	0	100	100
6	J	1/4 (25%)	1 (100%)	0	0	100	100
All	All	1296/1435 (90%)	1203 (93%)	90 (7%)	3 (0%)	43	73

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	311	PRO
5	E	111	PRO
3	C	993	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/283 (98%)	269 (98%)	7 (2%)	42	71
2	B	352/375 (94%)	342 (97%)	10 (3%)	38	69
3	C	122/131 (93%)	121 (99%)	1 (1%)	73	86
4	D	360/387 (93%)	353 (98%)	7 (2%)	50	75
5	E	38/54 (70%)	38 (100%)	0	100	100
5	F	36/54 (67%)	34 (94%)	2 (6%)	19	50
6	J	3/4 (75%)	3 (100%)	0	100	100
All	All	1187/1288 (92%)	1160 (98%)	27 (2%)	44	72

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	340	ASN
3	C	848	LEU
4	D	429	ILE
2	B	399	LEU
4	D	8	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	133	HIS
2	B	256	GLN
5	F	120	GLN
4	D	122	HIS
2	B	127	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	SAM	C	1101	-	27,29,29	1.12	4 (14%)	34,42,42	2.04	8 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SAM	C	1101	-	-	5/17/33/33	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	1101	SAM	C2-N3	2.75	1.38	1.33
7	C	1101	SAM	C2-N1	2.58	1.38	1.33
7	C	1101	SAM	C5-N7	-2.22	1.35	1.39
7	C	1101	SAM	OXT-C	-2.18	1.23	1.30

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	1101	SAM	N3-C2-N1	-5.39	120.42	128.58
7	C	1101	SAM	C5-C4-N3	-4.88	119.99	126.72
7	C	1101	SAM	N9-C8-N7	-3.61	108.81	113.94
7	C	1101	SAM	N3-C4-N9	3.61	133.31	127.17
7	C	1101	SAM	C2-N3-C4	3.43	120.21	111.83

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	C	1101	SAM	CA-CB-CG-SD
7	C	1101	SAM	CB-CG-SD-CE
7	C	1101	SAM	O-C-CA-CB
7	C	1101	SAM	CB-CG-SD-C5'
7	C	1101	SAM	OXT-C-CA-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	1101	SAM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/312 (97%)	0.75	22 (7%) 21 13	52, 84, 121, 138	0
2	B	382/405 (94%)	0.93	47 (12%) 8 6	55, 87, 130, 164	0
3	C	143/153 (93%)	0.91	17 (11%) 9 6	59, 85, 169, 187	0
4	D	408/439 (92%)	0.78	40 (9%) 13 8	51, 84, 122, 156	0
5	E	42/61 (68%)	1.68	12 (28%) 1 1	107, 136, 156, 162	0
5	F	43/61 (70%)	2.22	22 (51%) 0 0	90, 128, 155, 159	0
6	J	3/4 (75%)	1.87	2 (66%) 0 0	83, 83, 108, 109	1 (33%)
All	All	1326/1435 (92%)	0.91	162 (12%) 8 6	51, 87, 140, 187	1 (0%)

The worst 5 of 162 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	309	TYR	8.5
4	D	434	TRP	6.1
3	C	989	LEU	5.6
3	C	853	LEU	5.4
2	B	304	GLU	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	ZN	C	1102	1/1	0.78	0.12	115,115,115,115	1
7	SAM	C	1101	27/27	0.92	0.13	66,98,132,139	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.