



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:19 AM UTC

PDB ID : 7CHS / pdb_00007chs
Title : Crystal structure of SARS-CoV-2 antibody P22A-1D1 with RBD
Authors : Wang, X.; Zhang, L.; Ge, J.; Wang, R.; Zhang, Q.
Deposited on : 2020-07-06
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	217	13% 70% 16% • 12%
2	H	217	2% 86% 11% •
3	L	213	3% 82% 16% •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	190	Total	C	N	O	S	0	0	0
			1505	963	250	284	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	530	HIS	-	expression tag	UNP P0DTC2
E	531	HIS	-	expression tag	UNP P0DTC2
E	532	HIS	-	expression tag	UNP P0DTC2
E	533	HIS	-	expression tag	UNP P0DTC2
E	534	HIS	-	expression tag	UNP P0DTC2
E	535	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called antibody P22A-1D1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	211	Total	C	N	O	S	0	0	0
			1578	993	264	314	7			

- Molecule 3 is a protein called antibody P22A-1D1 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	212	Total	C	N	O	S	0	0	0
			1630	1026	273	327	4			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	69	Total	O	0	0
			69	69		
5	H	121	Total	O	0	0
			121	121		
5	L	85	Total	O	0	0
			85	85		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1



- Molecule 2: antibody P22A-1D1 heavy chain



- Molecule 3: antibody P22A-1D1 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	193.34Å 86.60Å 57.16Å 90.00° 99.25° 90.00°	Depositor
Resolution (Å)	47.71 – 2.40 47.71 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.71-2.40) 99.8 (47.71-2.40)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.191 , 0.224 0.199 , 0.230	Depositor DCC
R_{free} test set	1834 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.189	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5002	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	E	0.88	10/1546 (0.6%)	0.71	2/2101 (0.1%)
2	H	0.80	8/1614 (0.5%)	0.65	1/2199 (0.0%)
3	L	0.88	11/1665 (0.7%)	0.74	3/2260 (0.1%)
All	All	0.85	29/4825 (0.6%)	0.70	6/6560 (0.1%)

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	30	SER	C-O	-9.62	1.15	1.24
2	H	113	VAL	C-O	-9.04	1.14	1.24
2	H	139	THR	C-O	-8.35	1.14	1.24
1	E	492	LEU	C-O	-7.62	1.14	1.23
2	H	112	THR	C-O	-7.20	1.15	1.24
1	E	405	ASP	C-O	-7.02	1.15	1.24
2	H	102	TYR	C-O	-6.94	1.14	1.24
3	L	56	SER	C-O	-6.78	1.15	1.23
3	L	45	LYS	C-O	-6.62	1.15	1.23
2	H	210	LYS	C-O	-6.51	1.16	1.24
3	L	56	SER	CA-C	-6.49	1.44	1.52
1	E	357	ARG	C-O	-6.27	1.16	1.24
1	E	445	VAL	C-O	-6.23	1.17	1.24
1	E	448	ASN	C-O	-6.09	1.16	1.24
2	H	111	THR	C-O	-6.06	1.16	1.23
3	L	150	ASP	C-O	-5.88	1.16	1.23
3	L	30	SER	CA-C	-5.84	1.48	1.53
1	E	379	CYS	C-O	-5.77	1.16	1.24
3	L	160	GLU	C-O	-5.77	1.16	1.23
2	H	207	SER	C-O	-5.66	1.16	1.24
3	L	149	VAL	C-O	-5.65	1.17	1.24
1	E	446	GLY	C-O	-5.60	1.16	1.24
1	E	382	VAL	C-O	-5.58	1.17	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	2	ILE	CA-CB	-5.50	1.47	1.54
3	L	31	SER	C-O	-5.46	1.17	1.24
3	L	2	ILE	N-CA	-5.30	1.39	1.46
1	E	448	ASN	CG-ND2	-5.14	1.22	1.33
1	E	405	ASP	N-CA	-5.13	1.39	1.46
2	H	86	ARG	C-O	-5.00	1.17	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	404	GLY	N-CA-C	6.57	120.62	112.73
3	L	91	LEU	N-CA-C	6.36	118.44	108.96
1	E	446	GLY	N-CA-C	-6.04	106.72	113.79
3	L	2	ILE	CB-CA-C	-5.40	103.50	110.84
3	L	92	ASN	N-CA-CB	5.29	119.43	110.49
2	H	112	THR	N-CA-C	5.26	117.82	109.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1505	0	1422	29	0
2	H	1578	0	1537	12	0
3	L	1630	0	1599	19	0
4	E	14	0	13	0	0
5	E	69	0	0	4	0
5	H	121	0	0	1	0
5	L	85	0	0	2	0
All	All	5002	0	4571	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:391:CYS:SG	1:E:525:CYS:SG	1.35	1.33
1:E:379:CYS:SG	1:E:432:CYS:SG	1.40	1.24
3:L:107:ARG:NH1	3:L:108:THR:O	2.03	0.92
1:E:379:CYS:SG	1:E:432:CYS:CB	2.58	0.91
1:E:391:CYS:CB	1:E:525:CYS:SG	2.64	0.85
1:E:427:ASP:OD1	5:E:701:HOH:O	1.94	0.84
1:E:334:ASN:HD22	1:E:335:LEU:H	1.31	0.77
1:E:371:SER:OG	1:E:373:SER:OG	2.03	0.75
2:H:203:ASN:HD21	2:H:205:LYS:HE3	1.55	0.71
1:E:340:GLU:OE2	1:E:356:LYS:NZ	2.25	0.69
3:L:196:THR:OG1	5:L:302:HOH:O	2.12	0.68
3:L:209:ASN:OD1	5:L:301:HOH:O	2.11	0.67
2:H:32:ASN:HB2	2:H:34:MET:HE1	1.75	0.67
1:E:393:THR:O	1:E:523:THR:OG1	2.11	0.66
1:E:474:GLN:O	5:E:702:HOH:O	2.14	0.66
3:L:183:ALA:O	3:L:187:LYS:HG3	1.97	0.64
2:H:102:TYR:OH	5:H:301:HOH:O	2.14	0.64
1:E:383:SER:HB2	1:E:386:LYS:HB2	1.80	0.62
3:L:187:LYS:O	3:L:188:HIS:ND1	2.32	0.62
1:E:448:ASN:O	5:E:703:HOH:O	2.17	0.60
1:E:382:VAL:HG12	1:E:383:SER:N	2.17	0.60
1:E:357:ARG:NH2	1:E:394:ASN:ND2	2.50	0.59
2:H:33:TYR:HB2	2:H:98:ASP:O	2.04	0.56
2:H:99:ARG:HB2	2:H:103:GLY:O	2.04	0.56
1:E:395:VAL:HG22	1:E:515:PHE:HD1	1.71	0.56
2:H:203:ASN:ND2	2:H:205:LYS:HE3	2.19	0.55
3:L:187:LYS:O	3:L:188:HIS:CG	2.60	0.55
1:E:357:ARG:NH2	1:E:394:ASN:HD21	2.04	0.54
1:E:334:ASN:HD22	1:E:335:LEU:N	2.02	0.53
1:E:394:ASN:O	1:E:516:GLU:HB3	2.10	0.51
1:E:334:ASN:ND2	1:E:335:LEU:H	2.04	0.51
1:E:493:GLN:HE21	1:E:494:SER:H	1.59	0.50
2:H:4:LEU:HD22	2:H:22:CYS:SG	2.53	0.49
3:L:182:LYS:HZ3	3:L:186:GLU:CD	2.21	0.48
3:L:189:LYS:HD2	3:L:209:ASN:HB3	1.96	0.48
3:L:89:LEU:C	3:L:89:LEU:HD23	2.39	0.48
2:H:138:GLY:O	2:H:189:PRO:HA	2.15	0.47
3:L:6:GLN:NE2	3:L:86:TYR:O	2.43	0.47
1:E:369:TYR:CD1	1:E:369:TYR:O	2.68	0.47
1:E:403:ARG:HH11	3:L:92:ASN:HB3	1.80	0.46
1:E:382:VAL:CG1	1:E:383:SER:N	2.79	0.46
3:L:123:GLN:HG2	3:L:128:THR:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:34:MET:HB2	2:H:34:MET:HE2	1.54	0.45
1:E:369:TYR:O	1:E:369:TYR:CG	2.69	0.44
3:L:19:VAL:HG22	3:L:75:ILE:HB	1.98	0.43
1:E:458:LYS:NZ	5:E:702:HOH:O	2.50	0.43
1:E:365:TYR:HB2	1:E:387:LEU:CD1	2.48	0.43
3:L:21:ILE:O	3:L:72:THR:HA	2.19	0.43
3:L:21:ILE:HG12	3:L:101:THR:HG21	2.01	0.43
1:E:384:PRO:HA	1:E:387:LEU:HG	2.00	0.42
3:L:4:LEU:HD13	3:L:88:CYS:SG	2.58	0.42
3:L:165:GLN:HG3	3:L:172:TYR:CE2	2.55	0.42
1:E:444:LYS:HA	1:E:444:LYS:HD3	1.86	0.41
2:H:64:LYS:HB2	2:H:64:LYS:HE2	1.73	0.41
2:H:59:TYR:O	2:H:64:LYS:NZ	2.53	0.41
1:E:339:GLY:O	1:E:343:ASN:HB2	2.20	0.41
3:L:179:THR:O	3:L:180:LEU:HD12	2.21	0.40
2:H:102:TYR:HA	3:L:49:TYR:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	186/217 (86%)	177 (95%)	9 (5%)	0	100	100
2	H	207/217 (95%)	204 (99%)	3 (1%)	0	100	100
3	L	210/213 (99%)	201 (96%)	8 (4%)	1 (0%)	24	37
All	All	603/647 (93%)	582 (96%)	20 (3%)	1 (0%)	43	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	92	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	164/190 (86%)	163 (99%)	1 (1%)	78	89
2	H	177/182 (97%)	174 (98%)	3 (2%)	53	74
3	L	185/186 (100%)	180 (97%)	5 (3%)	39	62
All	All	526/558 (94%)	517 (98%)	9 (2%)	53	74

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	373	SER
2	H	131	SER
2	H	182	LEU
2	H	216	GLU
3	L	2	ILE
3	L	30	SER
3	L	31	SER
3	L	90	HIS
3	L	92	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	334	ASN
2	H	81	GLN
2	H	83	ASN
2	H	203	ASN
3	L	92	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	E	601	1	14,14,15	0.37	0	17,19,21	0.59	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	601	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	601	NAG	C1-O5-C5	2.03	114.91	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	190/217 (87%)	0.47	29 (15%) 5 4	27, 40, 87, 118	0
2	H	211/217 (97%)	-0.08	4 (1%) 66 62	27, 35, 54, 79	0
3	L	212/213 (99%)	0.10	6 (2%) 55 51	28, 39, 58, 81	0
All	All	613/647 (94%)	0.15	39 (6%) 25 22	27, 38, 77, 118	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	138	GLY	6.9
1	E	521	PRO	6.6
1	E	526	GLY	5.4
1	E	382	VAL	4.7
1	E	390	LEU	4.1
1	E	384	PRO	4.0
1	E	387	LEU	3.9
1	E	385	THR	3.8
1	E	333	THR	3.6
2	H	131	SER	3.6
1	E	335	LEU	3.6
1	E	391	CYS	3.4
1	E	381	GLY	3.1
1	E	362	VAL	2.9
3	L	92	ASN	2.9
1	E	525	CYS	2.8
1	E	516	GLU	2.8
1	E	386	LYS	2.7
1	E	523	THR	2.7
1	E	389	ASP	2.6
1	E	334	ASN	2.6
1	E	365	TYR	2.6
2	H	217	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	393	THR	2.3
2	H	101	TYR	2.3
1	E	369	TYR	2.3
3	L	66	GLY	2.3
1	E	371	SER	2.3
1	E	388	ASN	2.2
3	L	51	ALA	2.2
1	E	522	ALA	2.2
3	L	93	SER	2.1
1	E	368	LEU	2.1
3	L	68	GLY	2.1
1	E	383	SER	2.1
1	E	372	ALA	2.1
1	E	392	PHE	2.0
1	E	359	SER	2.0
3	L	52	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	E	601	14/15	0.64	0.19	59,69,75,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.