



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 10:05 PM UTC

PDB ID : 8CH7 / pdb_00008ch7
BMRB ID : 34787
Title : RDC-refined Interleukin-4 (wild type) pH 5.6
Authors : Vaz, D.C.; Rodrigues, J.R.; Loureiro-Ferreira, N.; Mueller, T.; Sebald, W.;
Redfield, C.; Brito, R.M.M.
Deposited on : 2023-02-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 65%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	129	

2 Ensemble composition and analysis ⓘ

This entry contains 2 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 2116 atoms, of which 1068 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Interleukin-4.


Mol	Chain	Residues	Atoms						Trace
1	A	129	Total	C	H	N	O	S	0
			2116	653	1068	192	196	7	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Interleukin-4




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Interleukin-4



4.2.2 Score per residue for model 2

- Molecule 1: Interleukin-4





5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 200 calculated structures, 2 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	3.6
X-PLOR NIH	structure calculation	3.6

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1207
Number of shifts mapped to atoms	1207
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	65%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1048	1068	1063	42±0
All	All	2096	2136	2126	83

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:46:CYS:SG	1:A:47:ARG:N	0.70	2.64	2	1
1:A:56:TYR:CE1	1:A:83:LEU:HD12	0.69	2.22	2	1
1:A:38:ASN:H	1:A:38:ASN:HD22	0.62	1.36	2	1
1:A:56:TYR:CG	1:A:57:SER:N	0.61	2.68	1	1
1:A:38:ASN:HD22	1:A:38:ASN:N	0.60	1.95	2	1
1:A:108:THR:O	1:A:110:GLU:N	0.59	2.34	2	2
1:A:56:TYR:CD2	1:A:56:TYR:C	0.59	2.79	1	1
1:A:69:THR:OG1	1:A:70:ALA:N	0.58	2.34	1	2
1:A:108:THR:O	1:A:109:LEU:C	0.57	2.47	1	2
1:A:38:ASN:H	1:A:38:ASN:ND2	0.56	1.98	2	1
1:A:55:PHE:O	1:A:59:HIS:CG	0.56	2.59	1	2
1:A:108:THR:O	1:A:111:ASN:N	0.55	2.39	2	2
1:A:11:ILE:O	1:A:15:ASN:ND2	0.54	2.41	1	2
1:A:75:ARG:O	1:A:75:ARG:NE	0.54	2.41	2	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:47:ARG:N	1:A:99:CYS:SG	0.53	2.81	2	1
1:A:22:THR:OG1	1:A:23:LEU:N	0.53	2.41	2	1
1:A:66:LEU:C	1:A:68:ALA:H	0.53	2.12	1	2
1:A:96:LEU:HD12	1:A:97:ASN:N	0.52	2.19	1	1
1:A:56:TYR:CE1	1:A:83:LEU:CD1	0.52	2.93	2	1
1:A:76:HIS:NE2	1:A:80:ILE:HD11	0.52	2.18	2	1
1:A:91:TRP:CE2	1:A:97:ASN:O	0.52	2.63	1	1
1:A:91:TRP:CZ2	1:A:97:ASN:O	0.52	2.63	1	1
1:A:38:ASN:N	1:A:38:ASN:ND2	0.49	2.60	2	1
1:A:68:ALA:HB3	1:A:72:GLN:HG3	0.49	1.85	1	1
1:A:98:SER:C	1:A:99:CYS:SG	0.48	2.96	1	1
1:A:24:CYS:SG	1:A:65:CYS:O	0.48	2.71	1	2
1:A:46:CYS:SG	1:A:99:CYS:N	0.48	2.86	1	1
1:A:43:GLU:O	1:A:46:CYS:SG	0.48	2.71	2	1
1:A:59:HIS:O	1:A:60:GLU:C	0.48	2.57	1	2
1:A:33:PHE:C	1:A:35:ALA:H	0.48	2.17	2	2
1:A:66:LEU:C	1:A:68:ALA:N	0.48	2.71	1	2
1:A:96:LEU:CD1	1:A:97:ASN:N	0.47	2.77	1	1
1:A:46:CYS:C	1:A:99:CYS:SG	0.47	2.97	2	1
1:A:96:LEU:N	1:A:96:LEU:CD2	0.47	2.78	2	1
1:A:72:GLN:O	1:A:73:PHE:C	0.47	2.57	2	2
1:A:96:LEU:N	1:A:96:LEU:HD22	0.47	2.25	2	1
1:A:47:ARG:O	1:A:48:ALA:C	0.46	2.58	2	2
1:A:69:THR:C	1:A:71:GLN:N	0.46	2.72	1	2
1:A:56:TYR:CD2	1:A:57:SER:N	0.45	2.85	1	1
1:A:80:ILE:CG2	1:A:84:LYS:NZ	0.45	2.80	2	1
1:A:86:LEU:O	1:A:87:ASP:C	0.45	2.60	1	2
1:A:116:LEU:O	1:A:116:LEU:HD13	0.45	2.11	1	1
1:A:93:LEU:O	1:A:94:ALA:C	0.45	2.59	1	1
1:A:29:VAL:HG23	1:A:30:THR:N	0.44	2.27	2	1
1:A:125:SER:C	1:A:127:CYS:H	0.44	2.21	1	1
1:A:62:ASP:OD1	1:A:62:ASP:O	0.44	2.35	2	1
1:A:54:GLN:O	1:A:55:PHE:C	0.44	2.59	1	1
1:A:52:LEU:O	1:A:53:ARG:C	0.44	2.60	2	1
1:A:48:ALA:O	1:A:49:ALA:C	0.43	2.61	2	1
1:A:109:LEU:O	1:A:109:LEU:HD23	0.43	2.13	2	1
1:A:86:LEU:O	1:A:89:ASN:N	0.43	2.51	1	1
1:A:11:ILE:O	1:A:15:ASN:CG	0.43	2.61	2	1
1:A:109:LEU:O	1:A:110:GLU:C	0.43	2.62	1	1
1:A:45:PHE:O	1:A:49:ALA:HB2	0.43	2.14	2	1
1:A:21:LYS:CB	1:A:21:LYS:NZ	0.42	2.82	1	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:54:GLN:N	1:A:54:GLN:CD	0.42	2.77	1	1
1:A:68:ALA:HB3	1:A:72:GLN:CG	0.42	2.44	1	1
1:A:19:GLU:O	1:A:19:GLU:CD	0.42	2.62	1	1
1:A:101:VAL:O	1:A:101:VAL:HG13	0.42	2.14	2	1
1:A:9:GLU:OE1	1:A:89:ASN:OD1	0.42	2.37	1	1
1:A:34:ALA:CB	1:A:105:ASN:HD22	0.42	2.28	1	1
1:A:76:HIS:O	1:A:77:LYS:C	0.42	2.61	1	1
1:A:72:GLN:O	1:A:75:ARG:N	0.41	2.53	1	1
1:A:101:VAL:HG13	1:A:101:VAL:O	0.41	2.15	1	1
1:A:53:ARG:O	1:A:56:TYR:CD1	0.40	2.74	1	1
1:A:15:ASN:HD22	1:A:15:ASN:N	0.40	2.13	2	1
1:A:113:LEU:N	1:A:113:LEU:CD2	0.40	2.83	1	1
1:A:1:HIS:NE2	1:A:124:TYR:OH	0.40	2.53	2	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/129 (98%)	98±0 (77±0%)	28±2 (22±1%)	2±1 (2±1%)	10	55
All	All	254/258 (98%)	195 (77%)	55 (22%)	4 (2%)	10	55

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	109	LEU	2
1	A	22	THR	1
1	A	34	ALA	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/118 (100%)	110±1 (93±1%)	8±1 (7±1%)	16	65
All	All	236/236 (100%)	220 (93%)	16 (7%)	16	65

All 14 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	24	CYS	2
1	A	29	VAL	2
1	A	18	THR	1
1	A	21	LYS	1
1	A	56	TYR	1
1	A	69	THR	1
1	A	116	LEU	1
1	A	28	THR	1
1	A	38	ASN	1
1	A	52	LEU	1
1	A	58	HIS	1
1	A	65	CYS	1
1	A	75	ARG	1
1	A	109	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 65% for the well-defined parts and 65% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *ChemicalShifts_WTIL4_pH56_BMRB-STAR_fev23.txt*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1207
Number of shifts mapped to atoms	1207
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	123	1.60 ± 0.10	Should be checked
$^{13}\text{C}_\beta$	59	2.15 ± 0.10	Should be checked
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	126	0.19 ± 0.40	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 65%, i.e. 1192 atoms were assigned a chemical shift out of a possible 1833. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	506/646 (78%)	257/260 (99%)	123/258 (48%)	126/128 (98%)
Sidechain	645/1057 (61%)	495/681 (73%)	137/324 (42%)	13/52 (25%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	41/130 (32%)	41/64 (64%)	0/55 (0%)	0/11 (0%)
Overall	1192/1833 (65%)	793/1005 (79%)	260/637 (41%)	139/191 (73%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 65%, i.e. 1192 atoms were assigned a chemical shift out of a possible 1833. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	506/646 (78%)	257/260 (99%)	123/258 (48%)	126/128 (98%)
Sidechain	645/1057 (61%)	495/681 (73%)	137/324 (42%)	13/52 (25%)
Aromatic	41/130 (32%)	41/64 (64%)	0/55 (0%)	0/11 (0%)
Overall	1192/1833 (65%)	793/1005 (79%)	260/637 (41%)	139/191 (73%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	12	LYS	HE2	1.40	1.95 – 3.88	-7.8
1	A	12	LYS	HE3	1.40	1.92 – 3.89	-7.7

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

