



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:59 PM UTC

PDB ID : 8CHE / pdb_00008che
Title : TLT-1 binding Fab of the bispecific antibody HMB-001 in complex with the TLT-1 stalk peptide
Authors : Johansson, E.; Schluckebier, G.
Deposited on : 2023-02-07
Resolution : 1.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 4% 96%
1	H	215	 3% 96%
2	B	219	 2% 98%
2	L	219	 5% 91% 9%
3	C	37	 5% 38% 59%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	37	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a small red segment on the left labeled '3%', a larger green segment in the middle labeled '38%', and a large grey segment on the right labeled '59%'. A small black dot is located at the end of the green segment.</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 14866 atoms, of which 6832 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	H	214	Total 3304	C 1045	H 1634	N 281	O 336	S 8	0	14	0
1	A	214	Total 3264	C 1032	H 1613	N 279	O 332	S 8	0	10	0

- Molecule 2 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	L	219	Total 3442	C 1086	H 1697	N 302	O 349	S 8	0	9	0
2	B	219	Total 3419	C 1080	H 1686	N 300	O 345	S 8	0	6	0

- Molecule 3 is a protein called Trem-like transcript 1 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	C	15	Total 210	C 68	H 101	N 19	O 22	0	0	0
3	D	15	Total 210	C 68	H 101	N 19	O 22	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	261	Total 261	O 261	0	0
4	L	227	Total 227	O 227	0	0
4	A	275	Total 275	O 275	0	0
4	B	215	Total 215	O 215	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	21	Total O 21 21	0	0
4	D	18	Total O 18 18	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab heavy chain




- Molecule 1: Fab heavy chain




- Molecule 2: Fab light chain




- Molecule 2: Fab light chain



- Molecule 3: Trem-like transcript 1 protein



- Molecule 3: Trem-like transcript 1 protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	53.23Å 65.38Å 67.14Å 91.88° 91.72° 92.89°	Depositor
Resolution (Å)	28.46 – 1.49 28.46 – 1.49	Depositor EDS
% Data completeness (in resolution range)	91.0 (28.46-1.49) 91.0 (28.46-1.49)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.49Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.157 , 0.174 0.156 , 0.174	Depositor DCC
R_{free} test set	1968 reflections (1.22%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.234	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.007 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14866	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1698	0.61	0/2315
1	H	0.47	0/1723	0.58	0/2352
2	B	0.42	0/1781	0.54	0/2417
2	L	0.38	0/1802	0.50	0/2445
3	C	0.33	0/111	0.40	0/149
3	D	0.38	0/111	0.48	0/149
All	All	0.44	0/7226	0.56	0/9827

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1651	1613	1591	6	0
1	H	1670	1634	1592	8	0
2	B	1733	1686	1675	5	0
2	L	1745	1697	1676	14	0
3	C	109	101	101	1	0
3	D	109	101	101	1	0
4	A	275	0	0	5	3
4	B	215	0	0	1	3
4	C	21	0	0	1	0
4	D	18	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	261	0	0	5	4
4	L	227	0	0	4	3
All	All	8034	6832	6736	34	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:162:CYS:SG	4:B:493:HOH:O	2.10	1.07
1:A:87:ARG:NH2	4:A:301:HOH:O	2.01	0.93
2:L:162:CYS:SG	4:L:518:HOH:O	2.36	0.83
1:A:131:SER:OG	4:A:302:HOH:O	2.04	0.75
1:H:205:ASN:ND2	4:H:301:HOH:O	2.26	0.69

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:513:HOH:O	4:B:438:HOH:O[1_455]	1.82	0.38
4:H:478:HOH:O	4:L:425:HOH:O[1_455]	1.97	0.23
4:A:538:HOH:O	4:B:485:HOH:O[1_554]	2.06	0.14
4:H:303:HOH:O	4:A:504:HOH:O[1_565]	2.09	0.11
4:H:327:HOH:O	4:L:367:HOH:O[1_455]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	222/215 (103%)	217 (98%)	5 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	224/215 (104%)	219 (98%)	5 (2%)	0	100	100
2	B	223/219 (102%)	217 (97%)	6 (3%)	0	100	100
2	L	226/219 (103%)	218 (96%)	7 (3%)	1 (0%)	30	12
3	C	13/37 (35%)	13 (100%)	0	0	100	100
3	D	13/37 (35%)	13 (100%)	0	0	100	100
All	All	921/942 (98%)	897 (97%)	23 (2%)	1 (0%)	48	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	32	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/182 (105%)	191 (100%)	0	100	100
1	H	194/182 (107%)	194 (100%)	0	100	100
2	B	202/196 (103%)	202 (100%)	0	100	100
2	L	205/196 (105%)	203 (99%)	2 (1%)	68	45
3	C	11/31 (36%)	11 (100%)	0	100	100
3	D	11/31 (36%)	11 (100%)	0	100	100
All	All	814/818 (100%)	812 (100%)	2 (0%)	86	77

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	218	GLU
2	L	219	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	33	GLN
2	B	47	GLN
2	B	58	ASN
2	L	165	GLN
2	L	194	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/215 (99%)	-0.30	8 (3%) 45 49	7, 18, 42, 136	9 (4%)
1	H	214/215 (99%)	-0.21	6 (2%) 55 59	8, 19, 42, 67	11 (5%)
2	B	219/219 (100%)	-0.14	4 (1%) 67 71	8, 23, 44, 75	5 (2%)
2	L	219/219 (100%)	0.02	10 (4%) 37 41	9, 24, 49, 118	7 (3%)
3	C	15/37 (40%)	0.38	2 (13%) 7 7	20, 25, 47, 95	0
3	D	15/37 (40%)	0.23	1 (6%) 24 26	18, 21, 53, 82	0
All	All	896/942 (95%)	-0.14	31 (3%) 47 51	7, 21, 45, 136	32 (3%)

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	130	ARG	6.9
2	L	219	CYS	4.1
1	A	128	CYS	4.0
1	A	132	THR	4.0
1	A	131	SER	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.