



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:00 AM UTC

PDB ID : 8CJ7 / pdb_00008cj7
Title : HDAC6 selective degraded (difluoromethyl)-1,3,4-oxadiazole substrate inhibitor
Authors : Sandmark, J.; Ek, M.; Ripa, L.
Deposited on : 2023-02-12
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	4% 91% 5% ..
1	B	364	6% 93% ..
1	C	364	32% 93% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone deacetylase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	7	0
			2807	1767	502	520	18			
1	B	352	Total	C	N	O	S	0	5	0
			2764	1740	494	512	18			
1	C	352	Total	C	N	O	S	0	3	0
			2756	1732	494	512	18			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	435	SER	-	expression tag	UNP F8W4B7
A	436	ASN	-	expression tag	UNP F8W4B7
A	437	ALA	-	expression tag	UNP F8W4B7
A	438	GLY	-	expression tag	UNP F8W4B7
A	439	GLY	-	expression tag	UNP F8W4B7
B	435	SER	-	expression tag	UNP F8W4B7
B	436	ASN	-	expression tag	UNP F8W4B7
B	437	ALA	-	expression tag	UNP F8W4B7
B	438	GLY	-	expression tag	UNP F8W4B7
B	439	GLY	-	expression tag	UNP F8W4B7
C	435	SER	-	expression tag	UNP F8W4B7
C	436	ASN	-	expression tag	UNP F8W4B7
C	437	ALA	-	expression tag	UNP F8W4B7
C	438	GLY	-	expression tag	UNP F8W4B7
C	439	GLY	-	expression tag	UNP F8W4B7

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	K	0	0
			2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	K	0	0
			2	2		
2	C	2	Total	K	0	0
			2	2		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 6-[(5-pyridin-2-yl-1,2,4-triazol-3-yl)methyl]pyridine-3-carbohydrazide (CCD ID: UTO) (formula: C₁₃H₁₂N₈O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			22	13	8	1		
4	B	1	Total	C	N	O	0	1
			44	26	16	2		

- Molecule 5 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	4	Total I 4 4	0	0
5	B	1	Total I 1 1	0	0
5	C	1	Total I 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	268	Total O 268 268	0	0
6	B	264	Total O 264 264	0	0
6	C	102	Total O 102 102	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone deacetylase 6



• Molecule 1: Histone deacetylase 6



• Molecule 1: Histone deacetylase 6



A791	F792	F793	W794	S795	S796	L797	R798
------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.98Å 84.06Å 96.13Å 90.00° 110.25° 90.00°	Depositor
Resolution (Å)	90.19 – 1.51 90.19 – 1.51	Depositor EDS
% Data completeness (in resolution range)	51.8 (90.19-1.51) 52.0 (90.19-1.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 1.50Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.206 , 0.227 0.213 , 0.213	Depositor DCC
R_{free} test set	5667 reflections (2.51%)	wwPDB-VP
Wilson B-factor (Å ²)	13.6	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for l,-k,h	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9042	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: UTO, K, IOD, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	1/2888 (0.0%)	1.12	5/3914 (0.1%)
1	B	0.87	0/2848	1.12	3/3860 (0.1%)
1	C	0.80	0/2833	1.19	4/3838 (0.1%)
All	All	0.86	1/8569 (0.0%)	1.14	12/11612 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	730	MET	SD-CE	6.22	1.95	1.79

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	583	PHE	CA-CB-CG	6.57	120.37	113.80
1	B	583	PHE	CA-CB-CG	5.94	119.74	113.80
1	A	494	ALA	N-CA-C	-5.75	102.39	110.50
1	A	583	PHE	CA-CB-CG	5.70	119.50	113.80
1	A	773	THR	N-CA-C	5.54	114.51	108.14
1	A	703	GLY	N-CA-C	-5.53	104.62	112.37
1	C	703	GLY	N-CA-C	-5.49	104.68	112.37
1	B	494	ALA	N-CA-C	-5.34	102.97	110.50
1	C	612	ASP	N-CA-C	-5.33	103.10	110.35
1	B	706	ALA	N-CA-C	-5.26	106.71	113.02
1	A	706	ALA	N-CA-C	-5.12	106.81	113.16
1	C	706	ALA	N-CA-C	-5.09	106.84	113.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2807	0	2739	10	0
1	B	2764	0	2693	7	0
1	C	2756	0	2677	3	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	22	0	0	1	0
4	B	44	0	0	1	0
5	A	4	0	0	1	0
5	B	1	0	0	1	0
5	C	1	0	0	0	0
6	A	268	0	0	0	0
6	B	264	0	0	0	0
6	C	102	0	0	0	0
All	All	9042	0	8109	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:565[B]:VAL:HG22	1:A:754:MET:HE2	1.68	0.74
1:B:565[B]:VAL:HG22	1:B:754:MET:HE2	1.69	0.72
1:A:565[B]:VAL:CG2	1:A:754:MET:HE2	2.32	0.59
1:B:565[B]:VAL:CG2	1:B:754:MET:HE2	2.34	0.58
1:C:565:VAL:HG13	1:C:761:LEU:HD12	1.86	0.57
1:B:565[A]:VAL:HG13	1:B:761:LEU:HD12	1.87	0.57
1:A:565[A]:VAL:HG13	1:A:761:LEU:HD12	1.88	0.55
1:C:572:GLY:HA2	1:C:584:CYS:HB3	1.90	0.51
1:A:769:LEU:HD13	1:A:772:LEU:HD21	1.93	0.50
1:A:481:ARG:HA	5:A:808:IOD:I	2.82	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:574:HIS:NE2	4:B:804[B]:UTO:N20	2.60	0.50
1:B:481:ARG:HA	5:B:805:IOD:I	2.83	0.48
1:B:624:GLU:HG3	1:B:656:LEU:HD12	1.96	0.47
1:A:672[A]:LYS:HG3	1:A:714:GLY:O	2.16	0.45
1:B:597:GLN:HG2	1:B:604:LEU:HB3	1.99	0.44
1:A:574:HIS:NE2	4:A:804:UTO:N20	2.65	0.44
1:A:676:PRO:HG3	1:A:772:LEU:HB2	1.99	0.44
1:A:597:GLN:HG2	1:A:604:LEU:HB3	2.00	0.43
1:A:441:SER:CB	1:A:442:PRO:HD3	2.49	0.43
1:C:597:GLN:HG2	1:C:604:LEU:HB3	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/364 (99%)	352 (98%)	5 (1%)	2 (1%)	21	6
1	B	353/364 (97%)	347 (98%)	6 (2%)	0	100	100
1	C	351/364 (96%)	342 (97%)	9 (3%)	0	100	100
All	All	1063/1092 (97%)	1041 (98%)	20 (2%)	2 (0%)	48	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	672[A]	LYS
1	A	672[B]	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/305 (100%)	298 (98%)	6 (2%)	48	19
1	B	300/305 (98%)	297 (99%)	3 (1%)	68	44
1	C	298/305 (98%)	292 (98%)	6 (2%)	48	19
All	All	902/915 (99%)	887 (98%)	15 (2%)	53	24

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	465	GLU
1	A	579	THR
1	A	583	PHE
1	A	701	SER
1	A	769	LEU
1	A	772	LEU
1	B	465	GLU
1	B	583	PHE
1	B	701	SER
1	C	465	GLU
1	C	579	THR
1	C	583	PHE
1	C	701	SER
1	C	768	SER
1	C	788	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	560	GLN
1	B	487	HIS
1	C	560	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 15 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	UTO	B	804[A]	3	24,24,24	1.60	4 (16%)	30,32,32	2.52	12 (40%)
4	UTO	B	804[B]	3	24,24,24	1.60	3 (12%)	30,32,32	2.53	10 (33%)
4	UTO	A	804	3	24,24,24	1.54	4 (16%)	30,32,32	2.74	11 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UTO	B	804[A]	3	-	0/14/14/14	0/3/3/3
4	UTO	B	804[B]	3	-	5/14/14/14	0/3/3/3
4	UTO	A	804	3	-	0/14/14/14	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	804[A]	UTO	O1-C2	5.42	1.36	1.23
4	B	804[B]	UTO	O1-C2	4.92	1.34	1.23
4	A	804	UTO	O1-C2	4.64	1.34	1.23
4	B	804[B]	UTO	C2-N20	-3.61	1.30	1.33
4	A	804	UTO	C2-N20	-3.31	1.30	1.33
4	B	804[B]	UTO	N7-N8	-2.84	1.29	1.33
4	B	804[A]	UTO	C2-N20	-2.80	1.31	1.33
4	A	804	UTO	N7-N8	-2.76	1.29	1.33
4	B	804[A]	UTO	N7-N8	-2.62	1.29	1.33
4	B	804[A]	UTO	N7-N17	-2.10	1.29	1.32
4	A	804	UTO	N7-N17	-2.01	1.29	1.32

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	804	UTO	C3-C2-N20	6.60	123.65	116.33
4	B	804[B]	UTO	C9-N8-N7	6.35	107.92	101.73
4	A	804	UTO	C9-N8-N7	5.66	107.25	101.73
4	B	804[B]	UTO	N8-N7-N17	-5.62	109.24	114.01
4	B	804[A]	UTO	C9-N8-N7	5.50	107.08	101.73
4	B	804[A]	UTO	N8-N7-N17	-5.25	109.55	114.01
4	B	804[B]	UTO	N8-C9-N16	-5.08	107.02	111.99
4	A	804	UTO	N8-N7-N17	-5.02	109.74	114.01
4	B	804[A]	UTO	N8-C9-N16	-4.96	107.14	111.99
4	A	804	UTO	N8-C9-N16	-4.96	107.14	111.99
4	A	804	UTO	O1-C2-N20	-4.86	116.63	122.46
4	B	804[A]	UTO	C3-C2-N20	4.40	121.21	116.33
4	B	804[A]	UTO	C12-N11-C10	3.95	122.03	116.93
4	B	804[A]	UTO	N7-N17-N16	3.77	109.09	106.03
4	B	804[B]	UTO	C3-C2-N20	3.74	120.48	116.33
4	B	804[A]	UTO	C19-N18-C5	3.65	122.64	118.00
4	A	804	UTO	C19-N18-C5	3.62	122.61	118.00
4	B	804[B]	UTO	N7-N17-N16	3.62	108.97	106.03
4	B	804[A]	UTO	C10-C9-N8	3.58	128.18	123.38
4	B	804[B]	UTO	C12-N11-C10	3.49	121.42	116.93
4	A	804	UTO	N7-N17-N16	3.44	108.83	106.03
4	A	804	UTO	C12-N11-C10	3.37	121.28	116.93
4	B	804[B]	UTO	C19-N18-C5	3.08	121.92	118.00
4	B	804[B]	UTO	O1-C2-N20	-3.04	118.81	122.46
4	B	804[A]	UTO	C6-N7-N8	2.73	125.91	122.33
4	B	804[B]	UTO	C10-C9-N16	2.64	127.64	124.11
4	B	804[B]	UTO	C22-C4-C3	-2.47	118.16	120.80
4	A	804	UTO	C10-C9-N8	2.34	126.52	123.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	804[A]	UTO	O1-C2-N20	-2.25	119.75	122.46
4	A	804	UTO	C6-N7-N8	2.23	125.26	122.33
4	B	804[A]	UTO	C13-C12-N11	-2.16	120.00	123.42
4	B	804[A]	UTO	C3-C19-N18	-2.07	120.54	123.59
4	A	804	UTO	C13-C12-N11	-2.01	120.24	123.42

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	804[B]	UTO	C15-C10-C9-N8
4	B	804[B]	UTO	N11-C10-C9-N8
4	B	804[B]	UTO	C15-C10-C9-N16
4	B	804[B]	UTO	N11-C10-C9-N16
4	B	804[B]	UTO	N20-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	804[B]	UTO	1	0
4	A	804	UTO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	356/364 (97%)	0.11	14 (3%)	43 50	5, 13, 30, 58	7 (1%)
1	B	352/364 (96%)	0.35	21 (5%)	27 32	9, 18, 34, 50	5 (1%)
1	C	352/364 (96%)	1.61	117 (33%)	1 1	12, 40, 72, 104	3 (0%)
All	All	1060/1092 (97%)	0.69	152 (14%)	6 7	5, 20, 61, 104	15 (1%)

All (152) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	772	LEU	11.5
1	C	769	LEU	11.2
1	A	769	LEU	8.7
1	A	442	PRO	8.0
1	B	767	PRO	7.7
1	C	775	LEU	7.0
1	C	767	PRO	7.0
1	C	522	LEU	6.7
1	C	797	LEU	6.6
1	C	442	PRO	6.5
1	A	773	THR	6.1
1	A	441	SER	5.9
1	C	511	ILE	5.6
1	C	777	THR	5.5
1	C	443	ILE	5.5
1	B	773	THR	5.5
1	C	459	TRP	5.5
1	B	442	PRO	5.3
1	C	658	LYS	5.3
1	C	518	LYS	5.3
1	C	510	ILE	5.2
1	C	783	ILE	5.2
1	C	781	VAL	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	516	HIS	5.0
1	C	519	PRO	5.0
1	C	768	SER	4.9
1	C	671	GLY	4.7
1	B	672	LYS	4.6
1	C	798	ARG	4.6
1	C	525	LEU	4.4
1	C	527	ASP	4.4
1	C	528	GLU	4.3
1	C	521	ASP	4.1
1	C	524	ARG	4.1
1	C	523	ASN	4.0
1	C	512	LYS	4.0
1	A	672[A]	LYS	3.9
1	C	724	HIS	3.9
1	C	796	SER	3.9
1	C	682	PHE	3.8
1	C	520	ARG	3.7
1	C	493	LEU	3.7
1	C	579	THR	3.7
1	C	780	THR	3.7
1	C	578	ASP	3.7
1	C	788	ARG	3.7
1	C	672	LYS	3.6
1	C	776	LYS	3.6
1	C	562	ARG	3.6
1	B	671	GLY	3.6
1	C	514	SER	3.5
1	C	515	GLU	3.5
1	C	793	PHE	3.5
1	C	526	GLY	3.5
1	A	671[A]	GLY	3.4
1	C	481	ARG	3.4
1	C	651	TYR	3.4
1	C	691	ARG	3.4
1	B	443	ILE	3.3
1	C	539	TYR	3.3
1	A	443	ILE	3.2
1	C	505	SER	3.2
1	C	766	PRO	3.2
1	C	517	MET	3.2
1	C	454	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	656	LEU	3.1
1	C	787	LEU	3.1
1	B	777	THR	3.1
1	C	679	MET	3.1
1	C	534	ILE	3.1
1	C	725	LEU	3.1
1	C	458	MET	3.1
1	C	509[A]	SER	3.0
1	C	580	ALA	3.0
1	C	779	ALA	3.0
1	C	508	ILE	3.0
1	C	499	LEU	3.0
1	C	728	GLN	3.0
1	C	786	VAL	2.9
1	C	560	GLN	2.9
1	C	577	LYS	2.9
1	C	784	ASN	2.9
1	C	506	LYS	2.9
1	C	529	TYR	2.8
1	C	789	ALA	2.8
1	B	724	HIS	2.8
1	C	732	LEU	2.8
1	B	658	LYS	2.7
1	C	462	HIS	2.7
1	B	776	LYS	2.7
1	B	578	ASP	2.7
1	C	532	ILE	2.7
1	C	533	PHE	2.7
1	C	778	SER	2.6
1	C	795	SER	2.6
1	C	765	SER	2.6
1	C	683	HIS	2.6
1	C	684	HIS	2.6
1	B	728	GLN	2.6
1	C	689	ILE	2.5
1	B	506	LYS	2.5
1	C	660	ARG	2.5
1	C	500	ALA	2.5
1	C	668	TRP	2.5
1	C	712	LEU	2.5
1	C	715	PHE	2.5
1	C	624	GLU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	777	THR	2.5
1	A	658	LYS	2.4
1	C	507	HIS	2.4
1	B	562	ARG	2.4
1	C	685	LEU	2.4
1	C	792	PRO	2.4
1	B	794	TRP	2.3
1	C	794	TRP	2.3
1	C	654	VAL	2.3
1	C	790	HIS	2.3
1	C	585	PHE	2.3
1	A	767	PRO	2.3
1	C	708	ARG	2.2
1	B	660	ARG	2.2
1	C	690	ALA	2.2
1	C	676	PRO	2.2
1	C	461	SER	2.2
1	B	481	ARG	2.2
1	B	798	ARG	2.2
1	C	536	ASN	2.2
1	C	716	GLN	2.2
1	C	460	ASP	2.2
1	C	494	ALA	2.2
1	B	793	PHE	2.2
1	C	686	VAL	2.2
1	C	688	PRO	2.1
1	C	659	GLY	2.1
1	C	662	TYR	2.1
1	C	643	PHE	2.1
1	C	719	PRO	2.1
1	B	656	LEU	2.1
1	C	604	LEU	2.1
1	C	680	ALA	2.1
1	A	775	LEU	2.1
1	B	775	LEU	2.1
1	A	728	GLN	2.1
1	C	693	PHE	2.0
1	C	791	ALA	2.0
1	A	798	ARG	2.0
1	C	729	LEU	2.0
1	C	622	ILE	2.0
1	C	670	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	540	THR	2.0
1	C	625	GLU	2.0
1	C	583	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	C	803	1/1	0.75	0.25	89,89,89,89	0
4	UTO	B	804[A]	22/22	0.80	0.18	24,26,27,28	22
4	UTO	B	804[B]	22/22	0.80	0.18	4,22,36,36	22
4	UTO	A	804	22/22	0.89	0.14	12,25,40,42	0
5	IOD	A	808	1/1	0.95	0.24	46,46,46,46	0
5	IOD	B	805	1/1	0.95	0.24	71,71,71,71	0
5	IOD	C	804	1/1	0.96	0.20	46,46,46,46	0
3	ZN	B	803	1/1	0.97	0.31	51,51,51,51	0
5	IOD	A	806	1/1	0.99	0.04	15,15,15,15	0
5	IOD	A	807	1/1	0.99	0.12	28,28,28,28	0
3	ZN	A	803	1/1	0.99	0.29	41,41,41,41	0
2	K	C	802	1/1	0.99	0.07	53,53,53,53	0
5	IOD	A	805	1/1	0.99	0.09	24,24,24,24	0
2	K	B	802	1/1	1.00	0.02	16,16,16,16	0
2	K	C	801	1/1	1.00	0.02	31,31,31,31	0
2	K	A	801	1/1	1.00	0.01	9,9,9,9	0
2	K	A	802	1/1	1.00	0.01	13,13,13,13	0
2	K	B	801	1/1	1.00	0.01	12,12,12,12	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around UTO B 804 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UTO A 804:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.