



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:38 AM UTC

PDB ID : 3CKA / pdb_00003cka
Title : The crystal structure of OspA mutant
Authors : Makabe, K.; Biancalana, M.; Terechko, V.; Koide, S.
Deposited on : 2008-03-14
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	8% 84% 12% ..
1	B	320	6% 86% 11% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer surface protein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	4	10	0
			2446	1530	391	524	1			
1	B	316	Total	C	N	O	S	0	2	0
			2416	1508	390	517	1			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLY	-	expression tag	UNP D2Y4L7
A	24	SER	-	expression tag	UNP D2Y4L7
A	25	HIS	-	expression tag	UNP D2Y4L7
A	26	MET	-	expression tag	UNP D2Y4L7
A	37	SER	GLU	engineered mutation	UNP D2Y4L7
A	45	SER	GLU	engineered mutation	UNP D2Y4L7
A	46	SER	LYS	engineered mutation	UNP D2Y4L7
A	48	ALA	LYS	engineered mutation	UNP D2Y4L7
A	60	ALA	LYS	engineered mutation	UNP D2Y4L7
A	64	SER	LYS	engineered mutation	UNP D2Y4L7
A	83	ALA	LYS	engineered mutation	UNP D2Y4L7
A	104	SER	GLU	engineered mutation	UNP D2Y4L7
A	107	SER	LYS	engineered mutation	UNP D2Y4L7
A	123	TYR	GLU	engineered mutation	UNP D2Y4L7
A	125	LEU	LYS	engineered mutation	UNP D2Y4L7
A	132	LEU	VAL	engineered mutation	UNP D2Y4L7
A	134	PHE	GLU	engineered mutation	UNP D2Y4L7
A	136	TYR	ILE	engineered mutation	UNP D2Y4L7
A	308	SER	LYS	engineered mutation	UNP Q45040
A	309	SER	GLU	engineered mutation	UNP Q45040
A	323	SER	LYS	engineered mutation	UNP Q45040
B	23	GLY	-	expression tag	UNP D2Y4L7
B	24	SER	-	expression tag	UNP D2Y4L7
B	25	HIS	-	expression tag	UNP D2Y4L7
B	26	MET	-	expression tag	UNP D2Y4L7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	37	SER	GLU	engineered mutation	UNP D2Y4L7
B	45	SER	GLU	engineered mutation	UNP D2Y4L7
B	46	SER	LYS	engineered mutation	UNP D2Y4L7
B	48	ALA	LYS	engineered mutation	UNP D2Y4L7
B	60	ALA	LYS	engineered mutation	UNP D2Y4L7
B	64	SER	LYS	engineered mutation	UNP D2Y4L7
B	83	ALA	LYS	engineered mutation	UNP D2Y4L7
B	104	SER	GLU	engineered mutation	UNP D2Y4L7
B	107	SER	LYS	engineered mutation	UNP D2Y4L7
B	123	TYR	GLU	engineered mutation	UNP D2Y4L7
B	125	LEU	LYS	engineered mutation	UNP D2Y4L7
B	132	LEU	VAL	engineered mutation	UNP D2Y4L7
B	134	PHE	GLU	engineered mutation	UNP D2Y4L7
B	136	TYR	ILE	engineered mutation	UNP D2Y4L7
B	308	SER	LYS	engineered mutation	UNP Q45040
B	309	SER	GLU	engineered mutation	UNP Q45040
B	323	SER	LYS	engineered mutation	UNP Q45040

- Molecule 2 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	5	2		
2	A	1	Total	C	O	0	0
			16	10	6		
2	B	1	Total	C	O	0	0
			13	8	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			13	8	5		
2	B	1	Total	C	O	0	0
			13	8	5		
2	B	1	Total	C	O	0	0
			9	6	3		
2	B	1	Total	C	O	0	0
			9	6	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	270	Total	O	0	0
			270	270		
3	B	247	Total	O	0	0
			247	247		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Outer surface protein A



- Molecule 1: Outer surface protein A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.29Å 83.65Å 106.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.65 20.00 – 1.65	Depositor EDS
% Data completeness (in resolution range)	98.4 (20.00-1.65) 99.7 (20.00-1.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.63Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.202 , 0.240 0.221 , 0.257	Depositor DCC
R_{free} test set	4102 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtrriage
Anisotropy	0.199	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5459	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.14 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1943e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	0/2494	1.03	1/3357 (0.0%)
1	B	1.12	0/2446	1.01	1/3289 (0.0%)
All	All	1.09	0/4940	1.02	2/6646 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	331	ILE	CB-CA-C	6.70	120.99	110.81
1	B	37	SER	N-CA-C	-5.66	98.75	110.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	36	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2446	0	2496	39	0
1	B	2416	0	2449	39	0
2	A	23	0	28	1	0
2	B	57	0	71	1	0
3	A	270	0	0	7	0
3	B	247	0	0	10	0
All	All	5459	0	5044	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:THR:HG21	3:B:1104:HOH:O	1.65	0.97
1:A:122[B]:THR:HG22	1:A:137:ILE:HG12	1.61	0.83
1:B:231:LEU:HD13	3:B:1093:HOH:O	1.78	0.83
1:A:251:GLU:O	1:A:254[A]:VAL:HG12	1.78	0.82
1:B:168:THR:HG23	3:B:951:HOH:O	1.84	0.78
1:A:60:ALA:HB2	1:B:334:LEU:CD2	2.15	0.76
1:A:60:ALA:HB2	1:B:334:LEU:HD21	1.69	0.72
1:A:316:GLN:NE2	3:A:976:HOH:O	2.23	0.72
1:B:27:LYS:HG3	1:B:28:ASN:H	1.55	0.71
1:B:231:LEU:HD12	1:B:234:TYR:CE1	2.29	0.68
1:A:122[A]:THR:HG21	1:A:135:LYS:HE3	1.76	0.67
1:A:254[A]:VAL:HG11	1:A:341:LEU:HB3	1.75	0.67
1:A:39:LYS:HD3	3:A:1058:HOH:O	1.95	0.66
1:B:51:LYS:NZ	1:B:68:ASP:OD1	2.28	0.66
1:B:28:ASN:O	1:B:44:LYS:HE2	1.97	0.64
1:A:226:LYS:HE3	3:A:1070:HOH:O	1.98	0.63
1:A:38:MET:HE1	1:A:99:LEU:HD22	1.81	0.63
1:B:158:LYS:HE2	1:B:160:ILE:HD11	1.80	0.63
1:A:182:TYR:CE2	1:A:184[B]:THR:HG23	2.36	0.61
1:B:262:LYS:CD	3:B:1168:HOH:O	2.50	0.58
1:B:240:LEU:HD22	1:B:241:THR:O	2.03	0.58
1:B:27:LYS:CG	1:B:28:ASN:H	2.16	0.58
1:A:28:ASN:N	1:A:28:ASN:HD22	2.01	0.58
1:B:27:LYS:HG3	1:B:28:ASN:N	2.18	0.57
1:A:122[A]:THR:HG21	1:A:135:LYS:CE	2.36	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248[B]:VAL:HG13	1:A:255:THR:HG23	1.88	0.56
1:B:232:LYS:HD3	3:B:1169:HOH:O	2.06	0.56
1:A:59:ASP:CG	1:B:235:VAL:HG21	2.31	0.55
1:A:185:ARG:NH1	3:A:993:HOH:O	2.39	0.55
1:A:137:ILE:HD12	3:A:998:HOH:O	2.05	0.55
1:B:181:LYS:HD3	3:B:1156:HOH:O	2.08	0.53
1:A:197:GLU:CD	1:B:129:LYS:HZ3	2.17	0.53
1:B:168:THR:HB	1:B:183:ILE:HG12	1.90	0.53
1:A:216:TYR:CE2	2:A:802:1PE:H151	2.44	0.53
1:A:194:LEU:HD21	3:B:1116:HOH:O	2.08	0.52
1:A:179:SER:HB2	1:B:151:GLU:OE1	2.10	0.52
1:B:262:LYS:HD2	3:B:1168:HOH:O	2.07	0.51
1:B:136:TYR:CZ	1:B:138:THR:CG2	2.95	0.50
1:A:32:VAL:HG22	1:A:40:VAL:CG1	2.43	0.49
1:A:162:ARG:NH1	3:A:1029:HOH:O	2.37	0.48
1:A:32:VAL:HG22	1:A:40:VAL:HG12	1.94	0.48
1:A:248[B]:VAL:HG13	1:A:255:THR:CG2	2.44	0.48
1:B:33:ASP:OD1	1:B:39:LYS:NZ	2.42	0.48
1:B:156:SER:O	1:B:171:LEU:HD12	2.14	0.48
1:A:170:GLU:OE2	1:A:181:LYS:HE3	2.15	0.47
1:B:136:TYR:CE2	1:B:138:THR:CG2	2.98	0.47
1:A:180:PHE:C	1:A:180:PHE:CD2	2.93	0.47
1:B:136:TYR:CE2	1:B:138:THR:HG23	2.50	0.47
1:A:244:LYS:NZ	1:A:259:ASN:ND2	2.63	0.46
1:A:302:LYS:HG2	1:A:316:GLN:NE2	2.30	0.46
1:B:52:TYR:CZ	1:B:70:ASN:HB3	2.51	0.46
1:A:171:LEU:HD23	1:A:179:SER:OG	2.16	0.45
1:B:241:THR:HG1	1:B:243:GLU:CD	2.24	0.45
1:B:211:GLY:O	1:B:213:ARG:HD2	2.17	0.45
1:B:243:GLU:O	1:B:244:LYS:HB3	2.17	0.45
1:A:140:ALA:C	1:A:142:LYS:N	2.76	0.44
2:B:805:1PE:H252	3:B:978:HOH:O	2.17	0.44
1:B:27:LYS:CG	1:B:28:ASN:N	2.79	0.44
1:A:28:ASN:N	1:A:28:ASN:ND2	2.64	0.44
1:B:228:LYS:NZ	1:B:237:GLU:OE2	2.51	0.43
1:B:117:LYS:C	1:B:119:LYS:H	2.26	0.43
1:A:87:LYS:NZ	3:A:899:HOH:O	2.34	0.43
1:B:117:LYS:C	1:B:119:LYS:N	2.76	0.43
1:B:180:PHE:C	1:B:180:PHE:CD2	2.97	0.43
1:A:182:TYR:CE2	1:A:184[B]:THR:CG2	3.01	0.42
1:B:28:ASN:ND2	1:B:44:LYS:HD2	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:TYR:CZ	1:B:138:THR:HG22	2.53	0.42
1:A:143:SER:HB3	1:A:162:ARG:HA	2.02	0.42
1:A:60:ALA:HB2	1:B:334:LEU:HD23	1.98	0.41
1:B:129:LYS:HD3	3:B:1078:HOH:O	2.21	0.40
1:A:140:ALA:O	1:A:142:LYS:N	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/320 (101%)	315 (98%)	6 (2%)	2 (1%)	21	8
1	B	316/320 (99%)	307 (97%)	9 (3%)	0	100	100
All	All	639/640 (100%)	622 (97%)	15 (2%)	2 (0%)	36	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	ASP
1	A	142	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/279 (102%)	271 (95%)	14 (5%)	22	5
1	B	278/279 (100%)	269 (97%)	9 (3%)	34	12
All	All	563/558 (101%)	540 (96%)	23 (4%)	28	6

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	32	VAL
1	A	54	LEU
1	A	61	LEU
1	A	85	LYS
1	A	142	LYS
1	A	143	SER
1	A	148	LEU
1	A	152	LYS
1	A	174	GLU
1	A	184[A]	THR
1	A	184[B]	THR
1	A	185	ARG
1	A	302	LYS
1	B	117	LYS
1	B	119	LYS
1	B	129	LYS
1	B	152	LYS
1	B	168	THR
1	B	174	GLU
1	B	240	LEU
1	B	262	LYS
1	B	263	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	259	ASN
1	A	297	ASN
1	A	316	GLN
1	A	339	ASN
1	B	297	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1PE	A	800	-	6,6,15	0.51	0	5,5,14	1.60	1 (20%)
2	1PE	B	804	-	12,12,15	0.58	0	11,11,14	1.56	4 (36%)
2	1PE	B	805	-	8,8,15	0.57	0	7,7,14	1.37	1 (14%)
2	1PE	B	806	-	8,8,15	0.47	0	7,7,14	1.37	1 (14%)
2	1PE	B	801	-	12,12,15	0.59	0	11,11,14	1.54	3 (27%)
2	1PE	B	803	-	12,12,15	0.61	0	11,11,14	1.52	3 (27%)
2	1PE	A	802	-	15,15,15	0.52	0	14,14,14	1.45	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	A	800	-	-	2/4/4/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	B	804	-	-	5/10/10/13	-
2	1PE	B	805	-	-	3/6/6/13	-
2	1PE	B	806	-	-	3/6/6/13	-
2	1PE	B	801	-	-	3/10/10/13	-
2	1PE	B	803	-	-	5/10/10/13	-
2	1PE	A	802	-	-	7/13/13/13	-

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	802	1PE	C25-OH5-C14	3.21	127.31	113.26
2	A	800	1PE	C26-OH6-C15	3.00	123.57	113.06
2	B	806	1PE	C26-OH6-C15	2.51	121.85	113.06
2	B	805	1PE	OH5-C25-C15	2.35	121.08	110.35
2	B	804	1PE	OH4-C24-C14	2.34	121.00	110.35
2	B	804	1PE	OH5-C25-C15	2.29	120.80	110.35
2	B	801	1PE	OH6-C26-C16	2.27	120.13	110.11
2	B	803	1PE	OH6-C26-C16	2.25	120.02	110.11
2	B	803	1PE	OH5-C25-C15	2.16	120.20	110.35
2	B	801	1PE	OH4-C24-C14	2.14	120.13	110.35
2	B	804	1PE	C24-OH4-C13	2.11	122.49	113.26
2	A	802	1PE	C23-OH3-C22	2.10	122.44	113.26
2	A	802	1PE	OH2-C12-C22	-2.09	99.50	111.82
2	B	803	1PE	OH4-C24-C14	2.08	119.83	110.35
2	B	804	1PE	OH6-C26-C16	2.08	119.27	110.11
2	B	801	1PE	OH5-C25-C15	2.02	119.57	110.35

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	806	1PE	OH6-C15-C25-OH5
2	A	802	1PE	OH6-C15-C25-OH5
2	A	800	1PE	OH6-C15-C25-OH5
2	B	803	1PE	OH6-C15-C25-OH5
2	B	804	1PE	OH7-C16-C26-OH6
2	B	804	1PE	OH4-C13-C23-OH3
2	A	802	1PE	OH7-C16-C26-OH6
2	B	806	1PE	C16-C26-OH6-C15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	802	1PE	C25-C15-OH6-C26
2	B	805	1PE	C15-C25-OH5-C14
2	B	805	1PE	C24-C14-OH5-C25
2	B	806	1PE	C25-C15-OH6-C26
2	A	802	1PE	C16-C26-OH6-C15
2	B	801	1PE	C16-C26-OH6-C15
2	A	802	1PE	OH5-C14-C24-OH4
2	B	804	1PE	C15-C25-OH5-C14
2	B	803	1PE	C14-C24-OH4-C13
2	A	802	1PE	C24-C14-OH5-C25
2	A	800	1PE	C16-C26-OH6-C15
2	B	805	1PE	C25-C15-OH6-C26
2	B	801	1PE	OH5-C14-C24-OH4
2	B	803	1PE	OH7-C16-C26-OH6
2	B	803	1PE	C16-C26-OH6-C15
2	B	801	1PE	C14-C24-OH4-C13
2	B	803	1PE	OH5-C14-C24-OH4
2	A	802	1PE	C12-C22-OH3-C23
2	B	804	1PE	C16-C26-OH6-C15
2	B	804	1PE	OH5-C14-C24-OH4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	805	1PE	1	0
2	A	802	1PE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	315/320 (98%)	0.51	25 (7%) 18 21	11, 24, 39, 50	10 (3%)
1	B	316/320 (98%)	0.70	19 (6%) 27 31	15, 27, 39, 47	2 (0%)
All	All	631/640 (98%)	0.61	44 (6%) 22 25	11, 26, 39, 50	12 (1%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	186	ALA	4.8
1	A	140	ALA	4.3
1	B	186	ALA	4.0
1	A	29	SER	4.0
1	A	60	ALA	3.7
1	B	28	ASN	3.6
1	A	36	GLY	3.5
1	A	118	ASP	3.3
1	A	141	ASP	3.2
1	A	117	LYS	3.1
1	A	154	GLU	3.1
1	A	192	TYR	3.1
1	B	174	GLU	3.1
1	B	243	GLU	3.0
1	A	120	SER	3.0
1	A	28	ASN	3.0
1	B	36	GLY	2.9
1	A	163	ALA	2.9
1	B	140	ALA	2.8
1	A	151	GLU	2.8
1	A	136	TYR	2.7
1	B	29	SER	2.7
1	B	192	TYR	2.6
1	A	139	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	132	LEU	2.5
1	A	137	ILE	2.5
1	B	197	GLU	2.5
1	A	121	SER	2.4
1	B	241	THR	2.3
1	B	118	ASP	2.3
1	A	142	LYS	2.3
1	A	138	THR	2.2
1	B	149	PHE	2.2
1	A	152	LYS	2.2
1	A	37	SER	2.1
1	A	114	VAL	2.1
1	A	243	GLU	2.1
1	B	137	ILE	2.1
1	B	60	ALA	2.1
1	B	242	ALA	2.1
1	B	155	LEU	2.1
1	B	154	GLU	2.0
1	B	176	GLY	2.0
1	A	119	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	1PE	B	801	13/16	0.72	0.16	53,55,56,56	0
2	1PE	B	805	9/16	0.73	0.15	56,61,62,63	0
2	1PE	B	806	9/16	0.75	0.17	53,54,56,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	1PE	B	804	13/16	0.77	0.13	54,59,61,62	0
2	1PE	B	803	13/16	0.79	0.13	59,61,64,64	0
2	1PE	A	802	16/16	0.81	0.13	31,37,48,48	0
2	1PE	A	800	7/16	0.83	0.15	47,47,50,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.