



Full wwPDB EM Validation Report ⓘ

Mar 8, 2026 – 02:45 AM UTC

PDB ID : 4CK6 / pdb_00004ck6
EMDB ID : EMD-2533
Title : Pseudo-atomic model of microtubule-bound human kinesin-5 motor domain in the ADP.AIFx state, based on cryo-electron microscopy experiment.
Authors : Goulet, A.; Major, J.; Jun, Y.; Gross, S.; Rosenfeld, S.; Moores, C.
Deposited on : 2013-12-30
Resolution : 9.20 Å (reported)
Based on initial models : 3HQD, 1JFF

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY



The reported resolution of this entry is 9.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Q-score	-	25397	239 (8.70 - 9.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	
2	B	445	
3	C	373	

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 1354 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TUBULIN ALPHA-1D CHAIN.

Mol	Chain	Residues	Atoms	AltConf	Trace
1	A	412	Total C 412 412	0	412

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ILE	VAL	conflict	UNP Q2HJ86
A	114	ILE	LEU	conflict	UNP Q2HJ86
A	136	SER	LEU	conflict	UNP Q2HJ86
A	137	VAL	ILE	conflict	UNP Q2HJ86
A	265	GLY	ILE	conflict	UNP Q2HJ86
A	358	GLU	ASP	conflict	UNP Q2HJ86
A	437	VAL	MET	conflict	UNP Q2HJ86
A	450	GLU	ASP	conflict	UNP Q2HJ86

- Molecule 2 is a protein called TUBULIN BETA-2B CHAIN.

Mol	Chain	Residues	Atoms	AltConf	Trace
2	B	426	Total C 426 426	0	426

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	55	ALA	THR	conflict	UNP Q6B856
B	170	VAL	MET	conflict	UNP Q6B856
B	296	ALA	SER	conflict	UNP Q6B856
B	316	VAL	ILE	conflict	UNP Q6B856

- Molecule 3 is a protein called KINESIN-LIKE PROTEIN KIF11.

Mol	Chain	Residues	Atoms	AltConf	Trace
3	C	361	Total C 361 361	0	361


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	25	VAL	CYS	conflict	UNP P52732
C	43	SER	CYS	conflict	UNP P52732
C	87	ALA	CYS	conflict	UNP P52732
C	99	ALA	CYS	conflict	UNP P52732
C	126	CYS	THR	engineered mutation	UNP P52732
C	368	HIS	-	expression tag	UNP P52732
C	369	HIS	-	expression tag	UNP P52732
C	370	HIS	-	expression tag	UNP P52732
C	371	HIS	-	expression tag	UNP P52732
C	372	HIS	-	expression tag	UNP P52732
C	373	HIS	-	expression tag	UNP P52732

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	AltConf
4	A	1	Total Mg 1 1	0
4	C	1	Total Mg 1 1	0

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).




Mol	Chain	Residues	Atoms					AltConf
5	A	1	Total	C	N	O	P	0
			32	10	5	14	3	

- Molecule 6 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues	Atoms				AltConf
7	B	1	Total	C	N	O	0
			62	47	1	14	

- Molecule 8 is ALUMINUM FLUORIDE (CCD ID: AF3) (formula: AlF_3).



Mol	Chain	Residues	Atoms			AltConf
8	C	1	Total	Al	F	0
			4	1	3	

- Molecule 9 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{10}\text{P}_2$).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
9	C	1	27	10	5	10	2	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: TUBULIN ALPHA-1D CHAIN




• Molecule 2: TUBULIN BETA-2B CHAIN



• Molecule 3: KINESIN-LIKE PROTEIN KIF11





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of particles used	10692	Depositor
Resolution determination method	Not provided	
CTF correction method	FREALIGN	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	18	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	50000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	9.523	Depositor
Minimum map value	-7.905	Depositor
Average map value	0.291	Depositor
Map value standard deviation	1.946	Depositor
Recommended contour level	1.1	Depositor
Map size (Å)	103.6, 103.6, 103.6	wwPDB
Map dimensions	37, 37, 37	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.8, 2.8, 2.8	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AF3, GDP, ADP, GTP, TA1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	412	0	0	2	0
2	B	426	0	0	4	0
3	C	361	0	0	2	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	32	0	12	0	0
6	B	28	0	12	0	0
7	B	62	0	51	3	0
8	C	4	0	0	0	0
9	C	27	0	12	0	0
All	All	1354	0	87	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436:GLY:CA	1:A:437:VAL:CA	2.46	0.94
2:B:243:PRO:CA	2:B:244:GLY:CA	2.55	0.84
3:C:267:ALA:CA	3:C:268:GLY:CA	2.67	0.71
2:B:95:SER:CA	2:B:96:GLY:CA	2.75	0.65
7:B:1211:TA1:H463	7:B:1211:TA1:H261	1.80	0.63
1:A:402:ARG:CA	1:A:403:ALA:CA	2.80	0.60
3:C:311:TYR:CA	3:C:312:ARG:CA	2.81	0.58
2:B:339:SER:CA	2:B:340:TYR:CA	2.93	0.47
7:B:1211:TA1:H463	7:B:1211:TA1:C26	2.46	0.44
2:B:360:GLY:CA	7:B:1211:TA1:H443	2.47	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
9	ADP	C	1207	8,4	28,29,29	1.38	3 (10%)	43,45,45	1.30	6 (13%)
6	GDP	B	1210	-	29,30,30	2.87	11 (37%)	45,47,47	3.48	16 (35%)
8	AF3	C	1206	9,4	0,3,3	-	-	-	-	-
7	TA1	B	1211	-	68,68,68	2.20	26 (38%)	105,105,105	1.50	14 (13%)
5	GTP	A	1209	4	33,34,34	1.90	3 (9%)	50,54,54	0.91	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	ADP	C	1207	8,4	-	5/16/32/32	0/3/3/3
6	GDP	B	1210	-	-	4/16/32/32	0/3/3/3
5	GTP	A	1209	4	-	4/22/38/38	0/3/3/3
7	TA1	B	1211	-	-	9/41/127/127	0/7/7/7

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1210	GDP	PA-O3A	7.57	1.67	1.59
7	B	1211	TA1	C06-C05	6.46	1.49	1.38
5	A	1209	GTP	PA-O3A	-5.99	1.53	1.59
6	B	1210	GDP	O6-C6	5.99	1.34	1.23
6	B	1210	GDP	C8-N9	5.82	1.50	1.37
7	B	1211	TA1	C08-C07	-5.71	1.25	1.38
5	A	1209	GTP	PB-O3B	5.63	1.65	1.59
5	A	1209	GTP	PB-O3A	-5.62	1.53	1.59
7	B	1211	TA1	C18-C10	5.37	1.69	1.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	1211	TA1	C05-C04	4.83	1.46	1.39
6	B	1210	GDP	C2-N1	4.78	1.49	1.37
7	B	1211	TA1	C45-C24	4.29	1.61	1.54
6	B	1210	GDP	PB-O2B	-3.92	1.40	1.54
7	B	1211	TA1	C36-C31	3.73	1.45	1.39
7	B	1211	TA1	O02-C03	3.51	1.41	1.34
7	B	1211	TA1	C25-C24	3.39	1.39	1.34
6	B	1210	GDP	O4'-C1'	3.34	1.49	1.42
7	B	1211	TA1	C46-C45	3.22	1.60	1.53
7	B	1211	TA1	C43-C01	3.21	1.60	1.54
7	B	1211	TA1	C11-C10	3.06	1.61	1.54
6	B	1210	GDP	C8-N7	3.04	1.41	1.32
6	B	1210	GDP	C5-N7	-2.95	1.33	1.39
9	C	1207	ADP	C2-N1	2.87	1.39	1.33
7	B	1211	TA1	C43-C26	2.81	1.58	1.52
6	B	1210	GDP	C5-C4	2.69	1.46	1.38
9	C	1207	ADP	C8-N7	2.64	1.36	1.31
7	B	1211	TA1	C26-C25	2.57	1.56	1.51
6	B	1210	GDP	C2-N3	-2.56	1.26	1.33
9	C	1207	ADP	C4-N3	2.56	1.39	1.34
7	B	1211	TA1	C01-C45	2.51	1.66	1.56
7	B	1211	TA1	C18-C20	2.50	1.62	1.55
7	B	1211	TA1	C04-C03	-2.44	1.44	1.50
6	B	1210	GDP	PB-O3B	2.42	1.63	1.54
7	B	1211	TA1	C41-C42	2.30	1.42	1.38
7	B	1211	TA1	C16-C15	2.28	1.56	1.52
7	B	1211	TA1	C37-C29	2.20	1.54	1.52
7	B	1211	TA1	C08-C09	2.20	1.42	1.38
7	B	1211	TA1	O09-C21	2.18	1.49	1.44
7	B	1211	TA1	C33-C32	2.17	1.42	1.38
7	B	1211	TA1	C10-C02	2.14	1.62	1.57
7	B	1211	TA1	C39-C38	2.10	1.42	1.38
7	B	1211	TA1	C07-C06	2.06	1.42	1.38
7	B	1211	TA1	C34-C33	2.03	1.42	1.38

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1210	GDP	N9-C8-N7	-10.72	93.51	113.40
6	B	1210	GDP	C8-N7-C5	9.21	120.67	104.26
6	B	1210	GDP	C6-C5-N7	8.62	145.97	130.29
6	B	1210	GDP	N2-C2-N3	6.33	132.03	119.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1210	GDP	C6-C5-C4	-6.25	109.43	118.83
7	B	1211	TA1	C06-C05-C04	-5.89	114.57	120.36
6	B	1210	GDP	C8-N9-C4	5.85	116.99	106.03
7	B	1211	TA1	C07-C08-C09	5.75	127.33	120.24
6	B	1210	GDP	C5-C6-N1	4.51	124.72	113.25
9	C	1207	ADP	O3B-PB-O3A	4.44	119.52	104.64
6	B	1210	GDP	N2-C2-N1	-4.30	107.69	116.76
6	B	1210	GDP	C2-N3-C4	4.18	119.49	112.30
7	B	1211	TA1	C05-C04-C03	-4.06	111.45	120.40
6	B	1210	GDP	O6-C6-C5	-4.00	115.97	126.53
6	B	1210	GDP	C4-C5-N7	-3.84	104.59	110.67
6	B	1210	GDP	C2-N1-C6	-3.81	118.20	125.11
7	B	1211	TA1	C09-C04-C03	3.59	128.31	120.40
6	B	1210	GDP	C2'-C3'-C4'	3.40	109.18	102.61
7	B	1211	TA1	C17-C18-C20	3.23	109.84	102.58
7	B	1211	TA1	O04-C11-C14	-3.04	101.69	108.13
7	B	1211	TA1	C45-C01-C02	2.99	115.22	111.93
9	C	1207	ADP	O3'-C3'-C2'	2.69	120.44	111.82
5	A	1209	GTP	O2G-PG-O3B	2.66	113.56	104.64
6	B	1210	GDP	C1'-N9-C8	-2.65	119.20	126.73
9	C	1207	ADP	O4'-C1'-N9	2.55	112.99	108.09
9	C	1207	ADP	C3'-C2'-C1'	2.50	106.18	101.46
7	B	1211	TA1	O01-C01-C43	2.48	113.37	107.04
7	B	1211	TA1	C08-C09-C04	-2.42	117.98	120.36
9	C	1207	ADP	O3A-PB-O1B	-2.42	98.32	111.04
9	C	1207	ADP	C4'-O4'-C1'	2.38	114.71	109.47
6	B	1210	GDP	O2'-C2'-C3'	2.28	119.11	111.82
7	B	1211	TA1	C14-C11-C15	-2.26	83.04	85.38
7	B	1211	TA1	C10-C18-C17	-2.26	102.26	106.55
7	B	1211	TA1	O06-C15-C11	2.21	93.00	90.58
6	B	1210	GDP	C4'-O4'-C1'	2.09	114.07	109.47
5	A	1209	GTP	O5'-C5'-C4'	2.08	116.07	108.99
5	A	1209	GTP	O3G-PG-O3B	2.04	111.49	104.64
7	B	1211	TA1	O04-C11-C10	2.01	112.36	109.24
7	B	1211	TA1	C21-O09-C22	2.01	120.10	115.95

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	1210	GDP	PA-O3A-PB-O2B
6	B	1210	GDP	C5'-O5'-PA-O3A

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
6	B	1210	GDP	C5'-O5'-PA-O1A
9	C	1207	ADP	PA-O3A-PB-O3B
9	C	1207	ADP	C5'-O5'-PA-O1A
9	C	1207	ADP	C5'-O5'-PA-O2A
9	C	1207	ADP	C5'-O5'-PA-O3A
7	B	1211	TA1	O02-C03-C04-C05
7	B	1211	TA1	O02-C03-C04-C09
7	B	1211	TA1	O03-C03-C04-C09
7	B	1211	TA1	O03-C03-C04-C05
7	B	1211	TA1	N01-C30-C31-C36
7	B	1211	TA1	O14-C30-C31-C36
7	B	1211	TA1	N01-C30-C31-C32
7	B	1211	TA1	O14-C30-C31-C32
5	A	1209	GTP	C3'-C4'-C5'-O5'
5	A	1209	GTP	O4'-C4'-C5'-O5'
5	A	1209	GTP	C5'-O5'-PA-O1A
9	C	1207	ADP	PA-O3A-PB-O1B
6	B	1210	GDP	PA-O3A-PB-O3B
7	B	1211	TA1	C15-C11-O04-C12
5	A	1209	GTP	PG-O3B-PB-O2B


There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	1211	TA1	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-2533. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 18



Y Index: 18



Z Index: 18


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 30



Y Index: 13




Z Index: 20


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 1.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 267 nm³; this corresponds to an approximate mass of 241 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.109 Å⁻¹

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-2533 and PDB model 4CK6. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (1.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8340	0.1060
A	0.8360	0.1060
B	0.8140	0.1030
C	0.8580	0.1100

