



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 03:07 AM UTC

PDB ID : 6CK9 / pdb_00006ck9
Title : Crystal Structure of HIV-1 ConC_Base0 Prefusion Env Trimer in Complex with Human Antibody Fragment 3H109L and 35O22 variants at 3.5 Angstrom
Authors : Lai, Y.-T.; Kwong, P.D.
Deposited on : 2018-02-27
Resolution : 2.71 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4348 (2.74-2.70)
Clashscore	190562	4665 (2.74-2.70)
Ramachandran outliers	187476	4584 (2.74-2.70)
Sidechain outliers	187428	4585 (2.74-2.70)
RSRZ outliers	180081	4348 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	153	14% 65% 20% 16%
2	D	134	40% 74% 19% . .
3	E	114	33% 69% 18% . 12%
4	G	463	16% 63% 29% . 7%
5	H	244	30% 74% 16% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	L	217	
7	A	5	
8	C	6	
9	F	4	
10	I	2	
10	K	2	
10	M	2	
10	O	2	
10	P	2	
11	J	2	
12	N	10	

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 10056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called gp41 ectodomain of Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	129	Total	C	N	O	S	0	0	0
			1030	655	173	196	6			

- Molecule 2 is a protein called 35O22 scFv heavy chain portion.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	128	Total	C	N	O	S	0	0	0
			994	630	168	191	5			

- Molecule 3 is a protein called 35O22 scFv light chain portion.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	100	Total	C	N	O	S	0	0	0
			762	480	125	151	6			

- Molecule 4 is a protein called gp120 of Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	G	430	Total	C	N	O	S	0	0	0
			3403	2140	597	639	27			

- Molecule 5 is a protein called 3H109L Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	H	219	Total	C	N	O	S	0	0	0
			1670	1065	271	328	6			

- Molecule 6 is a protein called 3H109L Fab light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	L	211	Total	C	N	O	S	0	0	0
			1604	1009	276	312	7			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	A	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	C	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	F	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 10 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
10	K	2	Total	C	N	O	0	0	0
			28	16	2	10			
10	M	2	Total	C	N	O	0	0	0
			28	16	2	10			
10	O	2	Total	C	N	O	0	0	0
			28	16	2	10			
10	P	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 11 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
11	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 12 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
12	N	10	Total	C	N	O	0	0	0
			116	64	2	50			

- Molecule 13 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	B	1	Total	C	N	O	0	0
			14	8	1	5		
13	B	1	Total	C	N	O	0	0
			14	8	1	5		
13	G	1	Total	C	N	O	0	0
			14	8	1	5		
13	G	1	Total	C	N	O	0	0
			14	8	1	5		
13	G	1	Total	C	N	O	0	0
			14	8	1	5		
13	G	1	Total	C	N	O	0	0
			14	8	1	5		
13	G	1	Total	C	N	O	0	0
			14	8	1	5		
13	G	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: gp41 ectodomain of Envelope glycoprotein gp160



- Molecule 2: 35O22 scFv heavy chain portion




- Molecule 3: 35O22 scFv light chain portion




- Molecule 4: gp120 of Envelope glycoprotein gp160








• Molecule 5: 3H109L Fab heavy chain



• Molecule 6: 3H109L Fab light chain





• Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain A:  20% 80%



- Molecule 8: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  17% 50% 33%




- Molecule 9: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%




- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%



- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%



- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  100%



- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:  100%

MAG1
MAG2

- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P:  100%

MAG1
MAG2

- Molecule 11: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

MAG1
MAG2

- Molecule 12: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  20% 70% 10%

MAG1
MAG2
BGA3
MAN4
MAN5
MAN6
MAN7
MAN8
MAN9
MAN10

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	131.90Å 131.90Å 314.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.17 – 2.71 43.17 – 2.71	Depositor EDS
% Data completeness (in resolution range)	35.5 (43.17-2.71) 35.5 (43.17-2.71)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.73Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.248 , 0.288 0.248 , 0.287	Depositor DCC
R_{free} test set	1472 reflections (1.76%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 24.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.076 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.82	EDS
Total number of atoms	10056	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.09	0/1049	0.28	0/1423
2	D	0.09	0/1021	0.27	0/1390
3	E	0.08	0/784	0.28	0/1070
4	G	0.10	0/3471	0.29	0/4708
5	H	0.09	0/1711	0.29	0/2330
6	L	0.08	0/1647	0.28	0/2247
All	All	0.09	0/9683	0.29	0/13168

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1030	0	1005	20	0
2	D	994	0	958	17	0
3	E	762	0	712	8	0
4	G	3403	0	3355	81	0
5	H	1670	0	1637	25	0
6	L	1604	0	1553	22	0
7	A	61	0	52	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	72	0	61	3	0
9	F	50	0	43	1	0
10	I	28	0	25	0	0
10	K	28	0	25	0	0
10	M	28	0	25	0	0
10	O	28	0	25	0	0
10	P	28	0	25	0	0
11	J	28	0	25	1	0
12	N	116	0	97	2	0
13	B	28	0	26	0	0
13	G	98	0	91	3	0
All	All	10056	0	9740	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:G:378:CYS:SG	4:G:379:ARG:N	2.52	0.83
4:G:71:THR:HG21	4:G:213:ILE:HD11	1.67	0.76
4:G:169:LYS:HB2	13:G:1601:NAG:H82	1.68	0.74
6:L:46:LEU:HD21	6:L:49:TYR:HB3	1.68	0.74
2:D:35:ASN:HB2	2:D:93:ALA:HB3	1.71	0.73

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	125/153 (82%)	114 (91%)	9 (7%)	2 (2%)	7 18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	126/134 (94%)	110 (87%)	16 (13%)	0	100	100
3	E	96/114 (84%)	82 (85%)	13 (14%)	1 (1%)	12	30
4	G	418/463 (90%)	359 (86%)	57 (14%)	2 (0%)	24	46
5	H	213/244 (87%)	194 (91%)	17 (8%)	2 (1%)	14	33
6	L	209/217 (96%)	191 (91%)	16 (8%)	2 (1%)	12	30
All	All	1187/1325 (90%)	1050 (88%)	128 (11%)	9 (1%)	16	35

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	L	51	ASN
6	L	25	ARG
5	H	144	ASP
1	B	548	ILE
1	B	625	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	111/129 (86%)	108 (97%)	3 (3%)	39	68
2	D	107/112 (96%)	103 (96%)	4 (4%)	30	57
3	E	87/100 (87%)	81 (93%)	6 (7%)	14	33
4	G	389/421 (92%)	367 (94%)	22 (6%)	18	41
5	H	190/212 (90%)	189 (100%)	1 (0%)	81	91
6	L	175/181 (97%)	172 (98%)	3 (2%)	53	77
All	All	1059/1155 (92%)	1020 (96%)	39 (4%)	30	57

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	G	331	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	H	63	LEU
4	G	350	LYS
4	G	377	ASN
6	L	160	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
4	G	195	ASN
5	H	31	ASN
6	L	184	GLN
5	H	200	HIS
6	L	66(C)	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

37 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	A	1	7,4	14,14,15	0.94	1 (7%)	17,19,21	2.03	3 (17%)
7	NAG	A	2	7	14,14,15	0.29	0	17,19,21	0.57	0
7	BMA	A	3	7	11,11,12	0.64	0	15,15,17	0.92	0
7	MAN	A	4	7	11,11,12	0.70	0	15,15,17	0.87	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	MAN	A	5	7	11,11,12	0.66	0	15,15,17	1.06	2 (13%)
8	NAG	C	1	8,4	14,14,15	0.33	0	17,19,21	0.45	0
8	NAG	C	2	8	14,14,15	0.48	0	17,19,21	1.34	2 (11%)
8	BMA	C	3	8	11,11,12	0.58	0	15,15,17	0.72	0
8	MAN	C	4	8	11,11,12	0.58	0	15,15,17	1.11	2 (13%)
8	MAN	C	5	8	11,11,12	0.75	0	15,15,17	1.25	2 (13%)
8	MAN	C	6	8	11,11,12	0.71	0	15,15,17	0.96	2 (13%)
9	NAG	F	1	9,4	14,14,15	0.22	0	17,19,21	0.46	0
9	NAG	F	2	9	14,14,15	0.26	0	17,19,21	0.43	0
9	BMA	F	3	9	11,11,12	0.60	0	15,15,17	0.82	0
9	MAN	F	4	9	11,11,12	0.62	0	15,15,17	0.96	2 (13%)
10	NAG	I	1	10,4	14,14,15	0.36	0	17,19,21	0.56	0
10	NAG	I	2	10	14,14,15	0.31	0	17,19,21	0.37	0
11	NAG	J	1	11,4	14,14,15	0.41	0	17,19,21	0.46	0
11	NAG	J	2	11	14,14,15	0.29	0	17,19,21	0.55	0
10	NAG	K	1	10,4	14,14,15	0.23	0	17,19,21	0.43	0
10	NAG	K	2	10	14,14,15	0.31	0	17,19,21	0.49	0
10	NAG	M	1	10,4	14,14,15	0.21	0	17,19,21	0.40	0
10	NAG	M	2	10	14,14,15	0.25	0	17,19,21	0.47	0
12	NAG	N	1	12,4	14,14,15	0.28	0	17,19,21	0.62	0
12	MAN	N	10	12	11,11,12	0.71	0	15,15,17	0.93	2 (13%)
12	NAG	N	2	12	14,14,15	0.22	0	17,19,21	0.44	0
12	BMA	N	3	12	11,11,12	0.70	0	15,15,17	0.99	0
12	MAN	N	4	12	11,11,12	0.74	1 (9%)	15,15,17	1.26	2 (13%)
12	MAN	N	5	12	11,11,12	0.62	0	15,15,17	1.03	2 (13%)
12	MAN	N	6	12	11,11,12	1.05	2 (18%)	15,15,17	1.90	4 (26%)
12	MAN	N	7	12	11,11,12	1.09	0	15,15,17	1.24	3 (20%)
12	MAN	N	8	12	11,11,12	0.91	0	15,15,17	1.19	1 (6%)
12	MAN	N	9	12	11,11,12	0.87	1 (9%)	15,15,17	1.37	2 (13%)
10	NAG	O	1	10,4	14,14,15	0.32	0	17,19,21	0.48	0
10	NAG	O	2	10	14,14,15	0.20	0	17,19,21	0.48	0
10	NAG	P	1	10,4	14,14,15	0.30	0	17,19,21	0.50	0
10	NAG	P	2	10	14,14,15	0.29	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	1	7,4	-	6/6/23/26	0/1/1/1
7	NAG	A	2	7	-	2/6/23/26	0/1/1/1
7	BMA	A	3	7	-	0/2/19/22	0/1/1/1
7	MAN	A	4	7	-	0/2/19/22	0/1/1/1
7	MAN	A	5	7	-	1/2/19/22	0/1/1/1
8	NAG	C	1	8,4	-	2/6/23/26	0/1/1/1
8	NAG	C	2	8	-	6/6/23/26	0/1/1/1
8	BMA	C	3	8	-	2/2/19/22	0/1/1/1
8	MAN	C	4	8	-	0/2/19/22	0/1/1/1
8	MAN	C	5	8	-	2/2/19/22	1/1/1/1
8	MAN	C	6	8	-	0/2/19/22	0/1/1/1
9	NAG	F	1	9,4	-	2/6/23/26	0/1/1/1
9	NAG	F	2	9	-	2/6/23/26	0/1/1/1
9	BMA	F	3	9	-	2/2/19/22	0/1/1/1
9	MAN	F	4	9	-	1/2/19/22	0/1/1/1
10	NAG	I	1	10,4	-	2/6/23/26	0/1/1/1
10	NAG	I	2	10	-	0/6/23/26	0/1/1/1
11	NAG	J	1	11,4	-	3/6/23/26	0/1/1/1
11	NAG	J	2	11	-	2/6/23/26	0/1/1/1
10	NAG	K	1	10,4	-	2/6/23/26	0/1/1/1
10	NAG	K	2	10	-	2/6/23/26	0/1/1/1
10	NAG	M	1	10,4	-	0/6/23/26	0/1/1/1
10	NAG	M	2	10	-	2/6/23/26	0/1/1/1
12	NAG	N	1	12,4	-	4/6/23/26	0/1/1/1
12	MAN	N	10	12	-	0/2/19/22	0/1/1/1
12	NAG	N	2	12	-	2/6/23/26	0/1/1/1
12	BMA	N	3	12	-	0/2/19/22	0/1/1/1
12	MAN	N	4	12	-	2/2/19/22	0/1/1/1
12	MAN	N	5	12	-	2/2/19/22	0/1/1/1
12	MAN	N	6	12	-	0/2/19/22	0/1/1/1
12	MAN	N	7	12	-	1/2/19/22	0/1/1/1
12	MAN	N	8	12	-	0/2/19/22	0/1/1/1
12	MAN	N	9	12	-	0/2/19/22	0/1/1/1
10	NAG	O	1	10,4	-	1/6/23/26	0/1/1/1
10	NAG	O	2	10	-	0/6/23/26	0/1/1/1
10	NAG	P	1	10,4	-	0/6/23/26	0/1/1/1
10	NAG	P	2	10	-	0/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1	NAG	O5-C1	3.30	1.49	1.43
12	N	9	MAN	C1-C2	2.63	1.58	1.52
12	N	6	MAN	C1-C2	2.28	1.57	1.52
12	N	4	MAN	C1-C2	2.17	1.57	1.52
12	N	6	MAN	C2-C3	2.10	1.55	1.52

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1	NAG	C1-O5-C5	5.96	120.17	112.19
7	A	1	NAG	C2-N2-C7	4.69	129.18	122.90
8	C	2	NAG	C2-N2-C7	4.56	129.01	122.90
12	N	6	MAN	C1-O5-C5	4.20	117.82	112.19
12	N	9	MAN	C1-O5-C5	4.05	117.61	112.19

There are no chirality outliers.

5 of 53 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	C	3	BMA	O5-C5-C6-O6
8	C	5	MAN	O5-C5-C6-O6
9	F	3	BMA	C4-C5-C6-O6
10	K	1	NAG	O5-C5-C6-O6
7	A	1	NAG	O5-C5-C6-O6


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	C	5	MAN	C1-C2-C3-C4-C5-O5


9 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	J	1	NAG	1	0
7	A	2	NAG	1	0
8	C	2	NAG	2	0
11	J	2	NAG	1	0
12	N	4	MAN	1	0
9	F	1	NAG	1	0
12	N	1	NAG	1	0
8	C	6	MAN	1	0
8	C	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.























5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
13	NAG	G	617	4	14,14,15	0.28	0	17,19,21	0.43	0
13	NAG	G	2411	4	14,14,15	0.24	0	17,19,21	0.38	0
13	NAG	G	1301	4	14,14,15	0.30	0	17,19,21	0.47	0
13	NAG	G	4421	4	14,14,15	0.20	0	17,19,21	0.43	0
13	NAG	G	2891	4	14,14,15	0.47	0	17,19,21	0.60	0
13	NAG	B	6111	1	14,14,15	0.29	0	17,19,21	0.49	0
13	NAG	G	1601	4	14,14,15	0.33	0	17,19,21	0.39	0
13	NAG	B	6371	1	14,14,15	0.26	0	17,19,21	0.48	0
13	NAG	G	2761	4	14,14,15	0.27	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	NAG	G	617	4	-	1/6/23/26	0/1/1/1
13	NAG	G	2411	4	-	1/6/23/26	0/1/1/1
13	NAG	G	1301	4	-	2/6/23/26	0/1/1/1
13	NAG	G	4421	4	-	2/6/23/26	0/1/1/1
13	NAG	G	2891	4	-	2/6/23/26	0/1/1/1
13	NAG	B	6111	1	-	2/6/23/26	0/1/1/1
13	NAG	G	1601	4	-	2/6/23/26	0/1/1/1
13	NAG	B	6371	1	-	0/6/23/26	0/1/1/1
13	NAG	G	2761	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
13	G	1601	NAG	O5-C5-C6-O6
13	B	6111	NAG	O5-C5-C6-O6
13	G	1601	NAG	C4-C5-C6-O6
13	B	6111	NAG	C4-C5-C6-O6
13	G	2761	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	G	2891	NAG	2	0
13	G	1601	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	129/153 (84%)	1.00	21 (16%) 4 4	14, 32, 63, 100	0
2	D	128/134 (95%)	1.90	54 (42%) 0 0	47, 82, 122, 136	0
3	E	100/114 (87%)	1.81	38 (38%) 1 0	64, 79, 107, 116	0
4	G	430/463 (92%)	1.02	73 (16%) 4 4	21, 42, 87, 113	0
5	H	219/244 (89%)	1.60	73 (33%) 1 1	48, 68, 98, 111	0
6	L	211/217 (97%)	0.72	16 (7%) 20 17	36, 50, 69, 95	0
All	All	1217/1325 (91%)	1.23	275 (22%) 2 2	14, 55, 101, 136	0

The worst 5 of 275 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	G	505	VAL	6.9
4	G	164	GLU	6.3
4	G	457	ASP	5.9
2	D	108	LEU	5.9
1	B	568	LEU	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NAG	A	1	14/15	-	-	30,36,55,56	0
7	NAG	A	2	14/15	-	-	50,63,93,93	0
7	BMA	A	3	11/12	-	-	103,115,125,127	0
7	MAN	A	4	11/12	-	-	95,102,107,111	0
7	MAN	A	5	11/12	-	-	69,88,102,108	0
8	NAG	C	1	14/15	-	-	30,35,63,75	0
8	NAG	C	2	14/15	-	-	32,40,63,70	0
8	BMA	C	3	11/12	-	-	32,34,48,69	0
8	MAN	C	4	11/12	-	-	86,98,107,109	0
8	MAN	C	5	11/12	-	-	107,116,122,127	0
8	MAN	C	6	11/12	-	-	30,39,59,62	0
9	NAG	F	1	14/15	-	-	39,51,57,65	0
9	NAG	F	2	14/15	-	-	57,73,93,97	0
9	BMA	F	3	11/12	-	-	106,115,129,131	0
9	MAN	F	4	11/12	-	-	116,125,135,136	0
10	NAG	O	2	14/15	0.55	0.22	108,121,130,132	0
11	NAG	J	2	14/15	0.64	0.16	102,107,115,115	0
10	NAG	M	2	14/15	0.66	0.21	93,112,125,127	0
12	MAN	N	10	11/12	0.73	0.17	71,84,99,108	0
12	MAN	N	9	11/12	0.76	0.21	86,98,106,108	0
10	NAG	P	2	14/15	0.76	0.19	53,101,112,112	0
11	NAG	J	1	14/15	0.84	0.17	38,89,105,106	0
10	NAG	K	2	14/15	0.85	0.21	117,136,159,162	0
10	NAG	K	1	14/15	0.87	0.17	38,50,69,91	0
12	MAN	N	7	11/12	0.88	0.15	78,87,97,105	0
10	NAG	M	1	14/15	0.89	0.13	32,72,86,92	0
12	MAN	N	8	11/12	0.90	0.12	58,68,78,90	0
12	NAG	N	1	14/15	0.90	0.12	37,51,67,68	0
10	NAG	O	1	14/15	0.90	0.14	63,71,92,104	0
10	NAG	I	2	14/15	0.91	0.12	82,102,110,111	0
10	NAG	P	1	14/15	0.91	0.10	54,63,88,101	0
12	NAG	N	2	14/15	0.92	0.11	41,47,61,71	0
12	MAN	N	6	11/12	0.92	0.14	41,60,67,76	0
10	NAG	I	1	14/15	0.92	0.11	44,69,92,102	0
12	BMA	N	3	11/12	0.94	0.09	40,41,60,73	0
12	MAN	N	4	11/12	0.95	0.09	38,43,49,65	0
12	MAN	N	5	11/12	0.96	0.09	39,40,51,62	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain A:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain I:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain M:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain O:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain P:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
13	NAG	G	1601	14/15	0.62	0.21	72,91,97,97	0
13	NAG	G	2761	14/15	0.67	0.20	86,101,118,118	0
13	NAG	B	6111	14/15	0.73	0.14	48,72,85,96	0
13	NAG	G	4421	14/15	0.75	0.27	105,117,125,130	0
13	NAG	G	1301	14/15	0.76	0.13	86,89,96,98	0
13	NAG	B	6371	14/15	0.81	0.13	79,91,98,106	0
13	NAG	G	2411	14/15	0.82	0.17	56,74,97,103	0
13	NAG	G	617	14/15	0.89	0.12	56,68,72,80	0
13	NAG	G	2891	14/15	0.90	0.11	41,45,75,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.