



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:11 PM UTC

PDB ID : 9CKJ / pdb_00009ckj
Title : Co-crystal structure of 53BP1 tandem Tudor domains in complex with UNC9512
Authors : Zeng, H.; Dong, A.; Shell, D.J.; Foley, C.; Arrowsmith, C.H.; Edwards, A.M.; Halabelian, L.; Structural Genomics Consortium (SGC)
Deposited on : 2024-07-09
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	124	
1	B	124	
1	C	124	
1	D	124	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	124	11% 86% 9% 5%
1	F	124	11% 90% 6% • •

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TP53-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	118	Total	C	N	O	S	0	0	0
			935	601	156	175	3			
1	B	120	Total	C	N	O	S	0	0	0
			929	599	152	175	3			
1	C	119	Total	C	N	O	S	0	1	0
			952	610	158	181	3			
1	D	119	Total	C	N	O	S	0	0	0
			903	581	148	171	3			
1	E	118	Total	C	N	O	S	0	0	0
			884	576	140	165	3			
1	F	121	Total	C	N	O	S	0	0	0
			936	605	150	178	3			

- Molecule 2 is (3S)-1-[4-(4-methylpiperazin-1-yl)pyridine-2-carbonyl]-N-[(3M)-3-(3-oxo-2,3-dihydro-1H-isoindol-5-yl)phenyl]piperidine-3-carboxamide (CCD ID: A1AY8) (formula: C₃₁H₃₄N₆O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			40	31	6	3		
2	B	1	Total	C	N	O	0	0
			40	31	6	3		
2	C	1	Total	C	N	O	0	0
			40	31	6	3		
2	D	1	Total	C	N	O	0	0
			40	31	6	3		
2	E	1	Total	C	N	O	0	0
			40	31	6	3		
2	F	1	Total	C	N	O	0	0
			40	31	6	3		

- Molecule 3 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	2	Total	X	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	56	Total	O	0	0
			56	56		
4	B	40	Total	O	0	1
			41	41		
4	C	48	Total	O	0	1
			49	49		
4	D	19	Total	O	0	0
			19	19		
4	E	12	Total	O	0	0
			12	12		
4	F	21	Total	O	0	0
			21	21		

- Molecule 1: TP53-binding protein 1



- GLY
 M1484
 V1492
 S1496
 S1497
 M1498
 G1499
 Y1500
 F1501
 K1514
 L1517
 D1550
 E1551
 Y1552
 E1564
 I1572
 Y1581
 K1582
 R1583
 L1588
 S1589
 G1603
 PRO
 TYR
 GLU


- | | | | | | | | | | | | | | | | | | | | |
|-----|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-----|-----|-----|
| GLY | ASN | S1485 | V1492 | N1498 | G1499 | Y1500 | F1501 | L1517 | L1533 | I1572 | Y1581 | K1582 | R1583 | L1588 | S1589 | G1603 | PRO | TYP | GLU |
|-----|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-----|-----|-----|

- [illegible]

- | GLY | ASN | PRO | TYR | GLU |
|-------|-----|-----|-----|-----|
| S1485 | | | | |
| F1486 | | | | |
| N1490 | | | | |
| F1491 | | | | |
| F1492 | | | | |
| A1493 | | | | |
| K1494 | | | | |
| N1495 | | | | |
| S1496 | | | | |
| S1497 | | | | |
| N1498 | | | | |
| F1501 | | | | |
| L1506 | | | | |
| G1511 | | | | |
| L1517 | | | | |
| F1527 | | | | |
| E1549 | | | | |
| D1550 | | | | |
| E1551 | | | | |
| Y1552 | | | | |
| L1572 | | | | |
| Y1580 | | | | |
| K1582 | | | | |
| R1583 | | | | |
| L1588 | | | | |
| L1602 | | | | |
| GLY | | | | |
| PRO | | | | |
| TYR | | | | |
| GLU | | | | |

● Molecule 1: TP53-binding protein 1

Chain F:  11% 90% 6% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.88Å 76.03Å 258.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.92 – 2.25 29.92 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.5 (29.92-2.25) 97.4 (29.92-2.25)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.219 , 0.261 0.232 , 0.282	Depositor DCC
R_{free} test set	1892 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	40.4	Xtriage
Anisotropy	0.341	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5979	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, A1AY8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	0/954	1.16	1/1284 (0.1%)
1	B	0.94	0/948	1.17	1/1279 (0.1%)
1	C	0.92	0/971	1.16	1/1306 (0.1%)
1	D	0.85	0/922	1.17	3/1248 (0.2%)
1	E	0.80	0/903	1.17	5/1223 (0.4%)
1	F	0.88	0/957	1.15	2/1295 (0.2%)
All	All	0.89	0/5655	1.16	13/7635 (0.2%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1598	GLU	CB-CG-CD	7.32	125.05	112.60
1	A	1498	ASN	CA-CB-CG	6.13	118.73	112.60
1	B	1498	ASN	CA-CB-CG	5.78	118.38	112.60
1	D	1498	ASN	CA-CB-CG	5.70	118.30	112.60
1	E	1511	GLY	CA-C-N	5.30	129.87	122.08
1	E	1511	GLY	C-N-CA	5.30	129.87	122.08
1	F	1571	SER	N-CA-C	-5.21	100.67	108.96
1	C	1498	ASN	CA-CB-CG	5.16	117.76	112.60
1	F	1498	ASN	CA-CB-CG	5.14	117.74	112.60
1	E	1493	ALA	CA-C-N	5.08	127.92	120.71
1	E	1493	ALA	C-N-CA	5.08	127.92	120.71
1	D	1572	ILE	CB-CG1-CD1	5.04	124.39	113.80
1	E	1495	TRP	N-CA-C	-5.01	99.12	107.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	935	0	908	8	0
1	B	929	0	882	4	0
1	C	952	0	919	5	0
1	D	903	0	826	6	0
1	E	884	0	800	4	0
1	F	936	0	871	5	0
2	A	40	0	0	0	0
2	B	40	0	0	0	0
2	C	40	0	0	0	0
2	D	40	0	0	0	0
2	E	40	0	0	0	0
2	F	40	0	0	0	0
3	F	2	0	0	0	0
4	A	56	0	0	1	0
4	B	41	0	0	0	0
4	C	49	0	0	0	0
4	D	19	0	0	0	0
4	E	12	0	0	0	0
4	F	21	0	0	1	0
All	All	5979	0	5206	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:1562:ARG:HD2	4:F:2120:HOH:O	2.01	0.61
1:A:1544:VAL:HG21	1:A:1572:ILE:HD12	1.82	0.60
1:A:1490:ARG:HH11	1:A:1490:ARG:HB3	1.68	0.57
1:A:1544:VAL:CG2	1:A:1572:ILE:HD12	2.35	0.57
1:F:1492:VAL:HG12	1:F:1501:PHE:HB3	1.91	0.53
1:E:1490:ARG:HH11	1:E:1490:ARG:HB3	1.76	0.51
1:D:1544:VAL:CG2	1:D:1572:ILE:HD12	2.40	0.51
1:D:1544:VAL:HG21	1:D:1572:ILE:HD12	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1501:PHE:HB2	1:A:1588:LEU:HB2	1.92	0.50
1:D:1492:VAL:HG12	1:D:1501:PHE:HB3	1.94	0.50
1:B:1572:ILE:HD12	1:B:1581:TYR:CD1	2.49	0.48
1:C:1501:PHE:HB2	1:C:1588:LEU:HB2	1.95	0.48
1:F:1562:ARG:HE	1:F:1564:GLU:HG3	1.79	0.48
1:E:1492:VAL:HG12	1:E:1501:PHE:HB3	1.96	0.47
1:A:1508:ARG:HH21	1:A:1516:LYS:HE3	1.78	0.47
1:D:1501:PHE:HB2	1:D:1588:LEU:HB2	1.97	0.47
1:A:1492:VAL:HG12	1:A:1501:PHE:HB3	1.96	0.47
1:B:1492:VAL:HG12	1:B:1501:PHE:HB3	1.96	0.47
1:B:1501:PHE:HB2	1:B:1588:LEU:HB2	1.98	0.45
1:C:1572:ILE:HD12	1:C:1581:TYR:CD1	2.52	0.45
1:E:1501:PHE:HB2	1:E:1588:LEU:HB2	1.99	0.45
1:E:1572:ILE:HD12	1:E:1581:TYR:CD1	2.52	0.44
1:A:1490:ARG:HG2	1:A:1505:LYS:HG2	2.00	0.43
1:A:1562:ARG:HB2	4:A:2155:HOH:O	2.17	0.43
1:F:1501:PHE:HB2	1:F:1588:LEU:HB2	2.01	0.43
1:B:1500:TYR:CD1	1:B:1589:SER:HA	2.54	0.43
1:F:1572:ILE:HD12	1:F:1581:TYR:CD1	2.55	0.41
1:C:1583[B]:ARG:HE	1:C:1583[B]:ARG:HB3	1.51	0.41
1:C:1492:VAL:HG12	1:C:1501:PHE:HB3	2.01	0.41
1:D:1493:ALA:HB3	1:D:1517:LEU:HD22	2.02	0.40
1:C:1500:TYR:CD2	1:C:1589:SER:HA	2.56	0.40
1:D:1486:PHE:HA	1:D:1489:LEU:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/124 (94%)	114 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	118/124 (95%)	113 (96%)	5 (4%)	0	100	100
1	C	118/124 (95%)	115 (98%)	3 (2%)	0	100	100
1	D	117/124 (94%)	113 (97%)	4 (3%)	0	100	100
1	E	116/124 (94%)	112 (97%)	4 (3%)	0	100	100
1	F	119/124 (96%)	117 (98%)	2 (2%)	0	100	100
All	All	704/744 (95%)	684 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	94/104 (90%)	92 (98%)	2 (2%)	47	58
1	B	91/104 (88%)	86 (94%)	5 (6%)	19	20
1	C	97/104 (93%)	94 (97%)	3 (3%)	35	44
1	D	83/104 (80%)	79 (95%)	4 (5%)	23	25
1	E	77/104 (74%)	75 (97%)	2 (3%)	40	50
1	F	89/104 (86%)	86 (97%)	3 (3%)	32	40
All	All	531/624 (85%)	512 (96%)	19 (4%)	31	39

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1498	ASN
1	A	1517	LEU
1	B	1498	ASN
1	B	1514	LYS
1	B	1517	LEU
1	B	1564	GLU
1	B	1583	ARG
1	C	1498	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	1517	LEU
1	C	1533	LEU
1	D	1498	ASN
1	D	1510	VAL
1	D	1517	LEU
1	D	1563	LYS
1	E	1498	ASN
1	E	1517	LEU
1	F	1498	ASN
1	F	1517	LEU
1	F	1564	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are unknown - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1AY8	D	2001	-	45,45,45	0.43	0	63,64,64	0.50	0
2	A1AY8	A	2001	-	45,45,45	0.49	0	63,64,64	0.55	0
2	A1AY8	E	2001	-	45,45,45	0.45	0	63,64,64	0.47	0
2	A1AY8	B	2001	-	45,45,45	0.49	0	63,64,64	0.56	1 (1%)
2	A1AY8	F	2001	-	45,45,45	0.44	0	63,64,64	0.54	0
2	A1AY8	C	2001	-	45,45,45	0.49	0	63,64,64	0.53	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1AY8	D	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	A	2001	-	-	0/24/53/53	0/6/6/6
2	A1AY8	E	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	B	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	F	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	C	2001	-	-	0/24/53/53	0/6/6/6

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	A1AY8	C6-N7-C8	2.15	123.96	118.11
2	C	2001	A1AY8	C6-N7-C8	2.02	123.60	118.11

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2001	A1AY8	C37-C22-N21-C20
2	E	2001	A1AY8	C37-C22-N21-C20
2	F	2001	A1AY8	C37-C22-N21-C20
2	D	2001	A1AY8	C37-C22-N21-C20


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In


addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/124 (95%)	0.16	3 (2%) 58 59	24, 38, 59, 75	0
1	B	120/124 (96%)	0.09	5 (4%) 40 40	25, 38, 65, 75	0
1	C	119/124 (95%)	-0.06	1 (0%) 82 84	19, 36, 50, 58	1 (0%)
1	D	119/124 (95%)	0.66	7 (5%) 28 26	36, 56, 80, 104	0
1	E	118/124 (95%)	0.93	14 (11%) 9 8	45, 62, 84, 101	0
1	F	121/124 (97%)	0.74	14 (11%) 9 9	34, 54, 79, 86	0
All	All	715/744 (96%)	0.42	44 (6%) 26 24	19, 48, 76, 104	1 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	1602	LEU	5.3
1	D	1603	GLY	4.4
1	A	1496	SER	4.2
1	D	1550	ASP	4.1
1	C	1603	GLY	3.8
1	E	1495	TRP	3.5
1	E	1485	SER	3.4
1	F	1603	GLY	3.3
1	F	1512	ALA	3.3
1	A	1603	GLY	3.2
1	E	1496	SER	3.0
1	F	1527	VAL	3.0
1	D	1485	SER	2.9
1	B	1603	GLY	2.8
1	B	1550	ASP	2.7
1	F	1529	GLY	2.7
1	E	1486	PHE	2.6
1	D	1486	PHE	2.6
1	D	1510	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	1527	VAL	2.6
1	E	1506	ILE	2.5
1	F	1486	PHE	2.5
1	B	1551	GLU	2.4
1	B	1496	SER	2.3
1	F	1497	SER	2.3
1	E	1583	ARG	2.3
1	E	1580	TRP	2.3
1	E	1497	SER	2.3
1	E	1550	ASP	2.2
1	E	1552	TYR	2.2
1	F	1487	VAL	2.2
1	B	1552	TYR	2.2
1	A	1550	ASP	2.2
1	F	1604	PRO	2.2
1	E	1549	GLU	2.2
1	F	1551	GLU	2.2
1	F	1605	TYR	2.2
1	F	1495	TRP	2.2
1	F	1498	ASN	2.2
1	D	1508	ARG	2.1
1	D	1551	GLU	2.1
1	E	1517	LEU	2.1
1	F	1552	TYR	2.0
1	F	1519	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UNX	F	2002	1/1	0.55	0.16	30,30,30,30	0
3	UNX	F	2003	1/1	0.80	0.12	30,30,30,30	0
2	A1AY8	F	2001	40/40	0.90	0.10	48,54,60,61	0
2	A1AY8	E	2001	40/40	0.91	0.11	46,66,74,74	0
2	A1AY8	C	2001	40/40	0.93	0.08	25,34,45,46	0
2	A1AY8	D	2001	40/40	0.93	0.09	28,36,47,48	0
2	A1AY8	A	2001	40/40	0.94	0.07	22,30,37,37	0
2	A1AY8	B	2001	40/40	0.95	0.07	23,28,36,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1AY8 F 2001:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1AY8 E 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1AY8 C 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1AY8 D 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1AY8 A 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1AY8 B 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.