



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 01:53 AM UTC

PDB ID : 4CLP / pdb_00004clp
Title : Crystal structure of human soluble Adenylyl Cyclase complex with adenosine
-3',5'-cyclic-monophosphate
Authors : Kleinboelting, S.; Weyand, M.; Steegborn, C.
Deposited on : 2014-01-15
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ADENYLATE CYCLASE TYPE 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	460	3678	2385	593	665	35	0	10	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	470	HIS	-	expression tag	UNP Q96PN6
A	471	HIS	-	expression tag	UNP Q96PN6
A	472	HIS	-	expression tag	UNP Q96PN6
A	473	HIS	-	expression tag	UNP Q96PN6
A	474	HIS	-	expression tag	UNP Q96PN6
A	475	HIS	-	expression tag	UNP Q96PN6

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ADENOSINE-3',5'-CYCLIC-MONOPHOSPHATE (CCD ID: CMP) (formula: $C_{10}H_{12}N_5O_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		
4	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	209	Total	O	0	0
			209	209		

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	100.59Å 100.59Å 97.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	87.11 – 1.90 87.11 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (87.11-1.90) 100.0 (87.11-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.161 , 0.207 0.172 , 0.212	Depositor DCC
R_{free} test set	2208 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3978	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	A	253	1	8,9,10	0.74	0	6,9,11	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	253	1	-	2/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	253	CME	SD-CE-CZ-OH
1	A	253	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	1472	-	5,5,5	1.33	0	5,5,5	1.83	2 (40%)
3	GOL	A	1471	-	5,5,5	0.74	0	5,5,5	1.55	1 (20%)
4	CMP	A	1474	-	25,25,25	2.09	6 (24%)	37,39,39	2.65	14 (37%)
2	ACT	A	1469	-	3,3,3	1.18	0	3,3,3	0.57	0
5	EDO	A	1478	-	3,3,3	0.81	0	2,2,2	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	A	1468	-	3,3,3	0.34	0	3,3,3	1.72	1 (33%)
5	EDO	A	1477	-	3,3,3	0.34	0	2,2,2	0.73	0
6	PEG	A	1479	-	6,6,6	0.53	0	5,5,5	0.67	0
5	EDO	A	1475	-	3,3,3	0.43	0	2,2,2	0.09	0
5	EDO	A	1476	-	3,3,3	0.29	0	2,2,2	0.57	0
4	CMP	A	1473	-	25,25,25	1.94	5 (20%)	37,39,39	3.50	21 (56%)
2	ACT	A	1470	-	3,3,3	1.76	1 (33%)	3,3,3	1.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1472	-	-	2/4/4/4	-
3	GOL	A	1471	-	-	4/4/4/4	-
4	CMP	A	1474	-	-	0/4/31/31	0/4/4/4
5	EDO	A	1478	-	-	1/1/1/1	-
5	EDO	A	1477	-	-	1/1/1/1	-
6	PEG	A	1479	-	-	0/4/4/4	-
5	EDO	A	1475	-	-	1/1/1/1	-
5	EDO	A	1476	-	-	1/1/1/1	-
4	CMP	A	1473	-	-	1/4/31/31	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1473	CMP	C5-C4	5.30	1.48	1.39
4	A	1474	CMP	C8-N7	4.67	1.40	1.31
4	A	1474	CMP	P-O5'	4.50	1.62	1.57
4	A	1474	CMP	C4-N9	-3.83	1.29	1.37
4	A	1474	CMP	C5-C4	3.83	1.45	1.39
4	A	1473	CMP	P-O5'	3.71	1.61	1.57
4	A	1474	CMP	P-O3'	3.67	1.63	1.57
4	A	1473	CMP	C5-N7	-3.66	1.32	1.39
4	A	1473	CMP	P-O3'	3.26	1.63	1.57
4	A	1473	CMP	O5'-C5'	-3.06	1.41	1.46
4	A	1474	CMP	C5-C6	2.53	1.48	1.41
2	A	1470	ACT	CH3-C	2.26	1.58	1.49

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1473	CMP	C4'-O4'-C1'	-9.51	88.47	109.47
4	A	1473	CMP	C5-C4-N3	-7.10	116.95	126.72
4	A	1473	CMP	N3-C4-N9	7.03	139.12	127.17
4	A	1473	CMP	C1'-N9-C8	-6.64	112.36	127.09
4	A	1474	CMP	O5'-P-O3'	-6.20	97.40	105.70
4	A	1473	CMP	O4'-C1'-N9	6.04	119.69	108.09
4	A	1473	CMP	C4-N9-C1'	5.33	139.09	126.63
4	A	1473	CMP	O5'-C5'-C4'	-5.10	93.81	105.71
4	A	1474	CMP	C4-N9-C1'	-5.05	114.83	126.63
4	A	1474	CMP	C1'-N9-C8	4.89	137.95	127.09
4	A	1474	CMP	C4'-O4'-C1'	-4.55	99.42	109.47
4	A	1474	CMP	C6-C5-N7	4.35	140.47	132.09
4	A	1474	CMP	O4'-C1'-N9	4.31	116.36	108.09
4	A	1474	CMP	C5-C4-N9	4.11	110.29	105.81
4	A	1474	CMP	C4-C5-N7	-4.05	105.95	110.58
4	A	1474	CMP	O3'-C3'-C2'	3.82	119.35	115.61
4	A	1473	CMP	C2-N3-C4	3.80	121.11	111.83
4	A	1473	CMP	C2'-C1'-N9	3.64	122.34	113.30
4	A	1474	CMP	O2P-P-O1P	3.41	119.05	108.56
4	A	1473	CMP	O2P-P-O1P	3.18	118.33	108.56
4	A	1474	CMP	C2-N3-C4	3.11	119.42	111.83
4	A	1474	CMP	C5-C4-N3	-3.09	122.47	126.72
4	A	1473	CMP	N3-C2-N1	-3.02	124.00	128.58
3	A	1471	GOL	O3-C3-C2	-2.93	97.18	110.38
4	A	1473	CMP	N6-C6-N1	2.83	124.68	118.38
3	A	1472	GOL	O2-C2-C1	2.69	120.32	109.18
4	A	1474	CMP	C6-C5-C4	-2.66	113.55	117.18
4	A	1473	CMP	O3'-C3'-C4'	2.64	112.71	110.71
4	A	1473	CMP	C4-N9-C8	2.57	108.44	105.74
3	A	1472	GOL	C3-C2-C1	-2.50	102.64	111.80
4	A	1473	CMP	C5-N7-C8	2.46	107.32	103.45
4	A	1473	CMP	O2P-P-O3'	2.43	112.71	107.04
4	A	1473	CMP	O5'-P-O3'	2.31	108.80	105.70
4	A	1473	CMP	O5'-P-O1P	-2.30	105.14	110.44
4	A	1473	CMP	C5-C6-N6	-2.23	117.76	123.29
2	A	1468	ACT	OXT-C-CH3	2.22	124.35	115.05
4	A	1474	CMP	N3-C2-N1	-2.19	125.26	128.58
4	A	1473	CMP	C2'-C3'-C4'	-2.07	99.61	103.24
4	A	1473	CMP	C4-C5-N7	-2.04	108.25	110.58

There are no chirality outliers.


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1471	GOL	O1-C1-C2-C3
3	A	1472	GOL	O1-C1-C2-C3
3	A	1471	GOL	C1-C2-C3-O3
3	A	1471	GOL	O1-C1-C2-O2
3	A	1472	GOL	O1-C1-C2-O2
5	A	1475	EDO	O1-C1-C2-O2
5	A	1476	EDO	O1-C1-C2-O2
5	A	1477	EDO	O1-C1-C2-O2
5	A	1478	EDO	O1-C1-C2-O2
4	A	1473	CMP	O4'-C1'-N9-C8
3	A	1471	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data ⓘ

5.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/475 (96%)	0.12	28 (6%) 27 29	14, 30, 67, 107	10 (2%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	351	PRO	6.3
1	A	350	PHE	6.1
1	A	356	PRO	5.7
1	A	8	PHE	4.9
1	A	355	VAL	4.4
1	A	51	PHE	4.2
1	A	139	LEU	4.0
1	A	338	PHE	3.7
1	A	352	GLY	3.5
1	A	130	PHE	3.2
1	A	285	TYR	2.8
1	A	283	GLN	2.8
1	A	467	GLU	2.7
1	A	184	MET	2.7
1	A	449	VAL	2.7
1	A	9	GLN	2.6
1	A	353	GLU	2.6
1	A	454	ALA	2.5
1	A	54	MET	2.5
1	A	466	THR	2.5
1	A	5	LYS	2.5
1	A	141	ILE	2.4
1	A	339	ASP	2.3
1	A	223	ASN	2.3
1	A	453	VAL	2.1
1	A	28	HIS	2.1
1	A	48	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	53	ALA	2.1

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	253	10/11	0.99	0.06	19,24,31,31	0

5.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	1476	4/4	0.75	0.15	66,67,73,74	0
5	EDO	A	1478	4/4	0.75	0.22	45,54,57,65	0
4	CMP	A	1474	22/22	0.76	0.18	32,66,95,101	22
5	EDO	A	1477	4/4	0.80	0.18	68,70,71,72	0
2	ACT	A	1470	4/4	0.84	0.17	37,37,44,46	0
6	PEG	A	1479	7/7	0.85	0.14	45,51,55,69	0
2	ACT	A	1469	4/4	0.87	0.10	27,35,42,44	0
5	EDO	A	1475	4/4	0.87	0.15	42,48,62,66	0
4	CMP	A	1473	22/22	0.89	0.11	43,61,67,75	0
3	GOL	A	1472	6/6	0.90	0.14	41,46,52,55	0
3	GOL	A	1471	6/6	0.90	0.11	33,41,45,45	0
2	ACT	A	1468	4/4	0.94	0.13	30,30,31,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CMP A 1474:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CMP A 1473:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



5.5 Other polymers [i](#)

There are no such residues in this entry.