



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:51 AM UTC

PDB ID : 1CML / pdb_00001cml
Title : CHALCONE SYNTHASE FROM ALFALFA COMPLEXED WITH MALONYL-COA
Authors : Ferrer, J.-L.; Jez, J.; Bowman, M.E.; Dixon, R.; Noel, J.P.
Deposited on : 1999-03-30
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3630 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (CHALCONE SYNTHASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	389	3154	2001	531	603	19	0	19	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	164	ALA	CYS	engineered mutation	UNP P30074

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is MALONYL-COENZYME A (CCD ID: MLC) (formula: C₂₄H₃₈N₇O₁₉P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	
			54	24	7	19	3	1	
								3	0

- Molecule 4 is PIPERAZINE-N,N'-BIS(2-ETHANESULFONIC ACID) (CCD ID: PIN) (formula: $C_8H_{18}N_2O_6S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S		
			18	8	2	6	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	399	Total 399	O 399	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (CHALCONE SYNTHASE)



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	97.77Å 97.77Å 65.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	84.52 – 1.69 84.52 – 1.69	Depositor EDS
% Data completeness (in resolution range)	97.4 (84.52-1.69) 97.3 (84.52-1.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.42 (at 1.69Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.167 , 0.200 0.275 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	15.5	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3630	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MLC, SO4, PIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/3213	1.33	21/4346 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	ARG	CD-NE-CZ	14.64	144.90	124.40
1	A	376	GLY	O-C-N	-10.78	108.69	122.70
1	A	342	VAL	N-CA-CB	-10.35	92.95	110.65
1	A	232	ILE	N-CA-C	-8.66	103.45	111.67
1	A	342	VAL	CB-CA-C	8.47	125.29	112.16
1	A	137	MET	CA-C-O	-7.07	113.07	119.86
1	A	341	CYS	CA-C-O	6.68	128.04	120.90
1	A	67	ARG	NE-CZ-NH1	6.37	127.87	121.50
1	A	356	ASN	N-CA-C	6.36	120.52	113.21
1	A	291	GLY	CA-C-N	-6.32	115.36	123.19
1	A	291	GLY	C-N-CA	-6.32	115.36	123.19
1	A	138	PRO	N-CA-CB	6.19	109.41	102.60
1	A	288	GLU	CB-CG-CD	5.83	122.51	112.60
1	A	167	GLY	N-CA-C	-5.65	105.80	113.37
1	A	137	MET	CG-SD-CE	-5.59	88.61	100.90
1	A	137	MET	O-C-N	5.57	126.72	121.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	ARG	NE-CZ-NH2	-5.55	114.21	119.20
1	A	284	VAL	N-CA-CB	5.46	118.77	110.58
1	A	26	ASN	CA-CB-CG	5.19	117.79	112.60
1	A	341	CYS	CB-CA-C	-5.17	102.55	110.81
1	A	303	HIS	CA-CB-CG	-5.11	108.69	113.80

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	265[B]	PHE	Mainchain
1	A	276[B]	SER	Mainchain
1	A	32[B]	GLU	Mainchain
1	A	354[B]	THR	Mainchain
1	A	82[B]	ASN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3154	0	3169	45	0
2	A	5	0	0	0	0
3	A	54	0	25	5	0
4	A	18	0	12	0	0
5	A	399	0	0	26	8
All	All	3630	0	3206	45	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:VAL:HB	5:A:763:HOH:O	1.65	0.94
1:A:59:MET:HE2	3:A:390:MLC:H1'	1.57	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:VAL:HG21	5:A:763:HOH:O	1.90	0.72
1:A:14[A]:GLU:HG3	1:A:227:ASP:OD1	1.91	0.70
1:A:38:ASP:O	1:A:42[B]:LYS:HD3	1.97	0.65
1:A:292:ILE:HG21	5:A:766:HOH:O	1.97	0.64
1:A:70:MET:HG2	5:A:753:HOH:O	1.98	0.63
1:A:59:MET:CE	3:A:390:MLC:H1'	2.28	0.63
1:A:301:ILE:HG21	5:A:755:HOH:O	2.02	0.60
1:A:58:ARG:HD2	3:A:390:MLC:P3	2.41	0.59
1:A:342:VAL:HG22	5:A:755:HOH:O	2.04	0.57
1:A:58:ARG:HD2	3:A:390:MLC:O33	2.07	0.56
1:A:54:GLU:HB3	1:A:58:ARG:NH2	2.24	0.53
1:A:265[B]:PHE:HB3	5:A:522:HOH:O	2.10	0.52
1:A:370:LEU:CD1	5:A:755:HOH:O	2.59	0.50
1:A:341:CYS:HB2	5:A:400:HOH:O	2.11	0.50
1:A:9[A]:LYS:HG3	5:A:614:HOH:O	2.11	0.50
1:A:301:ILE:CG2	5:A:755:HOH:O	2.56	0.49
1:A:48:HIS:HA	5:A:772:HOH:O	2.11	0.49
1:A:264[B]:THR:HG23	5:A:538:HOH:O	2.13	0.49
1:A:98:VAL:HG11	5:A:763:HOH:O	2.12	0.48
1:A:194:THR:HA	5:A:770:HOH:O	2.13	0.47
1:A:9[A]:LYS:HG3	5:A:618:HOH:O	2.15	0.47
1:A:342:VAL:HB	5:A:400:HOH:O	2.14	0.47
1:A:354[B]:THR:O	1:A:355:GLN:C	2.57	0.47
1:A:66:LYS:HB3	1:A:67:ARG:HH11	1.80	0.47
1:A:215[B]:PHE:HD2	5:A:548:HOH:O	1.98	0.46
1:A:277:LYS:HE2	5:A:703:HOH:O	2.15	0.46
1:A:352:LYS:HA	1:A:352:LYS:HD2	1.75	0.45
1:A:75:GLU:O	1:A:78:LYS:HD2	2.17	0.45
1:A:133:SER:HB3	5:A:589:HOH:O	2.15	0.45
1:A:229:VAL:HG12	1:A:232:ILE:HD13	1.99	0.45
1:A:370:LEU:HG	5:A:755:HOH:O	2.17	0.44
1:A:42[B]:LYS:HD2	1:A:47:GLU:OE2	2.18	0.44
1:A:55:LYS:HE2	5:A:727:HOH:O	2.18	0.44
1:A:370:LEU:HD11	5:A:755:HOH:O	2.18	0.44
1:A:78:LYS:HD3	1:A:79:GLU:HG3	2.01	0.43
1:A:188:VAL:O	1:A:221:ALA:HA	2.19	0.42
1:A:277:LYS:HB3	1:A:277:LYS:HE3	1.72	0.42
1:A:277:LYS:HG3	5:A:700:HOH:O	2.20	0.42
1:A:42[B]:LYS:HD2	5:A:765:HOH:O	2.20	0.41
1:A:271:VAL:HB	1:A:272:PRO:HD3	2.03	0.41
1:A:281:LYS:HB3	5:A:473:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:LYS:HB2	3:A:390:MLC:HPB1	2.02	0.41
1:A:29:ASN:HB3	1:A:70:MET:O	2.21	0.40

All (8) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:536:HOH:O	5:A:760:HOH:O[4_557]	1.20	1.00
5:A:620:HOH:O	5:A:620:HOH:O[4_557]	1.70	0.50
5:A:669:HOH:O	5:A:767:HOH:O[4_556]	1.95	0.25
5:A:546:HOH:O	5:A:616:HOH:O[6_656]	2.03	0.17
5:A:618:HOH:O	5:A:619:HOH:O[4_557]	2.05	0.15
5:A:596:HOH:O	5:A:780:HOH:O[6_656]	2.09	0.11
5:A:563:HOH:O	5:A:761:HOH:O[3_565]	2.10	0.10
5:A:552:HOH:O	5:A:566:HOH:O[5_666]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	406/389 (104%)	393 (97%)	12 (3%)	1 (0%)	43 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	343/324 (106%)	333 (97%)	10 (3%)	37 20

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42[A]	LYS
1	A	42[B]	LYS
1	A	67	ARG
1	A	78	LYS
1	A	133	SER
1	A	151	ARG
1	A	281	LYS
1	A	284	VAL
1	A	292	ILE
1	A	342	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	181	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	392	-	4,4,4	0.80	0	6,6,6	0.24	0
4	PIN	A	391	-	18,18,18	2.57	7 (38%)	24,26,26	2.27	7 (29%)
3	MLC	A	390	-	53,56,56	2.85	12 (22%)	77,83,83	2.30	19 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PIN	A	391	-	-	7/12/22/22	0/1/1/1
3	MLC	A	390	-	-	20/54/71/71	0/3/3/3

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	390	MLC	CM2-CM3	-17.04	1.24	1.51
4	A	391	PIN	C1-S1	-6.11	1.69	1.77
4	A	391	PIN	C1'-S1'	-5.01	1.70	1.77
4	A	391	PIN	O3-S1	4.86	1.65	1.47
3	A	390	MLC	P2-O6	-4.08	1.55	1.59
3	A	390	MLC	CP2-NP1	3.97	1.55	1.46
3	A	390	MLC	OM4-CM3	3.75	1.34	1.22
4	A	391	PIN	O3'-S1'	3.51	1.60	1.47
3	A	390	MLC	P2-O7	-3.33	1.46	1.59
3	A	390	MLC	OM2-CM1	3.17	1.26	1.21
3	A	390	MLC	CP5-NP2	3.14	1.53	1.46
3	A	390	MLC	OM3-CM3	-2.91	1.21	1.30
3	A	390	MLC	P3-O33	2.77	1.59	1.50
3	A	390	MLC	CPB-CPA	2.64	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	390	MLC	CM1-S	2.53	1.82	1.76
4	A	391	PIN	O1-S1	-2.37	1.38	1.45
4	A	391	PIN	O1'-S1'	-2.28	1.38	1.45
3	A	390	MLC	C2-N1	2.07	1.37	1.33
4	A	391	PIN	O2-S1	-2.06	1.39	1.45

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	390	MLC	CP2-NP1-CP3	-7.26	109.31	122.82
3	A	390	MLC	OP1-CP3-NP1	-6.30	110.66	123.03
3	A	390	MLC	CP5-NP2-CP6	-6.07	111.64	122.55
4	A	391	PIN	O1'-S1'-C1'	5.38	114.86	106.73
4	A	391	PIN	O2-S1-C1	5.22	114.62	106.73
3	A	390	MLC	CP9-CPA-CP7	5.12	117.50	108.77
3	A	390	MLC	CP1-S-CM1	4.75	115.89	101.84
3	A	390	MLC	OM4-CM3-CM2	-4.63	108.93	122.11
3	A	390	MLC	OM2-CM1-S	-4.42	117.06	122.68
3	A	390	MLC	CM2-CM1-S	4.25	119.03	113.63
3	A	390	MLC	O4'-C4'-C3'	3.81	112.96	104.92
3	A	390	MLC	C2'-C3'-C4'	-3.70	96.76	103.24
3	A	390	MLC	OM3-CM3-CM2	3.66	125.84	114.51
3	A	390	MLC	C4'-O4'-C1'	-3.50	101.75	109.47
4	A	391	PIN	O3-S1-O2	-3.44	102.78	111.40
4	A	391	PIN	O3'-S1'-C1'	-3.20	99.75	106.00
3	A	390	MLC	O4'-C1'-C2'	3.03	113.10	106.62
4	A	391	PIN	O3-S1-C1	-3.02	100.09	106.00
3	A	390	MLC	CP4-CP3-NP1	2.84	121.52	116.34
3	A	390	MLC	OP3-CP7-CP6	-2.79	97.72	109.38
3	A	390	MLC	O12-P1-O6	2.53	114.11	107.27
4	A	391	PIN	O1-S1-C1	2.44	110.41	106.73
4	A	391	PIN	O3-S1-O1	-2.27	105.73	111.40
3	A	390	MLC	CP5-CP4-CP3	-2.27	108.62	112.39
3	A	390	MLC	P3-O3'-C3'	-2.04	118.00	123.43
3	A	390	MLC	O3'-C3'-C4'	2.01	117.12	110.03

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	390	MLC	CPB-O7-P2-O22
3	A	390	MLC	CP7-CPA-CPB-O7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	390	MLC	CP9-CPA-CPB-O7
3	A	390	MLC	CP8-CPA-CPB-O7
3	A	390	MLC	CP2-CP1-S-CM1
3	A	390	MLC	CM2-CM1-S-CP1
3	A	390	MLC	OM2-CM1-S-CP1
4	A	391	PIN	C2-C1-S1-O1
4	A	391	PIN	C1-C2-N1-C3
4	A	391	PIN	C1'-C2'-N1'-C3'
3	A	390	MLC	OP1-CP3-NP1-CP2
3	A	390	MLC	C2'-C3'-O3'-P3
3	A	390	MLC	C4'-C3'-O3'-P3
3	A	390	MLC	CP3-CP4-CP5-NP2
3	A	390	MLC	O4'-C4'-C5'-O5'
4	A	391	PIN	C2-C1-S1-O3
3	A	390	MLC	OP1-CP3-CP4-CP5
3	A	390	MLC	C3'-C4'-C5'-O5'
4	A	391	PIN	C2-C1-S1-O2
4	A	391	PIN	S1-C1-C2-N1
3	A	390	MLC	C5'-O5'-P1-O11
3	A	390	MLC	CPB-O7-P2-O6
4	A	391	PIN	C1'-C2'-N1'-C4'
3	A	390	MLC	CP4-CP3-NP1-CP2
3	A	390	MLC	OP2-CP6-NP2-CP5
3	A	390	MLC	P2-O6-P1-O12
3	A	390	MLC	P1-O6-P2-O22


There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	390	MLC	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	389/389 (100%)	1.66	102 (26%) 1 1	6, 14, 26, 47	19 (4%)

All (102) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	389	ILE	4.8
1	A	1	MET	4.7
1	A	2	VAL	4.6
1	A	231	GLU	3.8
1	A	4	VAL	3.8
1	A	342	VAL	3.5
1	A	292	ILE	3.5
1	A	13	ALA	3.4
1	A	10	ALA	3.4
1	A	160	TYR	3.3
1	A	47	GLU	3.3
1	A	196	VAL	3.3
1	A	341	CYS	3.2
1	A	354[A]	THR	3.2
1	A	241	TRP	3.2
1	A	230	PRO	3.1
1	A	7	ILE	3.0
1	A	232	ILE	3.0
1	A	5[A]	SER	3.0
1	A	9[A]	LYS	3.0
1	A	284	VAL	2.9
1	A	356	ASN	2.9
1	A	240	VAL	2.9
1	A	295	TYR	2.9
1	A	277	LYS	2.9
1	A	265[A]	PHE	2.8
1	A	162	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	268	LEU	2.8
1	A	204	THR	2.8
1	A	48	HIS	2.8
1	A	102	VAL	2.8
1	A	229	VAL	2.8
1	A	142	TYR	2.7
1	A	246	ILE	2.6
1	A	12	ARG	2.6
1	A	3	SER	2.6
1	A	99	VAL	2.6
1	A	157	TYR	2.6
1	A	223	ILE	2.6
1	A	228	PRO	2.6
1	A	245	THR	2.5
1	A	39	PHE	2.5
1	A	62	LYS	2.5
1	A	210[A]	VAL	2.5
1	A	267	LEU	2.5
1	A	290	LEU	2.5
1	A	86	TYR	2.5
1	A	355	GLN	2.4
1	A	388	ALA	2.4
1	A	71	TYR	2.4
1	A	31	VAL	2.4
1	A	42[A]	LYS	2.4
1	A	222	LEU	2.4
1	A	237	PHE	2.4
1	A	359	LYS	2.3
1	A	117	TRP	2.3
1	A	78	LYS	2.3
1	A	17	ALA	2.3
1	A	8	ARG	2.3
1	A	183	GLY	2.3
1	A	318	ALA	2.3
1	A	20	LEU	2.3
1	A	15	GLY	2.3
1	A	193	VAL	2.3
1	A	37	PRO	2.2
1	A	201	PRO	2.2
1	A	289	PRO	2.2
1	A	56	PHE	2.2
1	A	225	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	138	PRO	2.2
1	A	370	LEU	2.2
1	A	14[A]	GLU	2.2
1	A	298	ILE	2.2
1	A	105	LEU	2.2
1	A	257	HIS	2.2
1	A	43	ILE	2.2
1	A	133	SER	2.2
1	A	58	ARG	2.1
1	A	177	LEU	2.1
1	A	367	TRP	2.1
1	A	384	LEU	2.1
1	A	82[A]	ASN	2.1
1	A	165	PHE	2.1
1	A	19	ILE	2.1
1	A	170	VAL	2.1
1	A	387	VAL	2.1
1	A	163	GLY	2.1
1	A	182	LYS	2.1
1	A	144	LEU	2.1
1	A	137	MET	2.1
1	A	38	ASP	2.1
1	A	235	PRO	2.1
1	A	287	PHE	2.1
1	A	113	ALA	2.1
1	A	45	ASN	2.1
1	A	279	ILE	2.1
1	A	376	GLY	2.0
1	A	110	ALA	2.0
1	A	164	ALA	2.0
1	A	361	THR	2.0
1	A	103	PRO	2.0
1	A	224	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MLC	A	390	54/54	0.59	0.33	20,41,50,51	54
4	PIN	A	391	18/18	0.78	0.20	14,30,39,39	18
2	SO4	A	392	5/5	0.82	0.17	27,28,28,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around PIN A 391:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.