



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:05 AM UTC

PDB ID : 8CMK / pdb_00008cmk
Title : Transportin-3 TNPO3 in complex with RSY region of CIRBP
Authors : Zhou, Q.; Sagmeister, T.; Pavkov-Keller, T.; Madl, T.
Deposited on : 2023-02-20
Resolution : 2.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1159 (2.96-2.92)
Clashscore	190562	1184 (2.96-2.92)
Ramachandran outliers	187476	1131 (2.96-2.92)
Sidechain outliers	187428	1131 (2.96-2.92)
RSRZ outliers	180081	1159 (2.96-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	923	
1	B	923	
2	C	35	
2	D	35	
2	E	35	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 29089 atoms, of which 14266 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transportin-3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	913	Total	C	H	N	O	S	0	0	0
			14304	4603	7075	1231	1342	53			
1	B	893	Total	C	H	N	O	S	0	0	0
			14011	4507	6932	1205	1314	53			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	511	ALA	CYS	engineered mutation	UNP Q9Y5L0
B	511	ALA	CYS	engineered mutation	UNP Q9Y5L0

- Molecule 2 is a protein called Cold-inducible RNA-binding protein.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	14	Total	C	H	N	O	0	0	0
			198	69	83	20	26			
2	D	14	Total	C	H	N	O	0	0	0
			198	69	83	20	26			
2	E	12	Total	C	H	N	O	0	0	0
			166	54	72	18	22			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			16	4	9	3		

- Molecule 4 is 2-(1H-INDOL-3-YL)ETHANAMINE (CCD ID: TSS) (formula: $C_{10}H_{12}N_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	N	0	0
			24	10	12	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	86	Total 86	O 86	0	0
5	B	73	Total 73	O 73	0	0
5	C	8	Total 8	O 8	0	0
5	D	5	Total 5	O 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transportin-3




- Molecule 1: Transportin-3




- Molecule 2: Cold-inducible RNA-binding protein






● Molecule 2: Cold-inducible RNA-binding protein



● Molecule 2: Cold-inducible RNA-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	97.53Å 101.81Å 114.13Å 90.00° 111.13° 90.00°	Depositor
Resolution (Å)	48.72 – 2.94 48.72 – 2.94	Depositor EDS
% Data completeness (in resolution range)	98.3 (48.72-2.94) 98.3 (48.72-2.94)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.8.0403, REFMAC 5.8.0403	Depositor
R, R_{free}	0.246 , 0.317 0.246 , 0.317	Depositor DCC
R_{free} test set	2107 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	66.9	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	29089	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.1067e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TSS, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/7374	1.06	3/10018 (0.0%)
1	B	0.56	0/7215	1.10	14/9792 (0.1%)
2	C	0.57	0/118	1.13	0/159
2	D	0.54	0/118	1.22	0/159
2	E	0.61	0/95	1.33	0/125
All	All	0.55	0/14920	1.09	17/20253 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	3
2	E	0	1
All	All	0	8

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	556	GLU	CB-CA-C	-8.98	95.89	110.79
1	A	395	ASP	CA-CB-CG	7.83	120.43	112.60
1	B	556	GLU	N-CA-CB	7.71	121.46	110.12
1	A	290	ASP	CB-CA-C	-7.14	98.54	110.68
1	B	394	ASP	CA-CB-CG	7.04	119.64	112.60
1	B	743	GLN	CB-CG-CD	-6.98	100.74	112.60
1	B	142	PHE	CA-CB-CG	-6.76	107.03	113.80
1	B	155	HIS	CA-CB-CG	6.35	120.15	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	753	PHE	CA-CB-CG	-5.43	108.36	113.80
1	B	81	ASP	CA-CB-CG	5.36	117.96	112.60
1	B	524	HIS	CA-CB-CG	-5.31	108.49	113.80
1	B	384	PRO	CA-C-N	5.27	127.60	120.38
1	B	384	PRO	C-N-CA	5.27	127.60	120.38
1	B	750	ASP	CA-CB-CG	5.24	117.84	112.60
1	A	607	ASP	CA-CB-CG	5.15	117.75	112.60
1	B	809	ASN	CA-CB-CG	5.04	117.64	112.60
1	B	394	ASP	CB-CA-C	-5.01	100.44	110.42

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	158	SER	Peptide
1	A	653	ARG	Sidechain
1	A	667	ARG	Sidechain
1	A	671	ARG	Sidechain
1	B	156	SER	Peptide
1	B	194	ASP	Peptide
1	B	671	ARG	Sidechain
2	E	161	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7229	7075	7257	26	3
1	B	7079	6932	7106	40	7
2	C	115	83	90	0	0
2	D	115	83	90	0	0
2	E	94	72	77	0	0
3	A	7	9	10	0	0
4	B	12	12	12	0	0
5	A	86	0	0	6	0
5	B	73	0	0	5	0
5	C	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	5	0	0	0	0
All	All	14823	14266	14642	64	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:94:ILE:HD12	1:B:142:PHE:CE1	1.97	0.97
1:B:94:ILE:HD12	1:B:142:PHE:CZ	2.13	0.83
1:A:730:ILE:HB	1:A:731:PRO:HD3	1.60	0.82
1:B:541:LEU:HB3	1:B:565:THR:HG22	1.71	0.71
1:B:599:GLU:OE2	5:B:1101:HOH:O	2.11	0.69
1:B:549:ASP:OD1	1:B:614:ARG:NH1	2.30	0.64
1:B:367:ALA:HB2	5:B:1141:HOH:O	1.98	0.63
1:A:54:ARG:HG2	1:A:97:LEU:HD21	1.81	0.62
1:A:373:LEU:CD2	5:A:1170:HOH:O	2.46	0.62
1:A:230:GLU:HG3	5:A:1111:HOH:O	2.00	0.60
1:A:513:LYS:O	5:A:1101:HOH:O	2.16	0.60
1:A:54:ARG:HD3	1:A:96:ASN:HB3	1.83	0.60
1:B:536:GLN:HG3	5:B:1132:HOH:O	2.03	0.57
1:B:589:VAL:HG13	1:B:644:VAL:HG11	1.87	0.57
1:B:94:ILE:HB	1:B:142:PHE:CE2	2.40	0.56
1:A:541:LEU:HB3	1:A:565:THR:HG22	1.89	0.55
1:A:174:ALA:HA	1:A:213:VAL:HB	1.89	0.53
1:B:94:ILE:HG22	1:B:108:LEU:HD13	1.90	0.53
1:B:277:GLU:OE1	1:B:322:ARG:NH1	2.39	0.53
1:A:373:LEU:HD21	5:A:1170:HOH:O	2.08	0.53
1:B:839:HIS:HD2	5:B:1136:HOH:O	1.93	0.51
1:A:424:TYR:CE2	1:A:428:LYS:HE3	2.46	0.51
1:B:673:VAL:O	1:B:673:VAL:HG12	2.12	0.50
1:B:101:SER:O	1:B:104:ILE:HG13	2.12	0.50
1:B:94:ILE:HD12	1:B:142:PHE:CD1	2.44	0.49
1:B:576:LYS:HE2	1:B:580:CYS:SG	2.52	0.48
1:B:641:ILE:O	1:B:644:VAL:HG12	2.13	0.48
1:A:753:PHE:CD2	1:A:795:VAL:HG22	2.48	0.48
1:B:803:ILE:HD13	1:B:860:ILE:CG1	2.44	0.48
1:B:87:ARG:HB2	1:B:122:TRP:CH2	2.49	0.47
1:B:753:PHE:CD2	1:B:795:VAL:HG22	2.50	0.47
1:A:384:PRO:HB3	5:A:1119:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:905:ALA:HB1	1:B:910:GLN:HB2	1.95	0.46
1:A:599:GLU:N	1:A:600:PRO:CD	2.78	0.46
1:A:373:LEU:HD22	5:A:1170:HOH:O	2.15	0.46
1:B:394:ASP:HB3	1:B:397:GLY:H	1.80	0.46
1:B:561:LEU:O	1:B:565:THR:HG23	2.15	0.46
1:A:87:ARG:HD2	1:A:122:TRP:CD2	2.51	0.46
1:B:839:HIS:CD2	5:B:1136:HOH:O	2.69	0.46
1:A:358:ASP:OD1	1:A:358:ASP:C	2.60	0.45
1:A:151:PRO:HB3	1:A:211:LEU:HD22	1.97	0.45
1:B:153:GLU:HA	1:B:156:SER:HB3	1.98	0.45
1:B:303:CYS:SG	1:B:323:THR:HB	2.57	0.44
1:A:424:TYR:CE2	1:A:428:LYS:CE	3.00	0.44
1:B:219:MET:HB2	1:B:265:LEU:HD13	2.00	0.43
1:A:561:LEU:O	1:A:565:THR:HG23	2.18	0.43
1:B:546:ARG:HA	1:B:587:VAL:HG21	2.01	0.43
1:B:411:ILE:HG23	1:B:419:CYS:SG	2.58	0.43
1:B:133:TYR:HB3	1:B:139:SER:HB2	1.99	0.43
1:A:75:PHE:CZ	1:A:83:HIS:CE1	3.07	0.43
1:B:321:LEU:HD12	1:B:324:LEU:HD12	2.01	0.43
1:B:500:ASP:HB3	1:B:501:PRO:HD3	2.00	0.42
1:B:424:TYR:CD2	1:B:428:LYS:HE2	2.55	0.42
1:A:667:ARG:HH21	1:A:667:ARG:HG3	1.83	0.42
1:A:157:ARG:NH2	1:B:917:ASP:OD2	2.52	0.42
1:A:456:ASN:N	1:A:457:PRO:HD2	2.35	0.42
1:B:191:ALA:C	1:B:193:THR:H	2.28	0.42
1:B:730:ILE:HB	1:B:731:PRO:HD3	2.01	0.41
1:A:894:GLN:NE2	1:B:155:HIS:O	2.53	0.41
1:A:424:TYR:CG	1:A:462:VAL:HG22	2.55	0.41
1:B:552:LEU:HD13	1:B:552:LEU:HA	1.83	0.41
1:B:400:ARG:HH12	1:B:438:GLU:CD	2.30	0.40
1:A:427:LEU:HD21	1:A:440:VAL:HB	2.03	0.40
1:B:630:GLN:OE1	1:B:630:GLN:HA	2.21	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:263:LEU:HD23	1:B:549:ASP:CG[2_656]	0.96	0.64
1:A:45:GLU:OE2	1:B:54:ARG:CB[2_556]	1.86	0.34
1:B:263:LEU:CD2	1:B:549:ASP:CG[2_656]	1.88	0.32

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:271:GLN:OE1	1:B:598:GLN:NE2[2_656]	1.95	0.25
1:B:263:LEU:HD23	1:B:549:ASP:OD2[2_656]	1.45	0.15
1:A:45:GLU:OE2	1:B:54:ARG:HB2[2_556]	1.46	0.14
1:A:45:GLU:CD	1:B:54:ARG:HB2[2_556]	1.55	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	909/923 (98%)	895 (98%)	14 (2%)	0	100	100
1	B	881/923 (95%)	849 (96%)	30 (3%)	2 (0%)	43	65
2	C	12/35 (34%)	12 (100%)	0	0	100	100
2	D	12/35 (34%)	12 (100%)	0	0	100	100
2	E	10/35 (29%)	10 (100%)	0	0	100	100
All	All	1824/1951 (94%)	1778 (98%)	44 (2%)	2 (0%)	48	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	120	PRO
1	B	119	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	812/820 (99%)	810 (100%)	2 (0%)	87	95
1	B	796/820 (97%)	791 (99%)	5 (1%)	78	89
2	C	12/30 (40%)	12 (100%)	0	100	100
2	D	12/30 (40%)	12 (100%)	0	100	100
2	E	10/30 (33%)	9 (90%)	1 (10%)	7	19
All	All	1642/1730 (95%)	1634 (100%)	8 (0%)	81	90

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	596	LEU
1	A	828	GLN
1	B	94	ILE
1	B	213	VAL
1	B	258	ASN
1	B	475	THR
1	B	627	GLU
2	E	161	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	HIS
1	A	96	ASN
1	A	118	GLN
1	A	155	HIS
1	A	268	GLN
1	A	294	ASN
1	A	455	ASN
1	A	456	ASN
1	A	495	ASN
1	A	539	ASN
1	A	588	GLN
1	A	598	GLN
1	A	652	HIS
1	A	695	HIS
1	A	745	HIS
1	A	804	HIS
1	A	828	GLN
1	B	13	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	49	GLN
1	B	155	HIS
1	B	234	GLN
1	B	271	GLN
1	B	525	ASN
1	B	635	GLN
1	B	745	HIS
1	B	828	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TSS	B	1001	-	13,13,13	0.42	0	17,17,17	0.53	0
3	PEG	A	1001	-	6,6,6	0.34	0	5,5,5	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TSS	B	1001	-	-	2/3/3/3	0/2/2/2
3	PEG	A	1001	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1001	TSS	CA-CB-CG-CD1
4	B	1001	TSS	CA-CB-CG-CD2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	913/923 (98%)	0.09	17 (1%) 66 58	41, 74, 116, 150	0
1	B	893/923 (96%)	0.36	35 (3%) 43 35	39, 83, 171, 235	0
2	C	14/35 (40%)	0.03	1 (7%) 22 19	43, 60, 84, 88	0
2	D	14/35 (40%)	0.20	0 100 100	41, 60, 80, 91	0
2	E	12/35 (34%)	0.47	1 (8%) 17 15	102, 123, 139, 144	0
All	All	1846/1951 (94%)	0.22	54 (2%) 53 44	39, 78, 151, 235	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	264	PRO	3.6
1	B	153	GLU	3.4
1	B	210	ASN	3.3
1	B	204	CYS	3.2
1	A	453	PRO	3.0
1	A	813	GLU	3.0
1	B	150	LEU	3.0
1	B	80	THR	2.9
1	B	173	LEU	2.9
1	B	208	TRP	2.9
1	B	152	GLU	2.9
1	B	141	PRO	2.8
1	B	541	LEU	2.8
1	B	457	PRO	2.7
1	B	154	VAL	2.7
1	B	255	ALA	2.7
1	B	159	LEU	2.6
1	B	18	LEU	2.6
1	A	560	GLY	2.6
1	B	314	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	214	LEU	2.5
1	A	443	ILE	2.5
1	B	502	VAL	2.4
1	B	888	VAL	2.4
1	A	857	LEU	2.4
1	A	416	SER	2.3
1	B	328	LEU	2.3
2	E	164	TYR	2.3
1	B	441	LEU	2.3
1	B	681	LEU	2.2
1	B	71	ILE	2.2
1	B	62	PHE	2.2
1	A	888	VAL	2.2
1	A	328	LEU	2.2
1	A	850	LEU	2.2
1	B	850	LEU	2.2
1	A	20	HIS	2.2
1	B	130	VAL	2.2
1	B	92	THR	2.2
1	B	420	PHE	2.2
1	A	890	VAL	2.1
1	B	811	HIS	2.1
2	C	171	ASN	2.1
1	A	889	THR	2.1
1	A	603	GLY	2.1
1	B	356	THR	2.1
1	B	15	VAL	2.1
1	B	147	LEU	2.1
1	A	492	VAL	2.0
1	A	807	VAL	2.0
1	B	902	VAL	2.0
1	A	629	GLY	2.0
1	A	455	ASN	2.0
1	B	212	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	A	1001	7/7	0.69	0.14	63,73,77,78	0
4	TSS	B	1001	12/12	0.77	0.15	68,74,78,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.